

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 62 (2016)
Heft: 1-2

Artikel: Euler equations on the general linear group, cubic curves, and inscribed hexagons
Autor: Aleshkin, Konstantin / Izosimov, Anton
DOI: <https://doi.org/10.5169/seals-685359>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Euler equations on the general linear group, cubic curves, and inscribed hexagons

Konstantin ALESHKIN and Anton IZOSIMOV

Abstract. We study integrable Euler equations on the Lie algebra $\mathfrak{gl}(3, \mathbb{R})$ by interpreting them as evolutions on the space of hexagons inscribed in a real cubic curve.

Mathematics Subject Classification (2010). Primary: 37J35; Secondary: 14H70.

Keywords. Integrable systems, Euler equations on Lie algebras, cubic curves.

1. Introduction

In this paper we study an integrable matrix differential equation

$$(1.1) \quad \frac{d}{dt}X = [X^2, A]$$

where $X \in \mathfrak{gl}(3, \mathbb{R})$ is a real 3 by 3 matrix depending on the time t , and $A \in \mathfrak{gl}(3, \mathbb{R})$ is a real, fixed, 3 by 3 matrix. While we will not need a precise definition of an integrable system, we will take the point of view of [HSW] according to which “*integrability of a system of differential equations should manifest itself through some generally recognizable features: i) the existence of many conserved quantities, ii) the presence of algebraic geometry, iii) the ability to give explicit solutions.*” Equation (1.1) shows all these properties. In particular, the algebraic geometry underlying this equation is geometry of real cubic curves. This algebraic geometry arises from the possibility to rewrite equation (1.1) in the so-called *Lax form with a spectral parameter* (see equation (2.1) below). A Lax representation with a spectral parameter for equations of type (1.1) was found in S.V.Manakov’s fundamental paper [Man].

Equation (1.1) can be regarded as a special case of several general constructions of integrable systems. In particular, it can be obtained by the argument shift method [Man, MF], or by the method based on loop algebras [AvM2, AvM1, RSTS1, RSTS2]. Depending on the restrictions imposed on the matrices X and

A , this equation is known under different names. If A is symmetric, and X is skew-symmetric, it becomes the classical Euler equation describing the rotation of a torque-free rigid body. If, on the contrary, A is skew-symmetric, and X is symmetric, this equation is known as the Bloch–Iserles system [BI, BBI⁺]. Finally, it is worth mentioning the case of skew-Hermitian X and Hermitian A . In this setting, Equation (1.1) describes travelling wave solutions for the famous three-wave equation arising in optics, fluid dynamics, and plasma physics [ALMR] (see also recent paper [RT] studying equations of type (1.1) for $n \times n$ skew-Hermitian matrices).

In the present paper, we regard equation (1.1) as a dynamical system on the whole space $\mathfrak{gl}(3, \mathbb{R})$ of real 3 by 3 matrices. A distinctive feature of this full system is that, in contrast to the symmetric and skew-symmetric cases, solutions on arbitrary matrices need not be bounded and, in particular, may blow up in finite time. Using algebro-geometric technique, we show that for a generic matrix A Equation (1.1) has both types of solutions, that is blow-up solutions, and solutions defined for any $t \in \mathbb{R}$. We also show that the behavior of a given solution can be understood in terms of a simple geometric construction. Namely, with each generic initial condition X we associate a real cubic curve C_X with fixed points at infinity (the *spectral curve* coming from the Lax representation), and a hexagon H_X inscribed in this curve in such a way that its sides are parallel to the asymptotes of the curve. Then, we show that the behavior of the solution of Equation (1.1) with initial condition X is completely determined by the number of ovals of the curve C_X and the distribution of vertices of the hexagon H_X among these ovals.

It is also worth mentioning that Equation (1.1) may be regarded as an Euler equation on the Lie algebra $\mathfrak{gl}(3, \mathbb{R})$, or, which is the same, the geodesic equation for a certain left-invariant metric on the general linear group $GL(3, \mathbb{R})$. The study of such metrics originates from V. Arnold’s fundamental paper [Arn1], where Arnold suggests a common geometric framework for the Euler equation governing the motion of an ideal fluid, and the Euler equation in rigid body dynamics. In Arnold’s approach, both equations describe the geodesic flow of a one-sided invariant metric on a certain Lie group G . Such a geodesic flow is a dynamical system on the cotangent bundle T^*G , and, thanks to the G -invariance, it descends to the quotient space T^*G / G , which is naturally identified with the dual Lie algebra $\mathfrak{g}^*$. The corresponding equation on $\mathfrak{g}^*$ is called an Euler equation.

Equation (1.1) is an example of an Euler equation on the Lie algebra $\mathfrak{gl}(3, \mathbb{R})$. It describes the geodesic flow of a left-invariant pseudo-Riemannian metric $(\cdot, \cdot)$ on the group $GL(3, \mathbb{R})$ given at the identity by

$$(X, X) := \operatorname{tr} X \mathcal{A}^{-1}(X)$$

where $\mathcal{A}(X) = \frac{1}{2}(AX + XA)$. In particular, the problem of existence of global solutions for equation (1.1) is equivalent to the problem of geodesic completeness for the metric $(\cdot, \cdot)$.

A distinctive feature of the Euler equation (1.1) is its integrability. Note that a general Euler equation need not be integrable, and integrable examples are in fact quite rare. In particular, the above equation seems to be the only known example of an integrable Euler equation on $\mathfrak{gl}(3, \mathbb{R})$.

The problem of geodesic completeness for left-invariant metrics on finite-dimensional Lie groups was studied, for example, in [AP, BM]. Note that for general invariant metrics, geodesic completeness or, equivalently, existence of global in time solutions of the Euler equation, seems to be a very difficult problem. For integrable metrics constructed by the argument shift method this problem was discussed in [BIKO].

We also remark that since equation (1.1) is integrable, its solutions can be explicitly expressed in terms of theta functions. So, global behavior of solutions can be, in principle, studied by finding and examining explicit formulas. However, as we show in the present paper, global properties of solutions can be in fact understood from purely geometrical considerations, and there is no need in the analysis of complicated theta-functional formulas.

We tried to make the exposition self-contained. In particular, we do not assume that the reader is familiar with the general theory of integrable systems and the algebro-geometric approach to such systems. For most statements which can be, in principle, derived from this general theory, we give geometric proofs (relations to the general theory are explained in remarks; see, in particular, Remark 2.18). The only exception is, perhaps, Proposition 3.4 where we follow the standard approach on linearization of an integrable flow on the Jacobian. It would be interesting to find a geometric proof for this statement as well.

Main results of the paper are in Section 2. Section 3 is devoted to proofs of these results. In Section 4, we discuss possible generalizations of our approach to the $\mathfrak{gl}(n)$ case and their relation to general questions of real algebraic geometry.

2. Main constructions and results

2.1. Reduction to diagonal matrices. In what follows, we assume that the eigenvalues of the matrix A are all distinct and real. The case of complex conjugate eigenvalues can be treated using similar ideas, but still needs a separate consideration, and we omit it.

Denote the eigenvalues of A by a_1, a_2, a_3 . Note that equation (1.1) is invariant under similarity transformations

$$X \mapsto BXB^{-1}, \quad A \mapsto BAB^{-1}.$$

For this reason, we may assume that A is a diagonal matrix with diagonal entries a_1, a_2, a_3 . Therefore, 9-dimensional family of equations (1.1) boils down to a 3-dimensional family parametrized by three real numbers a_1, a_2, a_3 .

So, in what follows, we always assume that A is diagonal with distinct diagonal entries. We call such diagonal matrices *generic*.

Provided that A is diagonal, equation (1.1) is invariant under transformations of the form $X \mapsto DXD^{-1}$ where D is an invertible diagonal matrix. Such transformations form a group which may be regarded as the quotient group of invertible diagonal matrices by scalar matrices. We shall denote this quotient group by $\mathbb{PD}(3, \mathbb{R})$. This group is isomorphic to $(\mathbb{R}^*)^2$, and in particular, it is disconnected. We denote its connected component of the identity by $\mathbb{PD}^+(3, \mathbb{R})$. The latter group consists of (cosets of) those diagonal matrices whose diagonal entries are of the same sign.

2.2. Lax representation and spectral curve. We begin our study of equation (1.1) by rewriting it as a so-called *Lax equation*

$$(2.1) \quad \frac{d}{dt} X_\lambda = [X_\lambda, Y_\lambda]$$

where

$$(2.2) \quad X_\lambda := X + \lambda A, \quad Y_\lambda := AX + XA + \lambda A^2$$

and $\lambda \in \mathbb{C}$ is an auxiliary time-independent parameter, called the *spectral parameter*. It is straightforward to verify that equations (1.1) and (2.1) are equivalent.

Remark 2.1. For details about Lax equations with a spectral parameter and their algebraic-geometric solutions see, e.g., the monograph [BBT].

The following proposition is well-known.

Proposition 2.2. *If a matrix X_λ evolves according to equation (2.1), then the eigenvalues of X_λ do not change with time.*

Proof. Using induction on k , one can show that

$$\frac{d}{dt} X_\lambda^k = [X_\lambda^k, Y_\lambda]$$

for any integer $k \geq 1$; therefore

$$\frac{d}{dt} \operatorname{tr} X_\lambda^k = \operatorname{tr} [X_\lambda^k, Y_\lambda] = 0$$

where in the last identity we used that the trace of a commutator is always equal to zero. Thus, since traces of powers of the matrix X_λ do not depend on t , neither do its eigenvalues, q.e.d. $\square$

Proposition 2.2 implies that the coefficients of the characteristic polynomial $f_X(\lambda, \mu) := \det(X + \lambda A - \mu \operatorname{Id})$ are conserved along the solutions of equation (1.1). Note that only six out of ten coefficients explicitly depend on X , so there are six conserved quantities. We will not need explicit expressions for these conserved quantities. Instead, we organize them into an algebraic curve, called the spectral curve. In affine coordinates, this curve is defined by the equation $f_X(\lambda, \mu) = 0$. However, it will be convenient for us to work in homogenous coordinates. For this reason, we give the following definition:

Definition 2.3. For a given $X \in \mathfrak{gl}(3, \mathbb{R})$, the curve

$$C_X := \{(z_1 : z_2 : z_3) \in \mathbb{CP}^2 \mid \det(z_3 X + z_1 A - z_2 \operatorname{Id}) = 0\}$$

is called the *spectral curve*.

By definition, the spectral curve C_X is conserved along solutions of equation (1.1).

Proposition 2.4. *The spectral curve C_X is a real¹ projective cubic intersecting the line at infinity $\{z_3 = 0\}$ at points*

$$\infty_1 = (1 : a_1 : 0), \quad \infty_2 = (1 : a_2 : 0), \quad \infty_3 = (1 : a_3 : 0)$$

where a_1, a_2, a_3 are the eigenvalues of A .

Proof. The proof is achieved by putting $z_3 = 0$ in the equation of the spectral curve. $\square$

As we show below, any smooth real cubic curve passing through the points $\infty_1, \infty_2, \infty_3$ is the spectral curve for a suitable matrix $X \in \mathfrak{gl}(3, \mathbb{R})$. Moreover, we explicitly describe the topology of the set of matrices X corresponding to the given curve C in terms of the geometry of C .

¹ Recall that an algebraic curve is called real if it is invariant under complex conjugation, or, equivalently, if it can be defined as the zero locus of a real polynomial.

2.3. Isospectral sets. Since the flow (1.1) preserves the spectral curve associated with the matrix X , we can restrict this flow to the set

$$\mathcal{T}_C = \{X \in \mathfrak{gl}(3, \mathbb{R}) : C_X = C\}$$

of matrices whose spectral curve is the given curve C . Note that the set $\mathcal{T}_C$ may also be defined as a joint level set for six conserved quantities of equation (1.1) (recall that these conserved quantities are, by definition, the coefficients of the equation of the spectral curve). Since the space $\mathfrak{gl}(3, \mathbb{R})$ is 9-dimensional, we should expect that the set $\mathcal{T}_C$ is generically of dimension $9 - 6 = 3$.

Further, note that the flow (1.1) restricted to the 3-dimensional manifold $\mathcal{T}_C$ has a 2-dimensional symmetry group $\mathbb{P}D^+(3, \mathbb{R})$ acting by conjugation. A dimension count suggests that the $\mathbb{P}D^+(3, \mathbb{R})$ orbits of solutions of equation (1.1) are exactly the connected components of $\mathcal{T}_C$. In particular, all solutions lying in the same connected component should have the same global behavior.

In what follows, we aim to answer the following questions.

- i) For a given cubic curve C , what is the topology of the set $\mathcal{T}_C$? In particular, how many connected components does it have?
- ii) What is the global behavior of solutions of (1.1) on each of these components? In particular, do these solutions blow up or exist for all times?
- iii) Given an initial condition $X \in \mathfrak{gl}(3, \mathbb{R})$, how do we determine whether the solution passing through X blows up, or exists for all times?

The answer to the first two of these questions is given by Theorem 2.5. The answer to the third question is given by Theorem 2.13.




FIGURE 1

Types of cubic curves and the topology of the corresponding isospectral sets

2.4. Topology of isospectral sets. The answer to the above questions i) and ii) is formulated in terms of the real part of the curve C . By definition, the real part $C_{\mathbb{R}}$ of a real projective curve C is the set of its real points: $C_{\mathbb{R}} = C \cap \mathbb{RP}^2$. If the curve C is smooth, then its real part consists of a finite number of closed curves, which are called ovals. An oval is called bounded if it does not intersect the line at infinity. Otherwise, it is unbounded. It is a classical result that a smooth projective cubic can have either one oval, which is then unbounded, or two ovals, at least one of which is unbounded (see Figure 1).

Before we describe the set $\mathcal{T}_C$, note that for this set to be non-empty, the curve C should have the properties listed in Proposition 2.4, i.e. it should be a real cubic passing through the points ∞_1 , ∞_2 , and ∞_3 where $\infty_i = (1 : a_i : 0)$, and a_1, a_2, a_3 are the eigenvalues of the matrix A . The following theorem in particular says that for smooth curves C these conditions are also sufficient for the set $\mathcal{T}_C$ to be non-empty.

Theorem 2.5. *Assume that A is a generic diagonal matrix, and let C be a smooth real cubic passing through the points ∞_1 , ∞_2 , and ∞_3 . Then the following statements hold.*

- (1) *If the real part of C has one oval, then the set $\mathcal{T}_C$ has four connected components each diffeomorphic to $\mathbb{R}^3$.*
- (2) *If the real part of C has two ovals and one of them is bounded, then $\mathcal{T}_C$ has four components diffeomorphic to $\mathbb{R}^3$ and four components diffeomorphic to $S^1 \times \mathbb{R}^2$.*
- (3) *Finally, if the real part of C has two unbounded ovals, then $\mathcal{T}_C$ has four components diffeomorphic to $\mathbb{R}^3$ and two components diffeomorphic to $S^1 \times \mathbb{R}^2$.*

Furthermore, all solutions of (1.1) lying on components of $\mathcal{T}_C$ diffeomorphic to $\mathbb{R}^3$ blow up², while all solutions lying on $S^1 \times \mathbb{R}^2$ components exist for all times.

Note that this theorem does not answer the third of the above questions. Namely, if the spectral curve C_X has two ovals, then Theorem 2.5 does not allow us to determine whether the solution with initial condition X blows up or exists for all times. As we discuss below, the answer to this question can also be given in terms of a simple geometric construction.

²In what follows, when we say that a solution blows up, we mean that it does so both forward and backward in time. Note that equation (1.1) does have solutions which blow up only in one direction, but these solutions correspond to singular spectral curves.

Remark 2.6. Note that despite the fact that solutions located on $\mathbb{R}^3$ components blow up, the topology of these components is still compatible with the Arnold–Liouville theorem [Arn2]. This phenomenon is explained in the first author’s paper [Ale].

2.5. Regularly inscribed hexagons. As was pointed out above, it is not in general possible to decide from the spectral curve C_X whether the solution of equation (1.1) with initial condition X blows up. So, we need to supplement the curve with certain additional data in order to be able to understand the behavior of a given solution. It turns out that as such additional data we can take a certain hexagon inscribed in the spectral curve. This hexagon is constructed as follows.

As before, we assume that A is a diagonal matrix with distinct diagonal entries a_1, a_2, a_3 . Under this assumption, the spectral curve C_X has three distinct real asymptotes which are, by definition, the tangent lines to C_X at the points $\infty_1, \infty_2, \infty_3$. Denote these asymptotes by l_1, l_2, l_3 . Let $X \in \mathfrak{gl}(3, \mathbb{R})$ be such that the spectral curve C_X is smooth. Consider the matrix

$$X_z := z_3 X + z_1 A - z_2 \text{Id}$$

$$= \begin{pmatrix} x_{11}z_3 + a_1z_1 - z_2 & x_{12}z_3 & x_{13}z_3 \\ x_{21}z_3 & x_{22}z_3 + a_2z_1 - z_2 & x_{23}z_3 \\ x_{31}z_3 & x_{32}z_3 & x_{33}z_3 + a_3z_1 - z_2 \end{pmatrix}.$$

Recall that the zero locus of the determinant of this matrix is, by definition, the spectral curve C_X . Furthermore, we have the following:

Proposition 2.7. *Asymptotes of the curve C_X are the zero loci of the diagonal entries of the matrix X_z . In other words, the equation of the asymptote l_i is $L_i = 0$ where*

$$L_i := x_{ii}z_3 + a_i z_1 - z_2.$$

Proof. We have

$$\det X_z = L_1 L_2 L_3 + z_3^2 L$$

where L is a linear function in z_1, z_2, z_3 . Therefore, the restriction of the function L_i to the curve $\det X_z = 0$ has a zero of order 2 at infinity, which means that $L_i = 0$ is an asymptote. $\square$

Further, let $M_{ij}(z_1, z_2, z_3)$ be the (i, j) minor of the matrix X_z . Then, for $i \neq j$, we have $M_{ij} = z_3 L_{ij}$ where L_{ij} is a linear function in z_1, z_2, z_3 . Explicitly, one has

$$(2.3) \quad L_{ij} = \pm \det \begin{pmatrix} x_{ki} & x_{kk}z_3 + a_k z_1 - z_2 \\ x_{ji} & x_{jk}z_3 \end{pmatrix}$$




FIGURE 2
Inscribed hexagon H_X

where (i, j, k) is any permutation of $(1, 2, 3)$. Note that the function L_{ij} cannot be identically equal to zero. Indeed, as follows from formula (2.3), if $L_{ij} \equiv 0$, then either the j 'th row or the i 'th column of the matrix X_z contains two zeros. The latter implies that the polynomial $\det X_z$ is reducible, which contradicts the smoothness of the curve C_X .

This way, we obtain six straight lines l_{ij} given by $L_{ij} = 0$ where $i \neq j$. Properties of these straight lines are described in the following proposition.

Proposition 2.8. *Let (i, j, k) be any permutation of $(1, 2, 3)$. Then:*

- (1) *The line l_{ij} is parallel to the asymptote l_k of the curve C_X . In other words, we have*

$$l_{12} \parallel l_{21} \parallel l_3, \quad l_{23} \parallel l_{32} \parallel l_1, \quad l_{31} \parallel l_{13} \parallel l_2.$$

- (2) *We have $l_{ij} \neq l_{ik}$, and $l_{ji} \neq l_{ki}$.*

- (3) *The points*

$$P_i := l_{ij} \cap l_{ik}, \quad Q_i := l_{ji} \cap l_{ki}$$

lie in the real part of the curve C_X . In other words, $P_1 Q_2 P_3 Q_1 P_2 Q_3$ is an inscribed hexagon (see Figure 2).

Proof. The first statement is straightforward and follows from Proposition 2.7 and formula (2.3) for the function L_{ij} . Let us prove the second statement. Assume that $l_{ij} = l_{ik} = l$. Then, since $l_{ij} \parallel l_k$ and $l_{ik} \parallel l_j$, we have $\infty_j, \infty_k \in l$. Therefore, l is the line at infinity. At the same time, it is easy to see from formula (2.3) that the line l_{ij} is the line at infinity if and only if $x_{ji} = 0$. So, we have $x_{ji} = x_{ki} = 0$. However, if this was so, then the curve C_X would be not smooth

but reducible. Therefore, our assumption is false, and $l_{ij} \neq l_{ik}$. The proof of the inequality $l_{ji} \neq l_{ki}$ is analogous.

Now, let us prove the third statement. We show that the point P_1 lies in the real part of C_X . The proof for other points is analogous. First note that P_1 is the intersection point of two real straight lines, therefore this point is real. So, it suffices to show that $P_1 \in C_X$. Let $P_1 = (z_1 : z_2 : z_3)$. Then, by definition of the point P_1 , the first two columns, as well as the first and the last column of the matrix

$$(2.4) \quad \begin{pmatrix} x_{21} & x_{22}z_3 + a_2z_1 - z_2 & x_{23}z_3 \\ x_{31} & x_{32}z_3 & x_{33}z_3 + a_3z_1 - z_2 \end{pmatrix}$$

are linearly dependent. Note that the first column of this matrix cannot be zero: if it is zero, then, again, C_X is a reducible curve. Therefore, the rank of this matrix is equal to one, which implies that $\det X_z = 0$, and thus $P_1 \in C_X$. $\square$

Remark 2.9. Note that since the rank of the matrix (2.4) is equal to 1 at the point P_1 , the diagonal minor M_{11} of the matrix X_z at P_1 is equal to zero. Similarly, M_{11} vanishes at the point Q_1 . Also note that the zero locus of the minor M_{11} is a conic whose asymptotes coincide with the asymptotes l_2, l_3 of the spectral curve. The latter implies that the conic $M_{11} = 0$ has at most two finite intersection points with the spectral curve, and these points are P_1 and Q_1 . Similarly, P_i and Q_i may be defined as finite intersection points of the quadric $M_{ii} = 0$ with the spectral curve.

Proposition 2.8 implies that to each matrix X such that the corresponding spectral curve C_X is smooth one can associate a hexagon inscribed in the real part of the spectral curve. We denote this hexagon by H_X . The sides of this hexagon are parallel to the asymptotes of the curve C_X . In what follows, hexagons with this property are called *regularly inscribed hexagons*. More precisely, we give the following definition:

Definition 2.10. Assume that C is a real smooth cubic curve which intersects the line at infinity at real points ∞_1, ∞_2 , and ∞_3 . A hexagon *regularly inscribed* in C is six points $P_1, Q_2, P_3, Q_1, P_2, Q_3 \in C_{\mathbb{R}}$ such that for any permutation (i, j, k) of $(1, 2, 3)$, the third intersection point of the line $P_i Q_j$ with the curve C is ∞_k .

Note that a regularly inscribed hexagon is uniquely determined by any of its vertices. Indeed, assume that we are given a point $P_1 \in C_{\mathbb{R}}$. Then we can reconstruct the point Q_2 as the intersection point of the cubic with the line passing through P_1 and parallel to the asymptote l_3 . In a similar way, we reconstruct

points P_3 , Q_1 , P_2 , and Q_3 . If it turns out that Q_3P_1 is parallel to the asymptote l_2 , then we obtain a regularly inscribed hexagon. In what follows, we show that this is always so. This, in fact, is a simple corollary of a classical result about nine points on a cubic, known as Chasles, or Cayley–Bacharach theorem. Thus, there exists exactly one regularly inscribed hexagon with a given vertex P_1 (this is, of course, true for other vertices as well). In particular, the set of hexagons regularly inscribed in a curve C can be, in principle, identified with the real part of C .

Also note that Definition 2.10 describes a slightly more general class of hexagons compared to Proposition 2.8. Indeed, setting $P_i = Q_i = \infty_i$ for $i = 1, 2, 3$, we obtain a regularly inscribed hexagon. In what follows, we shall refer to this hexagon as *the degenerate hexagon* (note that if $P_i = \infty_i$ or $Q_i = \infty_i$ at least for one value of i , then the hexagon is automatically degenerate). Since all sides of the degenerate hexagon coincide, the second statement of Proposition 2.8 implies that this hexagon does not correspond to any matrix X . As we show below, this situation is exceptional: any other regularly inscribed hexagon corresponds to a 2-dimensional family of matrices X .

Now, we need to discuss certain topological properties of regularly inscribed hexagons. Note that if the real part of C has two ovals, then different vertices of a regularly inscribed hexagon may lie on different ovals. To distinguish between possible configurations, we use the fact that if the real part of a cubic has two ovals, then exactly one of these two ovals is contractible in $\mathbb{RP}^2$. Namely, if one of the ovals is bounded, then it is contractible, and the other oval is not; if both ovals are unbounded, then the one which intersects the line at infinity at two points is contractible, and the other one is not.

Definition 2.11. We say that a regularly inscribed hexagon has *type* (m, n) if m of its vertices lie on the contractible oval, and n of its vertices lie on the non-contractible oval.

Proposition 2.12. *All possible types of regularly inscribed hexagons are depicted in Figure 3: if one of the ovals of $C_{\mathbb{R}}$ is bounded, then H has type $(0, 6)$ or $(6, 0)$, and if both ovals of $C_{\mathbb{R}}$ are unbounded, then H has type $(4, 2)$ or $(2, 4)$.*

Proof. The proof follows from simple topological considerations. □

2.6. Blow-up and global solutions. Now, we formulate a theorem which allows one to determine whether a given solution of equation (1.1) blows up. This result is stated in terms of the type of the hexagon H_X .

First note that, in contrast to the spectral curve C_X , the hexagon H_X is time-dependent. However, the type of this hexagon obviously cannot change with




FIGURE 3

Different types of regularly inscribed hexagons
and the behavior of the corresponding solutions

time. Moreover, the type of H_X stays the same if we vary X within a connected component of the isospectral set $\mathcal{T}_C$. It turns out that the topological type of H_X allows one to distinguish between $\mathbb{R}^3$ and $S^1 \times \mathbb{R}^2$ components of $\mathcal{T}_C$, i.e. between global and blow-up solutions.

Theorem 2.13. *Assume that A is a generic diagonal matrix, and let $X \in \mathfrak{gl}(3, \mathbb{R})$ be such that the spectral curve C_X is smooth. Let also $X(t)$ be the solution of equation (1.1) with initial condition X . Then the following statements hold.*

- (1) *If the real part of C_X has one oval, then $X(t)$ blows up.*
- (2) *If the real part of C_X has two ovals, then $X(t)$ blows up if and only if the hexagon H_X is of type $(0, 6)$ or $(4, 2)$; if H_X is of type $(6, 0)$ or $(2, 4)$, then $X(t)$ exists for all times.*

Example 2.14 (Rigid body). As we mentioned in the Introduction, for skew-symmetric matrices X equation (1.1) becomes the Euler equation governing the motion of a rigid body. Let us demonstrate how Theorem 2.13 works in this case.




FIGURE 4

Asymptotes of the spectral curve for the rigid body

The equation of the spectral curve, written in affine coordinates $\lambda = z_1/z_3$, and $\mu = z_2/z_3$, is $\det(X + \lambda A - \mu \text{Id}) = 0$. Using that $X^t = -X$, we have

$$\begin{aligned} \det(X + \lambda A - \mu \text{Id}) &= \det((X + \lambda A - \mu \text{Id})^t) = \\ &= \det(-X + \lambda A - \mu \text{Id}) = -\det(X - \lambda A + \mu \text{Id}), \end{aligned}$$

so, for a skew-symmetric X , the spectral curve C_X is symmetric with respect to the origin. The latter in particular implies that C_X has two ovals both of which are unbounded.

Further, let us show that if the matrix X is skew-symmetric, then the hexagon H_X is of type (2,4). Figure 4 depicts the asymptotes l_1, l_2, l_3 of the spectral curve and six sectors into which these asymptotes cut the affine plane. The asymptote l_i is given by the equation $\mu = a_i \lambda$ (without loss of generality, we may assume that $a_1 > a_2 > a_3$). According to Remark 2.9, the vertex P_1 of the hexagon H_X may be found as one of the intersection points of the spectral curve with the conic $M_{11} = 0$. In affine coordinates, this conic is given by

$$(\mu - a_2 \lambda)(\mu - a_3 \lambda) + x_{23}^2 = 0,$$

i.e. it is a hyperbola whose branches lie in sectors 1 and 1'. Therefore, the point P_1 lies in one of these sectors. The branch of the curve C_X lying in the corresponding sector joins the point P_1 with at least one of the points ∞_2 or ∞_3 . So, P_1 lies in the same oval as ∞_2 or ∞_3 , and thus the hexagon H_X is indeed of type (2,4) (see Figure 3). The latter allows us to conclude that all generic solutions of the rigid body equation exist for all times. This is, of course, very well known (these trajectories are in fact periodic).

Example 2.15 (Rigid body in a pseudo-Euclidian space). Equation (1.1) may also be restricted to the Lie algebra $\mathfrak{so}(1, 2)$, which consists of matrices satisfying the equation $XI + IX^t = 0$ where I is diagonal $I := \text{diag}(1, 1, -1)$. For such matrices X , equation (1.1) may be regarded as the equation of a rigid body in the pseudo-Euclidian space $\mathbb{R}^{1,2}$. (Rigid body dynamics in non-Euclidian spaces was considered, e.g., in a recent paper [BoM].)

As in the Euclidian case, the spectral curve C_X is symmetric with respect to the origin and thus has two unbounded ovals. The difference is that we can no longer assume that $a_1 > a_2 > a_3$ due to the special role of the first coordinate. So, there are two different cases. The first case is when a_1 does not lie in the interval (a_2, a_3) . Then, repeating the argument of the Euclidian case (see Example 2.14), one shows that the hexagon H_X is of type $(2, 4)$, and thus all generic trajectories exist for all times. The second case is $a_1 \in (a_2, a_3)$. In this case, a similar argument shows that the hexagon H_X is of type $(4, 2)$. Thus, if a_1 is between a_2 and a_3 , then all generic trajectories blow up in finite time.

Now, let us give an informal explanation why Theorem 2.13 is true. First, assume that the real part of the spectral curve C_X has one oval. Then, as the matrix X evolves according to equation (1.1), the hexagon H_X , and in particular its vertex P_1 , slide along the spectral curve C_X . At some point of time t' , the vertex P_1 hits the point ∞_1 , and the hexagon H_X becomes degenerate. However, as was pointed out above, the degenerate hexagon does not correspond to any matrix X . For this reason, the solution $X(t)$ can not be extended to $t = t'$.

Further, note that if the real part of the spectral curve has two ovals, but the hexagon H_X is of type $(0, 6)$ or $(4, 2)$, then the points P_1 and ∞_1 still lie in the same oval (see Figure 3). So, we arrive to exactly the same conclusion as in the one oval case.

Finally, if the real part of C_X has two ovals, and H_X is of type $(6, 0)$ or $(2, 4)$, then the points P_1 and ∞_1 lie in different ovals. For this reason, they can never meet each other, and the solution exists for all times.

Of course, to turn this explanation into a rigorous proof, one should understand the dynamics of the hexagon H_X . It turns out that this dynamics is, roughly speaking, a uniform rotation. More precisely, there exists an angular coordinate $\phi \in [0, 2\pi)$ on each oval of the curve C_X such that the evolution of all vertices of H_X is given by $d\phi/dt = \text{const} \neq 0$. Thus, if the points P_1 and ∞_1 lie in the same oval, they will inevitably meet each other.

Note that this consideration also implies that for each global in time solution of equation (1.1), the hexagon H_X returns to its initial position after some time T . In other words, the evolution of the hexagon H_X is periodic. However, the evolution of the matrix X itself is, in general, not periodic but quasi-periodic.

To be more precise, we have the following.

Theorem 2.16. *Assume that A is a generic diagonal matrix, and let C be a smooth real cubic passing through the points $\infty_1, \infty_2, \infty_3$. Further, assume that the real part of C has two ovals³. Then there exist a real number $T > 0$ and a diagonal matrix $M \in \mathbb{PD}(3, \mathbb{R})$ such that for each γ lying in $\mathcal{T}_C$ global in time solution of equation (1.1), the following statements hold.*

- (1) *The dynamics of the hexagon H_X is periodic with period T .*
- (2) *The dynamics of the matrix X is quasi-periodic:*

$$X(t + T) = MX(t)M^{-1}.$$

Furthermore, we have $M \in \mathbb{PD}^+(3, \mathbb{R})$ if H_X has type $(6, 0)$, and $M \notin \mathbb{PD}^+(3, \mathbb{R})$ if H_X has type $(2, 4)$.

Example 2.17 (Rigid body revisited). Let us again consider the case of a skew-symmetric matrix X . Then, as follows from considerations of Example 2.14, the hexagon H_X has type $(2, 4)$. Therefore, $X(t + T) = MX(t)M^{-1}$ where $M \in \mathbb{PD}(3, \mathbb{R}) \setminus \mathbb{PD}^+(3, \mathbb{R})$. On the other hand, all generic trajectories of the rigid body are closed, so we should have $M^k = \text{Id}$ for a suitable integer $k > 0$. Clearly, this is only possible when the diagonal entries of M are equal to ± 1 , and thus $M^2 = \text{Id}$. Note that M itself is not $\pm \text{Id}$ since $M \notin \mathbb{PD}^+(3, \mathbb{R})$. So, we have $X(t + 2T) = X(t)$, i.e. the period of a generic trajectory of the rigid body is *twice* the period of the corresponding hexagon.

Remark 2.18. Let us comment on the relation between the hexagon H_X and the general approach of the algebro-geometric integration theory. In this general approach, one considers eigenvectors of the Lax matrix X_λ as a line bundle E over the spectral curve. The fiber of the bundle E at the point (λ, μ) is the eigenspace of X_λ corresponding to the eigenvalue μ (one can show that for smooth spectral curves this eigenspace is always one-dimensional, and that the line bundle E extends to the points at infinity). The isomorphism class of the line bundle E defines a point in the Jacobian of the spectral curve. The main result of the algebro-geometric integration theory is that the evolution of this point according to the Lax equation is linear with respect to the addition law on the Jacobian (see, e.g., the above-mentioned monograph [BBT]; cf. Proposition 3.4 below). From the latter it, in particular, follows that equation (1.1) can be solved in terms of theta functions (cf. [Man]).

³ Recall that if the real part of C has one oval, then there are no global in time solutions lying in $\mathcal{T}_C$.

The relation between the line bundle E and the hexagon H_X is as follows. For any regularly inscribed hexagon H , we have

$$(2.5) \quad P_1 + \infty_2 + \infty_3 \sim \infty_1 + P_2 + \infty_3 \sim \infty_1 + \infty_2 + P_3,$$

where $D_1 \sim D_2$ denotes linear equivalence of divisors D_1, D_2 . Indeed, the divisor of the function L_{13}/L_{23} , where L_{ij} is given by (2.3), is $P_1 + \infty_2 - \infty_1 - P_2$ (see Figure 2). Therefore, $P_1 + \infty_2 \sim \infty_1 + P_2$, and $P_1 + \infty_2 + \infty_3 \sim \infty_1 + P_2 + \infty_3$, as desired. The proof of the equivalence $\infty_1 + P_2 + \infty_3 \sim \infty_1 + \infty_2 + P_3$ is analogous. Furthermore, one can show that the line bundle corresponding to divisors (2.5) is isomorphic to the eigenvector bundle E . Thus, the data contained in the hexagon H_X and the line bundle E are equivalent. However, it turns out that it is easier to read off the information about the matrix X from the corresponding hexagon rather than from the line bundle.

3. Proofs of the main results

3.1. Regularly inscribed hexagons and Chasles' theorem. In this section we prove that for any point lying in the real part of a cubic curve, there exists a unique regularly inscribed hexagon whose vertex P_1 is at that point. Of course, there is nothing special about the vertex P_1 , so the result is also true for any other vertex.

Let C be a smooth real cubic curve, and assume that C intersects the line at infinity at three real points $\infty_1, \infty_2, \infty_3$. Then C has three real asymptotes l_1, l_2, l_3 . Take any point $P_1 \in C_{\mathbb{R}}$ and consider the line passing through P_1 and parallel to l_3 , that is the line passing through P_1 and ∞_3 (if $P_1 = \infty_3$, then the line passing through P_1 and ∞_3 is, by definition, the tangent line to C at ∞_3 , i.e. the asymptote l_3). By Bézout's theorem, this line should meet the curve C at one more point which we call Q_2 (see Figure 2). Clearly, $Q_2 \in C_{\mathbb{R}}$. Now, consider the line passing through Q_2 and parallel to l_1 , and denote its third intersection point with C by P_3 . Continuing this procedure, we obtain points $P_1, Q_2, P_3, Q_1, P_2, Q_3, P_1^* \in C_{\mathbb{R}}$ such that

$$P_1 Q_2 \parallel l_3, \quad Q_2 P_3 \parallel l_1, \quad P_3 Q_1 \parallel l_2, \quad Q_1 P_2 \parallel l_3, \quad P_2 Q_3 \parallel l_1, \quad Q_3 P_1^* \parallel l_2.$$

Now, we need to show that the points P_1^* and P_1 in fact coincide, so that the polygon $P_1 Q_2 P_3 Q_1 P_2 Q_3$ is a regularly inscribed hexagon. For simplicity, assume that the nine points

$$P_1, Q_2, P_3, Q_1, P_2, Q_3, \infty_1, \infty_2, \text{ and } \infty_3$$

are pairwise distinct (the general case follows by continuity). We shall apply the following classical result, known as Chasles', or Cayley–Bacharach theorem:

Theorem 3.1. *Let C_1, C_2 , and C_3 be three cubic curves in $\mathbb{CP}^2$. Assume that C_1 intersects C_2 at nine distinct points $R_1, \dots, R_9$, and that C_3 passes through eight of these nine points. Then C_3 also passes through the ninth point.*

To apply this result in our setting, consider the curves

$$C_1 = C, \quad C_2 = P_1 Q_2 \cup P_3 Q_1 \cup P_2 Q_3, \quad \text{and} \quad C_3 = Q_2 P_3 \cup Q_1 P_2 \cup Q_3 P_1^*.$$

Then

$$C_1 \cap C_2 = \{P_1, Q_2, P_3, Q_1, P_2, Q_3, \infty_1, \infty_2, \infty_3\},$$

and thus C_3 passes through all points of $C_1 \cap C_2$ except, possibly, P_1 . Therefore, by Theorem 3.1, the curve C_3 also passes through the point P_1 , so

$$P_1 \in C_1 \cap C_3 = \{Q_2, P_3, Q_1, P_2, Q_3, P_1^*, \infty_1, \infty_2, \infty_3\},$$

and hence $P_1 = P_1^*$, q.e.d.

3.2. Reconstructing a matrix from a curve and a hexagon. In this section we show that a matrix X can be reconstructed from the spectral curve C_X and the hexagon H_X uniquely up to conjugation by diagonal matrices. For the sake of simplicity, we shall assume that the spectral curve satisfies the following genericity assumptions: (i) it does not have inflection points at infinity, (ii) intersection points of asymptotes do not lie on the curve. It is easy to see that under these assumptions every regularly inscribed hexagon which is not degenerate has at most one side at infinity. The proof of the general case is similar.

Proposition 3.2. *Let C be a smooth real projective cubic passing through the points $\infty_1, \infty_2, \infty_3$, and satisfying the above genericity assumptions. Let also $H \subset C$ be a regularly inscribed hexagon which is not degenerate. Then there exists a matrix $X \in \mathfrak{gl}(3, \mathbb{R})$, unique up to conjugation by a diagonal matrix, such that $C_X = C$, and $H_X = H$.*

Proof. First note that the spectral curve C_X and the hexagon H_X are invariant under transformations $X \mapsto DXD^{-1}$ where D is a diagonal invertible matrix. Therefore, uniqueness up to conjugation by diagonal matrices is the best result we can expect.

Now, let us show how to reconstruct X from C and H . Note that by Proposition 2.7 the diagonal entries x_{11}, x_{22}, x_{33} of the matrix X are uniquely determined by the spectral curve. So, we only need to reconstruct the off-diagonal entries.

First, assume that none of the sides of H are at infinity. This also implies that none of the sides of H are asymptotes of C . In terms of the matrix X to

be constructed, these conditions mean that all off-diagonal entries of X do not vanish (cf. formula (2.3)). Take any two non-zero real numbers $\alpha, \beta \in \mathbb{R}^*$, and set $x_{31} = \alpha, x_{32} = \beta$. Now, we find the remaining entries of X by using the equations of sides of H . First, we find x_{12} from the equation of l_{23} . Since the side l_{23} is parallel to the asymptote l_1 , its equation has the form

$$a(x_{11}z_3 + a_1z_1 - z_2) - bz_3 = 0$$

where a, b are constants. Note that since l_{23} is neither the line at infinity, nor an asymptote, these constants do not vanish. On the other hand, in terms of the matrix X to be constructed, the equation of l_{23} should be

$$x_{32}(x_{11}z_3 + a_1z_1 - z_2) - x_{31}x_{12}z_3 = 0,$$

so

$$\frac{x_{32}}{a} = \frac{x_{31}x_{12}}{b},$$

which allows us to find x_{12} . In a similar way, we find x_{21} from the equation of l_{13} , then x_{23} from l_{12} , and, finally, x_{13} from l_{32} .

Now, let us show that the curves C_X and C are the same. As follows from the construction of X , the hexagons H_X and H have four common consecutive sides l_{23} , l_{13} , l_{12} , and l_{32} . Therefore, they have three common vertices Q_3, P_1 , and Q_2 . By Proposition 2.8, these points are not collinear. So, the curves C and C_X are two cubics which have common asymptotes and three common non-collinear points. As it is easy to see, such cubics must coincide. This, in turn, implies that the hexagons H and H_X coincide as well: they are two hexagons which are regularly inscribed in the same cubic and have a common vertex P_1 .

Now, assume that H has one side at infinity. Without loss of generality, it is l_{31} . Then the two adjacent sides of H are necessarily asymptotes of the curve, namely $l_{32} = l_1$, and $l_{21} = l_3$. As above we fix $\alpha, \beta \in \mathbb{R}^*$ and set $x_{31} = \alpha, x_{32} = \beta$. Apart from this, since the side l_{31} is at infinity, we should set $x_{13} = 0$ (see the proof of Proposition 2.8). Further, similarly to the above, we find x_{12} , x_{21} , x_{23} using the equations of l_{23} , l_{13} , and l_{12} respectively. Note that since $x_{13} = 0$, the sides l_{32} and l_{21} of the hexagon H_X automatically coincide with the corresponding sides of H , that is with asymptotes l_1, l_3 . So, H_X and H have five sides in common and thus, similarly to the above, C_X coincides with C , and H_X coincides with H .

So, we showed that in both cases a matrix X satisfying the desired conditions is uniquely determined by its non-zero entries x_{31}, x_{32} . Conjugating such a matrix with a suitable diagonal matrix, we can always assume that $x_{31} = 1$ and $x_{32} = 1$. Therefore, X is indeed unique up to conjugation by a diagonal matrix. $\square$

3.3. Description of isospectral sets. In this section, we prove the topological part of Theorem 2.5. Namely, we describe the topology of the sets $\mathcal{T}_C = \{X \in \mathfrak{gl}(3, \mathbb{R}) : C_X = C\}$.

Let $\mathcal{H}_C$ be the set of hexagons regularly inscribed in the curve C . Let also $H_d \in \mathcal{H}_C$ be the degenerate hexagon. Proposition 3.2 allows us to conclude that the set $\mathcal{T}_C$ is the total space of a principal $\mathbb{PD}(3, \mathbb{R})$ bundle over $\mathcal{H}_C \setminus H_d$. Also note that, according to Section 3.1, the space $\mathcal{H}_C \setminus H_d$ can be identified with the real part of the curve C without a point.

A trivializing cover for the bundle $\mathcal{T}_C \rightarrow \mathcal{H}_C \setminus H_d$ can be constructed as follows. Let

$$U_1 = \{H \in \mathcal{H}_C : l_{13} \neq l_2\}, \quad U_2 = \{H \in \mathcal{H}_C : l_{31} \neq l_2\}.$$

It is easy to see that, under the genericity assumptions of the previous section, $\{U_1, U_2\}$ is indeed a cover of $\mathcal{H}_C \setminus H_d$. As coordinates on fibers over U_1 , we take the entries x_{21} and x_{32} of the matrix X . Since $l_{13} \neq l_2$, these entries are non-zero (cf. formula (2.3)), and thus they uniquely determine a matrix X within its $\mathbb{PD}(3, \mathbb{R})$ orbit. Similarly, we take x_{12} and x_{23} as coordinates on fibers over U_2 .

Now, let us prove the first statement of Theorem 2.5. If the real part $C_{\mathbb{R}}$ of the curve C has one oval, then the set $\mathcal{H}_C \simeq C_{\mathbb{R}}$ is diffeomorphic to a circle S^1 . Therefore, the set $\mathcal{H}_C \setminus H_d$ is diffeomorphic to $\mathbb{R}$, and hence $\mathcal{T}_C \rightarrow \mathcal{H}_C \setminus H_d$ is a trivial bundle: $\mathcal{T}_C \simeq \mathbb{R} \times \mathbb{PD}(3, \mathbb{R})$. The latter set has four connected components diffeomorphic to $\mathbb{R}^3$, q.e.d.

Further, let us prove the second statement. Assume that the real part of C has two ovals one of which is bounded. In this case, the set $\mathcal{H}_C$ has two connected components distinguished by the type of a hexagon. Let

$$\mathcal{H}_C^{i,j} := \{H \in \mathcal{H}_C : H \text{ is of type } (i, j)\}, \quad \mathcal{T}_C^{i,j} := \{X \in \mathcal{T}_C : H_X \in \mathcal{H}_C^{i,j}\}.$$

Then

$$\mathcal{H}_C = \mathcal{H}_C^{6,0} \sqcup \mathcal{H}_C^{0,6}, \quad \text{and} \quad \mathcal{T}_C = \mathcal{T}_C^{6,0} \sqcup \mathcal{T}_C^{0,6}.$$

Note that since the degenerate hexagon H_d has type $(0, 6)$, we have

$$\mathcal{H}_C \setminus H_d = \mathcal{H}_C^{6,0} \sqcup (\mathcal{H}_C^{0,6} \setminus H_d).$$

Therefore, the set $\mathcal{T}_C^{0,6}$ is the total space of a principal $\mathbb{PD}(3, \mathbb{R})$ bundle over $\mathcal{H}_C^{0,6} \setminus H_d$. Since the latter set $\mathcal{H}_C^{0,6} \setminus H_d$ is diffeomorphic to $\mathbb{R}$, this bundle is trivial, and thus $\mathcal{T}_C^{0,6} \simeq 4\mathbb{R}^3$.

Now, let us study the component $\mathcal{T}_C^{6,0}$. By definition of the set $\mathcal{H}_C^{6,0}$, all vertices of any hexagon $H_X \in \mathcal{H}_C^{6,0}$ lie on the bounded oval, and therefore none of the vertices are at infinity. In particular, the side l_{13} of H_X cannot coincide

with the asymptote l_2 . So, the component $\mathcal{H}_C^{6,0}$ is completely covered by the chart U_1 , and thus the bundle $\mathcal{T}_C^{6,0} \rightarrow \mathcal{H}_C^{6,0}$ is also trivial. Since $\mathcal{H}_C^{6,0} \simeq S^1$, we have $\mathcal{T}_C^{6,0} \simeq 4S^1 \times \mathbb{R}^2$, q.e.d.

Finally, let us prove the third statement of Theorem 2.5. Assume that the real part of C has two unbounded ovals. Then

$$\mathcal{H}_C \setminus H_d = \mathcal{H}_C^{2,4} \sqcup (\mathcal{H}_C^{4,2} \setminus H_d).$$

Similarly to the above, we have $\mathcal{H}_C^{4,2} \setminus H_d \simeq \mathbb{R}$, therefore the bundle $\mathcal{T}_C^{4,2} \rightarrow \mathcal{H}_C^{4,2} \setminus H_d$ is trivial, and $\mathcal{T}_C^{4,2} \simeq 4\mathbb{R}^3$.

Now, let us show that the bundle $\mathcal{T}_C^{2,4} \rightarrow \mathcal{H}_C^{2,4}$ is *not* trivial. Note that there are only two different principal $\mathbb{P}\mathcal{D}(3, \mathbb{R})$ bundles over the circle S^1 , namely the trivial one with the total space $4S^1 \times \mathbb{R}^2$, and the non-trivial one with the total space $2S^1 \times \mathbb{R}^2$. So, as soon as we prove that the bundle is non-trivial, the topology of the total space is uniquely determined.

Without loss of generality, we may assume that the points ∞_2, ∞_3 lie on the contractible oval O_1 , and that the point ∞_1 lies on the non-contractible oval O_2 (if this is not so, we renumber these points). Then $P_1 \in O_1$, and $Q_3 \in O_2$, as depicted in Figure 5. Let us consider the intersection of the chart U_1 with the component $\mathcal{H}_C^{2,4}$. The complement to U_1 consists of those hexagons for which $l_{13} = l_2$. The latter is possible if either $P_1 = \infty_2$, or $Q_3 = \infty_2$. However, for $H_X \in \mathcal{H}_C^{2,4}$, the points Q_3 and ∞_2 lie on different ovals (see Figure 5), therefore the chart U_1 covers the whole set $\mathcal{H}_C^{2,4}$ except one hexagon H_0 distinguished by the condition $P_1 = \infty_2$. This hexagon can be obtained by moving the point Q_3 in Figure 5 to the right till it reaches the asymptote l_2 . The domain $\mathcal{H}_C^{2,4} \setminus H_0$ is covered by the chart U_1 , therefore in this domain we have a trivialization of




FIGURE 5

the bundle given by x_{21}, x_{32} . Let us analyze what happens to these coordinates as the hexagon H_X passes through H_0 , or, which is the same, as the vertex Q_3 crosses the asymptote l_2 . First, note that the side l_{12} of the hexagon H_0 is at infinity, therefore for this hexagon we have $x_{21} = 0$ (see the proof of Proposition 2.8), which shows that the trivialization $\{x_{21}, x_{32}\}$ is no longer valid for $H_X = H_0$. However, since H_0 has only one side at infinity, we have $x_{32} \neq 0$. Therefore, to determine whether the bundle $\mathcal{T}_C^{2,4} \rightarrow \mathcal{H}_C^{2,4}$ is trivial, we should study what happens to the sign of x_{21} as the point Q_3 crosses the asymptote l_2 . By formula (2.3), the side l_{13} is given by the equation

$$x_{31}(x_{22}z_3 + a_2z_1 - z_2) - x_{21}x_{32}z_3 = 0.$$

At the same time, $x_{22}z_3 + a_2z_1 - z_2 = 0$ is the equation of the asymptote l_2 . Therefore, provided that $x_{31} \neq 0$, i.e. that the side l_{13} is not at infinity, the sign of the product $x_{21}x_{32}$ has the following geometric meaning: it is positive if the line l_{13} lies on one side of the asymptote l_2 , and negative if it lies at the other side. Now, notice that as Q_3 crosses the asymptote l_2 , the line l_{13} gets from one side of the asymptote to the other (see Figure 5), therefore the sign of the product $x_{21}x_{32}$ changes to its negative. Since x_{32} does not vanish as Q_3 crosses the asymptote, this means that the sign of x_{21} changes, and therefore the bundle is non-trivial. So, we have $\mathcal{T}_C^{2,4} \simeq 2S^1 \times \mathbb{R}^2$, thus the third statement of Theorem 2.5 is proved.

3.4. Dynamics of the hexagon. Let us fix the spectral curve C and describe the evolution of the hexagon H_X under the flow (1.1). As before, consider the matrix $X_z = z_3X + z_1A - z_2\text{Id}$. Recall that the zero locus of the determinant of this matrix is the spectral curve C . We shall describe the dynamics of the hexagon H_X by relating its vertices to eigenvectors of X_z . Note that since a regularly inscribed hexagon is uniquely determined by any of its vertices, it suffices to describe the dynamics of one vertex. As before, let M_{ij} be the (i, j) minor of X_z . Then the vector $(M_{11}, -M_{12}, M_{13})^t$ belongs to the kernel of X_z . Normalizing this vector by dividing its components by M_{11} , we obtain the following meromorphic vector-function ψ on C :

$$\psi := \begin{pmatrix} 1 \\ -\frac{M_{12}}{M_{11}} \\ \frac{M_{13}}{M_{11}} \end{pmatrix}$$

Denote the components of this vector by ψ_1, ψ_2, ψ_3 . By construction, we have $X_z\psi = 0$ identically on C .

Proposition 3.3. *The component ψ_2 of ψ has poles at ∞_2 and Q_1 and zeros at ∞_1 and Q_2 ; the component ψ_3 has poles at ∞_3 and Q_1 and zeros at ∞_1 and Q_3 .*

Proof. As follows from Remark 2.9, zeros of M_{11} on C are the points P_1, Q_1 , and the points ∞_2, ∞_3 taken with multiplicity 2. Further, we have $M_{12} = z_3 L_{12}$ where L_{12} is a linear function whose zero locus is the line l_{12} . Therefore, zeros of M_{12} are the points $P_1, Q_2, \infty_1, \infty_2$, and the point ∞_3 taken with multiplicity 2. Dividing M_{12} by M_{11} , we obtain the desired statement about zeros and poles of ψ_2 . Zeros and poles of ψ_3 are computed analogously. $\square$

So, the vertex Q_1 of the hexagon H_X is the only movable pole of the eigenvector ψ (clearly, vertices Q_2 and Q_3 can be interpreted in the same way; to obtain these vertices as poles, one needs to renormalize the vector ψ by setting $\psi_2 = 1$ or $\psi_3 = 1$). This allows us to describe the dynamics of Q_1 using standard technique (see e.g. the reviews [DKN, DMN, RSTS3]). Let ω be a holomorphic 1-form on C ; such a form is unique up to a constant factor. In the affine chart $\lambda = z_1/z_3$ and $\mu = z_2/z_3$, it is given by

$$(3.1) \quad \omega = \frac{d\mu}{\partial_\lambda f_X} = -\frac{d\lambda}{\partial_\mu f_X}$$

where $f_X(\lambda, \mu) = 0$ is the equation of the affine part of the curve C .

Proposition 3.4. *Assume that X evolves according to equation (1.1). Then, for the holomorphic form ω normalized by (3.1), we have*

$$\omega \left(\frac{dQ_1}{dt} \right) = 1.$$

Proof. We work in affine coordinates $\lambda = z_1/z_3$ and $\mu = z_2/z_3$. Equation $X_z \psi = 0$ can be rewritten as

$$(3.2) \quad (X_\lambda - \mu \text{Id}) \psi = 0$$

where $X_\lambda = X + \lambda A$. Assuming that X_λ evolves according to Lax equation (2.1) and differentiating (3.2) with respect to time, we get

$$(3.3) \quad (X_\lambda - \mu \text{Id}) \left(\frac{d\psi}{dt} + Y_\lambda \psi \right) = 0.$$

Note that $\text{rank}(X_\lambda - \mu \text{Id}) = \text{rank } X_z = 2$ at every point of C . Indeed, if $\text{rank } X_z = 1$ at some point $P \in C$, then this point is a common zero for all 2×2 minors M_{ij} of the matrix X_z , which means that all vertices of the hexagon

H_X coincide. However, as follows from Proposition 2.8, this is impossible. Thus, (3.3) implies that

$$(3.4) \quad \frac{d\psi}{dt} = (\xi \cdot \text{Id} - Y_\lambda)\psi,$$

where ξ is a function on the curve C . Using that $\psi_1 \equiv 1$, we find that ξ is equal to the first coordinate $(Y_\lambda \psi)_1$ of the vector $Y_\lambda \psi$.

Now, let us consider the point Q_1 at some moment of time t_0 , and let u be a local coordinate on C near this point. Let also $u_q(t)$ be the u -coordinate of Q_1 at moment t . Provided that Q_1 does not coincide with ∞_2 or ∞_3 , we have

$$(3.5) \quad \psi(t) = \frac{h(t)}{u - u_q(t)} + \text{terms holomorphic in } u,$$

where $h(t)$ is a vector holomorphic in u . Substituting (3.5) into (3.4) and equating coefficients in $(u - u_q(t))^{-2}$, we get

$$\frac{du_q}{dt} = (Y_\lambda h)_1,$$

where $(Y_\lambda h)_1$ is the first coordinate of the vector $Y_\lambda h$. The coordinate free-form of this equation is

$$(3.6) \quad \omega \left(\frac{dQ_1}{dt} \right) = \text{Res}_{Q_1} ((Y_\lambda \psi)_1 \omega),$$

where ω is the holomorphic form defined above. Now, note that the form $(Y_\lambda \psi)_1 \omega$ may only have poles at those points where either Y_λ or ψ have a pole, i.e. at points $Q_1, \infty_1, \infty_2, \infty_3$. Therefore, by Cauchy's residue theorem, we have

$$(3.7) \quad \text{Res}_{Q_1} ((Y_\lambda \psi)_1 \omega) = - \sum_{i=1}^3 \text{Res}_{\infty_i} ((Y_\lambda \psi)_1 \omega).$$

Further, note that $Y_\lambda = \lambda^{-1}(X_\lambda^2 - X^2)$ (cf. Formula (2.2)), therefore,

$$\begin{aligned} \text{Res}_{\infty_i} ((Y_\lambda \psi)_1 \omega) &= \text{Res}_{\infty_i} ((X_\lambda^2 \psi)_1 \lambda^{-1} \omega) - \text{Res}_{\infty_i} ((X^2 \psi)_1 \lambda^{-1} \omega) \\ &= \text{Res}_{\infty_i} (\mu^2 \lambda^{-1} \omega) \end{aligned}$$

where we used the identities $X_\lambda \psi = \mu \psi$, $\psi_1 = 1$, and that the function $(X^2 \psi)_1 \lambda^{-1}$ does not have a pole at the point ∞_i . Combining the last formula with (3.6) and (3.7), we conclude that

$$(3.8) \quad \omega \left(\frac{dQ_1}{dt} \right) = - \sum_{i=1}^3 \text{Res}_{\infty_i} (\mu^2 \lambda^{-1} \omega).$$

Note that although we assumed in the proof that $Q_1 \neq \infty_2$ and $Q_1 \neq \infty_3$, formula (3.8) still holds for these points by continuity argument. Now, to complete the proof, it suffices to compute the residues. This can be easily done using formula (3.1) and the explicit expression for the polynomial f_X (note that only cubic terms of f_X affect the residues). $\square$

Note that the conclusion of Proposition 3.4 is obviously true for other Q -vertices of the hexagon H_X as well, i.e.

$$\omega\left(\frac{dQ_1}{dt}\right) = \omega\left(\frac{dQ_2}{dt}\right) = \omega\left(\frac{dQ_3}{dt}\right) = 1.$$

For P -vertices, we have

$$\omega\left(\frac{dP_1}{dt}\right) = \omega\left(\frac{dP_2}{dt}\right) = \omega\left(\frac{dP_3}{dt}\right) = -1.$$

The latter can be proved by noting that (1.1) is anti-invariant with respect to the transformation $X \mapsto X^t$ which preserves the spectral curve C and interchanges P -vertices with Q -vertices.

Now, for each oval O_i of the curve C , fix a point $R_i \in O_i$ and consider the function

$$\phi(R) = \int_{R_i}^R \omega.$$

Then ϕ is a periodic coordinate on O_i . In terms of the coordinate ϕ , the dynamics of vertices of H_X is linear: $d\phi/dt = \pm 1$. This in particular implies that the dynamics of the hexagon H_X is monotonous and periodic. The period is given by the integral of ω along any of the ovals of C (note that this integral is the same for both ovals since they are homologous cycles in C).

3.5. Complete and blow-up solutions. In this section, we prove Theorems 2.13 and 2.16, i.e. we investigate the dynamics of (1.1) at each connected component of the set $\mathcal{T}_C$. As we know from Section 3.3, the set $\mathcal{T}_C$ is the total space of a principal $\mathbb{P}D(3, \mathbb{R})$ bundle over the set $\mathcal{H}_C \setminus H_d$ where $\mathcal{H}_C$ is the set of all hexagons regularly inscribed in C , and H_d is the degenerate hexagon. Furthermore, flow (1.1) is invariant with respect to the $\mathbb{P}D(3, \mathbb{R})$ action on $\mathcal{T}_C$. This allows us to apply the following classical result.

Theorem 3.5 (A. Lichnerowicz [Lic]). *Let $\pi: E \rightarrow B$ be a principal G -bundle, and let v be a vector field on E which is invariant with respect to the G -action. Then an integral trajectory $x(t)$ of the field v is complete if and only if the corresponding trajectory of the field π_*v on the base is complete.*

This result implies that one can study the behavior of solutions of (1.1) by considering their projections to the space of hexagons. Dynamics on the space of hexagons was studied in the previous section: it is linear.

To prove Theorem 2.13, we consider separately each of the components $\mathcal{T}_C^{i,j}$ defined in Section 3.3. For example, consider a solution $X(t)$ of equation (1.1) such that the corresponding hexagon H_X has type $(0, 6)$. The latter means that $X(t) \in \mathcal{T}_C^{0,6}$ in the notation of Section 3.3. The set $\mathcal{T}_C^{0,6}$ is a principal bundle over $\mathcal{H}_C^{0,6} \setminus H_d$. Furthermore, as follows from Section 3.4, the dynamics on $\mathcal{H}_C^{0,6}$ is linear in terms of the coordinate ϕ , so the projection of $X(t)$ to the base $\mathcal{H}_C^{0,6} \setminus H_d$ meets the degenerate hexagon H_d and thus blows up in finite time. Therefore, by Theorem 3.5, the trajectory $X(t)$ itself also blows up.

An analogous consideration shows that trajectories of (1.1) corresponding to hexagons of type $(4, 2)$ also blow up, while trajectories corresponding to types $(6, 0)$ or $(2, 4)$ exist for all times. Thus, Theorem 2.13 is proved.

Note that this consideration also proves the dynamical part of Theorem 2.5 since all $\mathbb{R}^3$ components of $\mathcal{T}_C$ correspond to hexagons of type $(0, 6)$ or $(4, 2)$, while all $S^1 \times \mathbb{R}^2$ components correspond to $(6, 0)$ or $(2, 4)$ (see Section 3.3).

Now, let us prove Theorem 2.16. As follows from the previous section, if a trajectory $X(t)$ exists for all times, then the dynamics of the corresponding hexagon is periodic with a certain period T . Now, the formula

$$X(t + T) = MX(t)M^{-1}.$$

easily follows from the $\mathbb{PD}(3, \mathbb{R})$ invariance of the flow (1.1). Further, let us show that $M \in \mathbb{PD}^+(3, \mathbb{R})$ if H_X has type $(6, 0)$. Consider the bundle $\mathcal{T}_C^{6,0} \rightarrow \mathcal{H}_C^{6,0}$. By definition of the number T , the matrices $X(t + T)$ and $X(t)$ lie in the same fiber of this bundle. Furthermore, since it is a trivial bundle, $X(t + T)$ and $X(t)$ should lie in the same connected component of the fiber, and thus $M \in \mathbb{PD}^+(3, \mathbb{R})$.

Analogously, since the bundle $\mathcal{T}_C^{2,4} \rightarrow \mathcal{H}_C^{2,4}$ is not trivial, we have $M \notin \mathbb{PD}^+(3, \mathbb{R})$ if H_X has type $(2, 4)$, so Theorem 2.16 is proved.

4. Discussion

Note that equation (1.1), as well as the definitions of the spectral curve and the corresponding set $\mathcal{T}_C$, can be without any difficulty generalized to $\mathfrak{gl}_n(\mathbb{R})$. It is an interesting question how to generalize our results to this case. In particular, is it always true that the topology of $\mathcal{T}_C$ is completely determined by the geometry of $C_{\mathbb{R}}$? It is particularly interesting whether the structure of the set $\mathcal{T}_C$ depends on the way the ovals of C are nested into each other. (Recall that description of

all possible relative positions of ovals for a real algebraic curve of degree n is the Hilbert 16th problem which is still unsolved in full generality.)

We also note that the problem of description of the set $\mathcal{T}_C$ can be reformulated in purely algebro-geometric terms. Similarly to the $\mathfrak{gl}(3, \mathbb{R})$ case, let $a_1, \dots, a_n$ be the eigenvalues of the matrix A . For simplicity, let all eigenvalues $a_1, \dots, a_n$ be real, so that we may assume that A is a diagonal matrix. Further, let $\infty_i = (1 : a_i : 0) \in \mathbb{CP}^2$, and let C be a smooth real projective curve of degree n passing through the points $\infty_1, \dots, \infty_n$. Then, as follows from the results of [Bea], the manifold $\mathcal{T}_C = \{X \in \mathfrak{gl}(n, \mathbb{R}) : C_X = C\}$ can be described as a principal $(\mathbb{R}^*)^{n-1}$ bundle over the real part of $\text{Jac}(C) \setminus \Theta$ where $\text{Jac}(C)$ is the Jacobian of the curve C , and $\Theta \subset \text{Jac}(C)$ is the theta-divisor (the latter can also be deduced from the earlier papers [VMM, AvM2, RSTS2]).

Further, it is a classical result by Comessatti (see e.g. [SS]) that the topology of the real part of $\text{Jac}(C)$ is uniquely determined by the number of ovals of C . However, one still needs to understand the structure of the $(\mathbb{R}^*)^{n-1}$ bundle $\mathcal{T}_C \rightarrow (\text{Jac}(C) \setminus \Theta)_{\mathbb{R}}$. For $n = 3$ this is (in much more elementary terms) done in the present paper.

Also note that there is a more explicit algebro-geometric description of the set $\mathcal{T}_C$ due to L. Gavrilov [Gav]. Namely, consider the singular curve C_s obtained from the smooth curve C by identifying the points $\infty_1, \dots, \infty_n$, and let $\text{Jac}(C_s)$ be the generalized Jacobian of C_s . Then, as follows from [Gav], the set $\mathcal{T}_C$ is diffeomorphic to the real part of $\text{Jac}(C_s) \setminus \pi^{-1}(\Theta)$ where $\pi : \text{Jac}(C_s) \rightarrow \text{Jac}(C)$ is the canonical projection. Thus, to describe the set $\mathcal{T}_C$, one needs to describe the real part of the generalized Jacobian $\text{Jac}(C_s)$, and to study how it intersects the preimage of the theta divisor under the projection π .

Acknowledgments. The second author was partially supported by the Dynasty Foundation Scholarship and an NSERC research grant. The authors are grateful to Alexey Bolsinov and Boris Khesin for useful remarks.

References

- [Ale] K. R. ALESHKIN, The topology of integrable systems with incomplete fields. *Sbornik: Mathematics* **205** (2014), 1264. Zbl1338.37070 MR3288424
- [ALMR] M. S. ALBER, G. G. LUTHER, J. E. MARSDEN, and J. M. ROBBINS, Geometric phases, reduction and Lie-Poisson structure for the resonant three-wave interaction. *Physica D: Nonlinear Phenomena* **123** (1998), 271–290. Zbl1038.37510 MR1664938

- [AP] D. V. ALEKSEEVSKII and B. A. PUTKO, Completeness of left-invariant metrics on Lie groups. *Functional Analysis and Its Applications* **21** (1997), 233–234. Zbl 0657.53028
- [Arn1] V. ARNOLD, Sur la géométrie différentielle des groupes de lie de dimension infinie et ses applications à l'hydrodynamique des fluides parfaits. In *Annales de l'institut Fourier*, volume 16, pages 319–361. Institut Fourier, 1966. Zbl 0148.45301 MR 0202082
- [Arn2] V. I. ARNOLD, *Mathematical methods of classical mechanics*. Springer, 1989. Zbl 0692.70003 MR 0997295
- [AvM1] M. ADLER and P. Van MOERBEKE, Completely integrable systems, Euclidean Lie algebras, and curves. *Advances in mathematics* **38** (1980), 267–317. Zbl 0455.58017 MR 0597729
- [AvM2] M. ADLER and P. Van MOERBEKE, Linearization of Hamiltonian systems, Jacobi varieties and representation theory. *Advances in Mathematics* **38** (1980), 318–379. Zbl 0455.58010 MR 0597730
- [BBI⁺] A. M. BLOCH, V. BRÎNZĂNESCU, A. ISERLES, J. E. MARSDEN, and T. S. RATIU, A class of integrable flows on the space of symmetric matrices. *Communications in Mathematical Physics* **290** (2009), 399–435. Zbl 1231.37030 MR 2525626
- [BBT] O. BABELON, D. BERNARD and M. TALON, *Introduction to Classical Integrable Systems*. Cambridge University Press, 2003. Zbl 1045.37033 MR 1995460
- [Bea] A. BEAUVILLE, Jacobiennes des courbes spectrales et systemes hamiltoniens complètement intégrables. *Acta Mathematica*, 164(1):211–235, 1990. Zbl 0712.58031 MR 1049157
- [BI] A. M. BLOCH and A. ISERLES, On an isospectral lie–poisson system and its lie algebra. *Foundations of Computational Mathematics* **6** (2006), 121–144. Zbl 1105.37033 MR 2198217
- [BIKO] A. V. BOLSINOV, A. M. IZOSIMOV, A. YU. KONYAEV and A. A. OSJEMKOV, Algebra and topology of integrable systems: Problems for investigation. *Tr. Sem. Vektor. Tenzor. Anal* **28** (2012), 119–191.
- [BoM] A. V. BORISOV and I. S. MAMAEV, Rigid body dynamics in non-Euclidean spaces. *Russian Journal of Mathematical Physics* **23** (2016): 431–454.
- [BM] Sh. BROMBERG and A. MEDINA, Complétude de l'équation d'euler. In *Algebra and Operator Theory*, pages 127–144. Springer, 1998. Zbl 0923.58037 MR 1643439
- [DKN] B. A. DUBROVIN, I. M. KRICHEVER, and S. P. NOVIKOV, Integrable systems. I. *Dynamical systems IV, Encyclopaedia Math. Sci* **4** (1985), 179–284. Zbl 0780.58019 MR 0842910
- [DMN] B. A. DUBROVIN, V. B. MATVEEV, and S. P. NOVIKOV, Non-linear equations of Korteweg-de Vries type, finite-zone linear operators, and Abelian varieties. *Russian Mathematical Surveys* **31** (1976), 55–136. Zbl 0326.35011 MR 0427869

- [Gav] L. GAVRILOV, Generalized Jacobians of spectral curves and completely integrable systems. *Math. Zeitschrift* **230** (1999), 487–508. Zbl 0922.58033 MR 1679969
- [HSW] N. J. HITCHIN, G. B. SEGAL, and R. S. WARD, *Integrable Systems: Twistors, Loop Groups, and Riemann Surfaces*. Oxford University Press, 2013. Zbl 1268.37001
- [Lic] A. LICHNEROWICZ, *Global Theory of Connections and Holonomy Groups*. Noordhoff, 1976. Zbl 0337.53031 MR 0413000
- [Man] S. V. MANAKOV, Note on the integration of euler’s equations of the dynamics of an n -dimensional rigid body. *Functional analysis and its applications* **10** (1976), 328–329. Zbl 0358.70004
- [MF] A. S. MISHCHENKO and A. T. FOMENKO, Euler equations on finite-dimensional lie groups. *Mathematics of the USSR-Izvestiya* **12** (1978), 371. Zbl 0405.58031
- [RT] T. S. RATIU and D. TARAMA, The $U(n)$ free rigid body: Integrability and stability analysis of the equilibria. *Journal of Differential Equations* **259** (2015): 7284—7331.
- [RSTS1] A. G. REYMAN and M. A. SEMENOV-TIAN-SHANSKY, Reduction of Hamiltonian systems, affine Lie algebras and Lax equations. *Inventiones mathematicae* **54** (1979), 81–100. Zbl 0403.58004
- [RSTS2] A. G. REYMAN and M. A. SEMENOV-TIAN-SHANSKY, Reduction of Hamiltonian systems, affine Lie algebras and Lax equations II. *Inventiones mathematicae* **63** (1981), 423–432. Zbl 0442.58016 MR 0620678
- [RSTS3] A. G. REYMAN and M. A. SEMENOV-TIAN-SHANSKY, Group-theoretical methods in the theory of finite-dimensional integrable systems. In *Dynamical systems VII*, pages 116–225. Springer, 1994.
- [SS] M. SEPPÄLÄ and R. SILHOL, Moduli spaces for real algebraic curves and real abelian varieties. *Mathematische Zeitschrift* **201** (1989), 151–165. Zbl 0777.14014 MR 1135469
- [VMM] P. Van MOERBEKE and D. MUMFORD, The spectrum of difference operators and algebraic curves. *Acta Mathematica* **143** (1979), 93–154. Zbl 0502.58032 MR 0533894

(Reçu le 16 avril 2015)

Konstantin ALESHKIN, SISSA, via Bonomea 265, 34136 Trieste, Italy and Landau Institute for Theoretical Physics, 119334 Moscow, Russia

e-mail: kaleshkin@sissa.it

Anton IZOSIMOV, Department of Mathematics, University of Toronto, 40 St. George Street, Toronto, Ontario M5S 2E4, Canada

e-mail: izosimov@math.utoronto.ca