

Non-locality of the gapless plasma excitations in the QHE

Autor(en): **Grodnensky, I. / Heitmann, D. / Klitzing, K. von**

Objekttyp: **Article**

Zeitschrift: **Helvetica Physica Acta**

Band (Jahr): **65 (1992)**

Heft 2-3

PDF erstellt am: **26.09.2024**

Persistenter Link: <https://doi.org/10.5169/seals-116436>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*

ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

Non-locality of the gapless plasma excitations in the QHE

By I. Grodnensky¹, D. Heitmann² and K. von Klitzing²

¹Institute of Radioengineering and Electronics of the Academy of Sciences of the USSR,
Marx avenue 18, 103907, GSP-3, Moscow, USSR.

²Max-Planck Institut für Festkörperforschung, Heisenbergstrasse 1, D-7000 Stuttgart 80,
Federal Republic of Germany

Abstract. Edge magnetoplasma excitations in a two-dimensional electron system (2DES) have been studied in radio-frequency experiments. The plasmon frequency is shown to be governed by length l which characterizes the spatial distribution of the plasma charge oscillations in the direction perpendicular to the plasmon wavevector and demonstrates a nonlocal behaviour in the QHE regime.

The collective excitations spectrum of an electronic system at large wavevectors q is governed by the well known nonlocal effects. Here we report about a very different nonlocal effect which is an unique property of a finite size 2DES when the edges become important. We have observed this nonlocal effect in the QHE experiments on 2DES of macroscopic size ($1 \times 1 \text{ cm}^2$) for the low frequency ($\omega \ll \omega_c$) branch of the plasma excitations, i.e., the edge magnetoplasmons (EMP). The nonlocal effect arises at low values of the length l which is shown to govern the spatial extent of the plasma charge oscillations in the direction perpendicular to the q direction, i.e., to the edge of a 2DES.

The EMP frequency ω_{EMP} has been calculated by different authors [1, 2] and is shown to be essentially proportional to the Hall conductivity σ_{xy} and the wavevector q :

$$\omega_{\text{EMP}} = (2q\sigma_{xy}/\epsilon) \ln(2e/ql), \quad (1)$$

where ϵ is the effective dielectric function of the surrounding medium. The interesting quantity in the logarithmic term is l which in [1] is

$$l = 2\pi\sigma_{xx}/\epsilon\omega, \quad (2)$$

where σ_{xx} is the diagonal conductivity. However no significant influence of the logarithmic term with l has ever been observed in the previous experiments [3].

We have studied the EMP in a GaAs-AlGaAs heterostructure with electron density $n = 3 \cdot 10^{11} \text{ cm}^{-2}$ and mobility $\mu = 2 \cdot 10^5 \text{ cm}^2/\text{V s}$ in a non-resonant radio-frequency (1–1000 MHz) measurement cell [3].

The filling factor dependence $\omega_{\text{EMP}}(\nu)$ (for an arrangement with a large ϵ) demonstrates a very unusual behaviour. ω_{EMP} decreases and strongly oscillates with ν while the EMP damping γ_{EMP} has correspondingly deep minima. With increasing

temperature this behaviour becomes less pronounced. In the temperature regime where we observe in dc transport measurements an activated conductivity, that at $\nu = 2$ and $\nu = 4$ the frequencies ω_{EMP} exhibit a linear dependence on the inverse temperature $1/T$.

These novel results clearly demonstrate that EMP excitations are governed not only by σ_{xy} but also strongly influenced by σ_{xx} . Moreover, we have shown that ω_{EMP} depends on σ_{xx} via a logarithmic factor as follows from (1).

Finally we would like to discuss the relevant length l in our experiments. With decreasing T the plasmon "width", i.e., l , decreases as coincides with (2). However at low T ($T \leq 2K$) l approaches a saturation value which is extremely small and is $\approx 0.05 \mu\text{m}$ at $\nu = 1$ and $\nu = 2$ while the magnetic length is $\approx 0.01 \mu\text{m}$. That means that in this regime of the strong localization of the EMP we have reached the limits of the local approximation for $\sigma_{\alpha\beta}$ which is used in [1]. The length l now characterizes the non-local effects which govern the dynamical transport in the edge region of a 2DES in the QHE. We can suggest that in this case l corresponds to the variation of the electron density near the edge of a 2DES or a localization length in the QHE.

REFERENCES

- [1] V. A. VOLKOV and S. A. MIKHAILOV, Sov. Phys. JETP. 67, 1639 (1988).
- [2] M. WASSERMEIER et al., Phys. Rev. B41, 10287 (1990).
- [3] I. M. GRODNENSKY and A. YU. KAMAEV, Surf. Sci. 229, 522 (1990).