

**Zeitschrift:** Botanica Helvetica  
**Herausgeber:** Schweizerische Botanische Gesellschaft  
**Band:** 107 (1997)  
**Heft:** 1

**Artikel:** Response of the carbohydrate metabolism and fructose-2,6-bisphosphate to environmental changes : effect of different light treatments  
**Autor:** Praag, Esther van / Monod, David / Greppin, Hubert  
**DOI:** <https://doi.org/10.5169/seals-72640>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 11.07.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## Response of the carbohydrate metabolism and fructose-2,6-bisphosphate to environmental changes. Effect of different light treatments

Esther van Praag, David Monod, Hubert Greppin, Robert Degli Agosti

Laboratoire de Biochimie et Physiologie Végétale, Pavillon des Isotopes, 20, bd d'Yvoy, CH-1211 Genève 4, Switzerland

Manuscript accepted January 29, 1997

### Abstract

van Praag E., Monod D., Greppin H., Degli Agosti R. 1997. Response of the carbohydrate metabolism and fructose-2,6-bisphosphate to environmental changes. Effect of different light treatments. Bot. Helv. 107: 103–112.

Important and rapid variations of Fru-2,6-P<sub>2</sub> were observed during light-on and light-off transitions in both spinach and mustard: Fru-2,6-P<sub>2</sub> underwent tremendous variations showing an 80% decrease within 13 minutes in both plants. In mustard cotyledons, during both day and night periods, Fru-2,6-P<sub>2</sub> showed a decrease during the first hour, followed by a regular 240% increase. In spinach, the same behavior of Fru-2,6-P<sub>2</sub> was observed, except that once the concentration of the sugar had doubled, it remained constant. In contrast to mustard, spinach Fru-2,6-P<sub>2</sub> presented a much higher level during the night than during the light period. In 3 day old germinating oat, red light illumination for 5 min was accompanied by a 48 h halt of elongation and by a 180% increase of Fru-2,6-P<sub>2</sub> 20 min after irradiation, followed by the total disappearance of this sugar 80 minutes after light treatment.


The present work shows the highly dynamic capacity of the regulatory sugar Fru-2,6-P<sub>2</sub> in plants with respect to short, mid and long term light treatments. This indicates an important cellular messenger role of Fru-2,6-P<sub>2</sub> in the regulation of carbohydrate metabolism in response to light changes.

**Key words:** *Spinacia oleracea*, *Sinapis alba*, *Avena sativa*, Fructose-2,6-bisphosphate, carbohydrate metabolism, photoperiodism.

### Introduction

During the 80's, it was shown that Fru-2,6-P<sub>2</sub> plays a central role in sugar metabolism as it regulates the activities of several enzymes: those connected to glycolysis are generally activated, while those involved in gluconeogenesis are inhibited (Huber, 1986). Interactions of this molecule with its target enzymes could thus readjust the fluxes and metabolite pool in the cytosol (Stitt, 1990). The role of Fru-2,6-P<sub>2</sub> has become particularly important in plants since it regulates the activities of key enzymes: P<sub>i</sub>-dependent fruc-

tose-phosphate 1-phosphotransferase (PFP) (Sabulase and Anderson, 1981; van Schaftingen et al., 1982) and cytoplasmic fructose-1,6-bisphosphatase (cFBPase) (Stitt, 1990; Kruger and Beevers, 1984). PFP, is catalyzing the following interconversion:



This enzyme has a high affinity for its activator with a  $K_a$  generally situated between 1.56 nM and 50 nM (van Schaftingen et al., 1982; Bertagnolli et al., 1986; Degli Agosti et al., 1992; van Praag, 1996). It is present in the cytosol of plant cells as either  $\alpha_2$  or  $\beta_2$  homodimers or as a  $\alpha_2\beta_2$  heterotetramer. The  $\alpha$ -subunit is suggested to play a regulatory role by binding Fru-2,6-P<sub>2</sub>, while the  $\beta$ -subunit contains the catalytic activity (Carlisle et al., 1990). Recent computer models of the tetrameric  $\alpha_2\beta_2$  form of PFP confirm that Fru-2,6-P<sub>2</sub> binds between the  $\alpha$ - and  $\beta$ -subunits, with interactions between the 6'-phosphoryl group and the oxygen of the furanose ring with the  $\alpha$ -chain (van Praag, 1996).

Fru-2,6-P<sub>2</sub> is generally present in micromolar quantities in plant tissues with concentrations ranging between 1 and 30  $\mu$ M in spinach (Cséke et al., 1982; Stitt, 1990), maize (Stitt, 1990) or wheat leaves (Stitt, 1990) as well as in citrus juice cell protoplasts (van Praag, 1996). The level of this phosphorylated sugar further seems to follow an oscillating pattern in plant tissues (Stitt, 1990; Paz et al., 1985). This is rather interesting as Fru-2,6-P<sub>2</sub> generally correlates negatively with the accumulation of sucrose in photosynthetic leaves (Stitt, 1990). Such accumulation of sucrose during the day was shown in primary leaves and petioles of spinach plants (Steiner, 1968a), followed by a drop of sugar content during the night (dark period) (Steiner, 1968b). Interestingly, a regular increase in the concentrations of glucose and fructose but not of sucrose was observed in plants exposed to a prolongation of the light period (transfer treatment) (Degli Agosti et al., 1990). The same response was also noticed in etiolated mustard hypocotyls exposed to five minutes of red light. Indeed, the levels of free hexoses increased continuously in mustard hypocotyls from the moment they were shortly illuminated with red light (Steiner, 1968a, b), but it was concluded that it was an indirect consequence of irradiation.

In an effort to understand the mechanism that induce the accumulation of glucose and fructose in plant cells exposed to a sudden modification of the light regime, we decided to examine the possible involvement of Fru-2,6-P<sub>2</sub> in spinach or mustard leaves and in etiolated oat germs.

## Materials and methods

### Plant material

Spinach (*Spinacia oleraceae* L. cv. Nobel) plants, grown on soil, and mustard seeds (*Sinapis alba* L.), grown on vermiculite, were cultivated in a growth chamber with standard irradiation conditions

( $20 \text{ W m}^{-2}$ ) provided by Sylvania fluorescent lamps ("day-light" F40T12, 40 W). The temperature was maintained at  $20 \pm 0.5^\circ\text{C}$  and the relative humidity varied between  $70 \pm 5\%$  during the day and  $50 \pm 5\%$  at night. Plants were grown in a short day photoperiod: 8 hours light and 16 hours darkness. The transfer experiments were done with 30 day old spinach plants and 10 day old mustard plants.

Oat seeds (*Avena sativa* L. cv. Garry) were planted on wet vermiculite in closed boxes, which were kept in complete darkness for a number of days. Temperature of germination was  $21 \pm 1^\circ\text{C}$ .

#### *Extraction of Fru-2,6-P<sub>2</sub>*

Extraction was done at  $0^\circ\text{C}$ . The tissue samples were transferred into a cold grinding glass, 1 ml of cold NaOH 0.5 M and 3 ml of chloroform were added before extraction. After grinding extensively, the extract was vortexed and centrifuged at  $4000 \times g$  during 5 min. The aqueous phase was collected and stored at  $-80^\circ\text{C}$ . Recovery of added Fru-2,6-P<sub>2</sub> in control samples was 98.8%.

#### *Measurement of Fru-2,6-P<sub>2</sub>*

The possible interaction of other plant constituents can modify the activity of PFP and its response to Fru-2,6-P<sub>2</sub>. It is thus essential that activation of Fru-2,6-P<sub>2</sub> is calibrated by comparison with an internal standard curve using pure Fru-2,6-P<sub>2</sub> plus the extracts. The standard assay medium (1 ml) contained: 50 mM Hepes-NaOH, pH 7.5; 2 mM Mg-acetate; 1 mM Fru-6-P; 10 mM Fru-2,6-P<sub>2</sub>; 0.15 mM NADH; 0.45 U aldolase (EC 4.1.2.13); 5 U triose-phosphate isomerase (EC 5.3.1.1); 1.7 U glycerol-3-phosphate dehydrogenase (EC 1.1.1.8) and 0.01 U of potato PFP. The mixture was preincubated in a water bath for 5 min at  $25^\circ\text{C}$  and the reaction was initiated upon addition of PPI (final concentration: 0.5 mM) and 10 to 20  $\mu\text{l}$  of sample to the cuvette. Oxidation of NADH was followed in a spectrometer at 340 nm.

## Results

### *1. Fru-2,6-P<sub>2</sub>: fast and medium term response to changeable photoperiods*

The levels of Fru-2,6-P<sub>2</sub> were followed in mustard and spinach leaves when light was turned on and off (Fig. 1). Rapid sampling revealed that the level of this sugar showed tremendous changes in concentration within a short period of time. Indeed, some time before the light was switched on in the morning, the level of Fru-2,6-P<sub>2</sub> showed a 2.5 fold increase in mustard cotyledons, followed by a rapid disappearance at the moment the light was turned on (Fig. 1 A). Similar results were obtained for spinach leaves (Fig. 1 C). In both cases, the responses to light transition were extremely fast: a 77% decrease of Fru-2,6-P<sub>2</sub> within 9 min in mustard cotyledons and a 80% decrease within 13 min in spinach leaves.

After the photoperiod, when the light was switched off again, comparable fluctuations were noted in mustard (Fig. 1 B) and in spinach leaves (Fig. 1 D). In mustard tissue, the level of Fru-2,6-P<sub>2</sub> increased instantly within 3–4 minutes after nightfall, followed by a sharp decline and recovery. A similar increase after darkness was observed in spinach leaves, however, the level of Fru-2,6-P<sub>2</sub> stayed high when plants remained in darkness (Fig. 1 D).

Fru-2,6-P<sub>2</sub> concentrations were further followed over longer periods of time, during the light and dark phases of the short day photoperiod. In mustard cotyledons (Fig. 2), a decrease of the sugar diphosphate content occurs during the first hour of light, followed by a regular increase of 233% during the next 5 hours. Two hours before nightfall, the level of Fru-2,6-P<sub>2</sub> started to decline (Fig. 2).

During the dark period, the pattern of Fru-2,6-P<sub>2</sub> is rather similar to that obtained during the light period. During the first 60 min a strong decrease is observed, after which it increases with peak-concentrations every two hours in the case of mustard cotyledons.




Fig. 1. Changes in the level of Fru-2,6-P<sub>2</sub> in mustard cotyledons (A + B) and spinach leaves (C + D) kept at a short day photoperiod during the dark/light (A + C) and light/dark (B + D) transition periods.




Fig. 2. Variation in the content of Fru-2,6-P<sub>2</sub> in mustard cotyledons during a short day photoperiod (8:16 L:D) in light (○) and dark (●).

The concentrations of Fru-2,6-P<sub>2</sub> are comparable to those reached during the day (0.5 nmole/mg chlorophyll (CHY)), representing a 240% increase.

Similar results were obtained with spinach leaves (Fig. 3). However, the level of Fru-2,6-P<sub>2</sub> reached during the night is almost twice that reached after 4 hours of illumination (Fig. 3).

## 2. Germinating oat: growth, effect of light and concentrations of Fru-2,6-P<sub>2</sub>

It was shown that day light changes induce a rapid response in Fru-2,6-P<sub>2</sub> levels in leaves, but could 5 min of red light illumination influence the concentration of this sugar in 3 day old etiolated oat? The mean concentration of Fru-2,6-P<sub>2</sub> in the germs exposed to green light before irradiation is equal to  $0.34 \pm 0.02$  nmole/gFW. Twenty minutes after irradiation, the level of Fru-2,6-P<sub>2</sub> increased by 180% (Fig. 4), followed by a steady decrease till it cannot be detected anymore after 80 minutes.

The growth of the irradiated germs was further followed to check if a 5 min irradiation was long enough to induce morphological modifications in 3 day old etiolated germs. At this age, the epicotyl is about 10 mm long (Fig. 5C). Twenty four hours later, clear differences in growth were observed. Indeed the germs grown in complete darkness and green light were 2.5 times longer than those exposed to 5 min of light (Fig. 5). However, growth of the irradiated samples resumed after 48 hour as in those who had remained in darkness (Fig. 5).

The concentrations of Fru-2,6-P<sub>2</sub> has also been examined in growing germs to study long term fluctuations (Fig. 6). The concentration of Fru-2,6-P<sub>2</sub> was followed from the moment that the germ appears on the 3rd day up to the 6th day and it showed large differences: up to 0.65 nmole/gFW on the 4th and 6th day and 0.18 nmole/gFW the 5th day (Fig. 6). At this stage the roots are well developed (Fig. 5C). After the fast drop of




Fig. 3. Variation in the content of Fru-2,6-P<sub>2</sub> in spinach leaves during a short day photoperiod (8:16 L:D) in light (○) and dark (●).




Fig. 4. Variation in the content of Fru-2,6-P<sub>2</sub> in germinating oat (3 day old) grown in complete darkness (●), after receiving 5 min of red light on the 3rd day.




Fig. 5. Growth of germinating oat kept in complete darkness (●) and after irradiation with red light during 5 min on the 3rd day (○). A: oat seed after a 24 hour period of soaking. B: appearance of the germ, 48 hours after sowing. C: growth of the oat germ and appearance of the root at the age of 72 hours. D: the developing germs separates into hypocotyl and primary leaf. The roots are well developed and present absorbing hairs.






Fig. 6. Long term evolution of Fru-2,6-P<sub>2</sub> content in germinating oat grown in complete darkness (●) and moved to a short day photoperiod at the age of 6 days (○).

Fru-2,6-P<sub>2</sub> observed in the beginning of the 6th day, the etiolated oat germs were placed in a short day-light photoperiod. Till the 8th day, the level of Fru-2,6-P<sub>2</sub> remains low, with an average value of 0.15 nmole/gFW. After the elevation observed on the 9th day, Fru-2,6-P<sub>2</sub> seems to reach a plateau-concentration (Fig. 6).

## Discussion

In this work, it was shown that light is a very important environmental factor as oat germs, spinach leaves or mustard primary leaves modify their endogenous amount of Fru-2,6-P<sub>2</sub> in its presence.

### *Levels of Fru-2,6-P<sub>2</sub> in leaves during the day and the night photoperiod*

Measurement of Fru-2,6-P<sub>2</sub> content in spinach and mustard leaves show that the levels vary according to the day and night phases. While it took 6 hours to reach the maximum level of Fru-2,6-P<sub>2</sub> during the light period, it took only 5 hours to reach this value in the dark period in mustard leaves. In spinach leaves, the level of Fru-2,6-P<sub>2</sub> was immediately much higher than those noticed during the light period. During the periods of light changes (light-on and light-off), the level of Fru-2,6-P<sub>2</sub> underwent tremendous variations within minutes. Such rapid fluctuations were also observed in other plant tissues (Baysdorfer and Robinson, 1985; Fahrendorfer et al., 1987; Sicher et al., 1986; Stitt et al., 1984; Usuda et al., 1987). The results obtained for the Fru-2,6-P<sub>2</sub> concentrations after light-off (Fig. 1 B and 2) seem contradictory. From different experiments with short sampling intervals (data not shown) it became clear that the level of Fru-2,6-P<sub>2</sub> regularly decreased to a threshold concentration after a maximum was reached during



the photoperiod. Generally this minimal concentration was attained before the moment of light-off (see e.g. Fig. 3). However, if light-off occurs before the minimal threshold is reached (see Fig. 2, minimum of Fru-2,6-P<sub>2</sub> around 17 h), the short term fluctuations are present while the mid-term adaptation is delayed.

These results, combined with those obtained previously for hexose, sucrose and starch in the same plants (Degli Agosti and Greppin, 1987; Degli Agosti et al., 1989) seem to suggest that carbohydrate metabolism is regulated by endogenous factors that are synchronized with the photoperiod. One of those factors could be related to the accumulation of triose-phosphates in the cytosol of plant cells. These sugars are exported from the chloroplast into the cytosol of photosynthetic tissues. Semiquantitative models were developed to understand the role of Fru-2,6-P<sub>2</sub> in the regulation of the glycolysis/gluconeogenesis (Stitt, 1990). It was found that triose-P needs to accumulate in the cytosol, until a "threshold" is reached to activate cFBPase. The sensitive response to rising triose-P is generated as the concentration of Fru-2,6-P<sub>2</sub> increases and that of Fru-2,6-P<sub>2</sub> decreases. The last compound is indeed known to regulate the activity of cytosolic FBPase, which catalyses the irreversible dephosphorylation of Fru-1,6-P<sub>2</sub> into Fru-6-P. The enzyme is totally inhibited in presence of micromolar concentrations of Fru-2,6-P<sub>2</sub>. This value is nearby an order of magnitude higher than the activation constant of PFP. Assuming that the cytosol represents 4 to 10% of the cell volume, the higher levels of Fru-2,6-P<sub>2</sub> measured in oat germs, mustard or spinach leaves are thus high enough to activate PFP and completely inhibit cFBPase (van Schaftingen et al., 1982; Degli Agosti et al., 1992; van Praag, 1996; Stitt, 1990).

#### *Mid and long term response of Fru-2,6-P<sub>2</sub> in germinating oat*

It is known that red light affects numerous physiological and morphological processes in plants: e.g. a reduction of stem growth. During the post-germinative stages of oat seeds, an important regulator of carbohydrate metabolism, Fru-2,6-P<sub>2</sub>, was present at high concentrations in oat germs kept dark. The high level of this constituent can be compared with those measured in previous experiments (Sicher and Bunce, 1987) and agrees with the needs of the growing plant: sugars stored in the seeds are transported to the growing tissues where they are metabolized through the glycolytic pathway. Glucose, fructose and triose-phosphates are produced and enable the biosynthesis of additional molecules needed for the growth process. Decreased levels of Fru-2,6-P<sub>2</sub> on the 5th day correspond to the stage where more than 60% of the starch stored in the seed has been exported to the growing tissues (Degli Agosti et al., 1991). This stage also shows already well developed roots, that should be able to take up the necessary minerals from the soil. Transition from sink tissue to source tissue can thus be expected to take place, enabling the etiolated germs to start gluconeogenesis, a metabolic pathway that can only be performed when the Fru-2,6-P<sub>2</sub> level is low.

The level of Fru-2,6-P<sub>2</sub> present in the germs that were placed in a SD photoperiod corresponds to those measured in mustard and spinach leaves. The greening of the coleoptile is accompanied by the development of chloroplasts. Once photosynthesis takes place, the chloroplasts start to export triose-phosphates to the cytosol and sucrose can be formed through gluconeogenesis.

The results obtained after exposing 3 day old etiolated germs to 3 min of red light show that the amount of Fru-2,6-P<sub>2</sub> is rapidly affected. Indeed, a sudden increase is noticed 20 min after irradiation, followed by total disappearance 60 min later. This dramatic effect of the Fru-2,6-P<sub>2</sub> content, accompanied by the increase of glucose and

fructose (Steiner, 1968 a, b) could possibly indicate that Fru-2,6-P<sub>2</sub> acts as an initial signal for further metabolic events that help the plant cell to reorganize itself in order to adapt to the new environmental conditions. The rapid fluctuations of Fru-2,6-P<sub>2</sub> could indeed be the signal that triggers the dissociation of PFP from its tetrameric glycolytic form into its gluconeogenic dimer. This hypothesis is strengthened by the fact that almost no increase of reducing sugars was observed in germs placed in continuous FR light after the R irradiation (Steiner, 1968 a, b).

The results further show that some specific parameters (time of the day, rate of sampling) need to be respected in experiments dealing with measurements of Fru-2,6-P<sub>2</sub> levels in plant tissues. Particularly in long term studies, it is important to collect the daily samples at exactly the same time of the day to be able to compare the evolution of the Fru-2,6-P<sub>2</sub> levels of this specific moment. Studies dealing with the Fru-2,6-P<sub>2</sub> levels during the light and night periods in plants will, however, only indicate the general tendency of the effector molecule, as it is shown in the present study that its concentrations can vary considerably in very short periods of time.

## Resumé

Le Fru-2,6-P<sub>2</sub>, régulateur du métabolisme des sucres, présente des modifications de sa concentration à court, moyen et long terme chez diverses plantes lors de variations lumineuses. Il fluctue de manière journalière dans les cotylédons de moutarde et les feuilles d'épinard. Lors des phases lumineuses, on observe une diminution durant les premières heures, suivie d'une augmentation régulière d'environ 240%. On observe également des fluctuations importantes et rapides au moment des transitions nuit/jour et jour/nuit chez ces deux plantes. L'irradiation par 5 min de lumière rouge sur des germes d'avoine étiolés modifie aussi bien la croissance que le niveau de Fru-2,6-P<sub>2</sub>. Celui-ci augmente de 180% 20 min. après le traitement et diminue progressivement par la suite.

Les résultats obtenus montrent que le Fru-2,6-P<sub>2</sub> possède une remarquable capacité dynamique d'adaptation à court, moyen et long terme. Ces adaptations sont présentes aussi bien chez des dicotylédones (moutarde et épinard) que chez une monocotylédone (avoine). Ceci met en évidence le rôle important joué par le Fru-2,6-P<sub>2</sub> en tant que messenger cellulaire du métabolisme des sucres en réponse à diverses variations lumineuses de l'environnement.

## References

- Baysdorfer C. and J. M. Robinson 1985. A rapid increase in spinach leaf fructose 2,6-bisphosphate occurs during a light to dark transition. *Plant Physiol.* 79: 911–913.
- Bertagnolli B. L., Younathan E. S., Voll R. J., and Cook P. F. 1986. Kinetic studies on the activation of pyrophosphate-dependent phosphofructokinase from mung bean by fructose 2,6-bisphosphate and related compounds. *Biochemistry* 25: 4682–4687. Waverley Press, Baltimore, pp. 44–62.
- Carlisle S., Blakeley S. D., Hemmingen S. M., Trevanion S. J., Hiyoshi T., Kruger N. K., Cséke C., Weeden N. F., Buchanan B. B., and Uyeda K. 1982. A special fructose bisphosphate functions as a cytoplasmic regulatory metabolite in green leaves. *Proc. Natl. Acad. Sci. USA* 79: 4322–4326.
- Degli Agosti R. and Greppin H. 1987. Extraction, identification et dosage des sucres du pétiole le l'épinard (*Spinacia oleracea* L. cv. Nobel) pendant la variation de photopériode. *Bot. Helv.* 97: 329–340.

- Degli Agosti R., Bonzon M., and Greppin H. 1989. Photoperiodic floral induction and glucose content changes in spinach, mustard and *Chenopodium rubrum* plants. Bot. Helv. 99: 73–79.
- Degli Agosti R., Bonzon M., and Greppin H. 1990. Carbohydrates evolution in spinach plants during early acclimation to various changes in photoperiod. Archs. Sci. Genève 43: 401–417.
- Degli Agosti R., Lenk R., and Greppin H. 1991. Etude par la résonance magnétique nucléaire de la germination d'une céréale. Archs. Sci. Genève 44: 235–243.
- Degli Agosti R., van Praag E., and Greppin H. 1992. Effect of chloride ions on the kinetic parameters of the potato tuber and mung bean pyrophosphate-dependent phosphofructokinase. Biochem. Int. 26: 707–713.
- Fahrendorfer T., Joseph A., Holtum M., Mukherjee U., and Latzko E. 1987. Fructose 2,6-bisphosphate partitioning and Crassulacean acid metabolism. Plant Physiol. 84: 182–187.
- Huber S. C. 1986. Fructose 2,6-bisphosphate as a regulatory molecule in plants. Annu. Rev. Plant Physiol. 37: 122–146.
- Kruger N. J. and Beevers M. 1984. Effect of fructose 2,6-bisphosphate on the kinetic properties of cytoplasmic fructose 1,6-bisphosphatase from germinating castor endosperm. Plant Physiol. 76: 49–54.
- Paz N., Xu D. P., and Black C. C. Jr. 1985. Rapid oscillations of fructose-2,6-bisphosphate levels in plant tissues. Plant Physiol. 79: 1133–1136.
- van Praag E. 1996. Fructose 2,6-bisphosphate: its role in the regulation of carbohydrate metabolism in Citrus fruits. Ph.D. The Hebrew University of Jerusalem, Israel.
- Sabulase D. C. and Anderson R. L. 1981. Inorganic pyrophosphate: fructose-6-phosphate 1-phosphotransferase in mung beans and its activation by fructose 1,6-bisphosphate and D-glucose 1,6-bisphosphate. Biochem. Biophys. Res. Comm. 100: 1423–1429.
- van Schaftingen E., Lederer B., Bartrons R., and Hers H. G. 1982. A kinetic study of pyrophosphate: fructose 6-phosphate phosphotransferase from potato tubers. Eur. J. Biochem. 129: 191–195.
- Sicher R. C., Kremer D. F., and Harris W. G. 1986. Control of photosynthetic sucrose synthesis in barley primary leaves. Role of fructose 2,6-bisphosphate. Plant Physiol. 82: 15–18.
- Sicher R. C. and Bunce J. A. 1987. Effects of light and CO<sub>2</sub> on fructose 2,6-bisphosphate levels in barley primary leaves. Plant. Physiol. Biochem. 25: 525–530.
- Steiner A. M. 1968 a. Rasch ablaufende Änderungen im Gehalt an löslichen Zuckern und Zellwandkohlenhydraten bei der phytochrominduzierten Photomorphogenese des Senfkeimlings (*Sinapis alba* L.) Planta 82: 223–234.
- Steiner A. M. 1968 b. Änderungen im Kohlenhydratgehalt der Organe des Senfkeimlings (*Sinapis alba* L.) im Dauer-Dunkelrot unter dem Einfluß von Phytochrom. Z. Pflanzenphysiol. 59: 401–414.
- Stitt M., Herzog B., and Heldt H. W. 1984. Control of photosynthetic sucrose synthesis by fructose 2,6-bisphosphate. I. Coordination of CO<sub>2</sub> fixation and sucrose synthesis. Plant Physiol. 75: 548–553.
- Stitt M. 1990. Fructose 2,6-bisphosphate as a regularory molecule in plants. Annu. Rev. Plant Physiol. Plant Mol. Biol. 41: 153–185.
- Usuda H., Kalt-Torres W., Kerr P. S., and Huber S. C. 1987. Diurnal changes in maize-leaf photosynthesis. II. Levels of metabolic intermediates of sucrose synthesis and the regulatory metabolite fructose 2,6-bisphosphate. Plant Physiol. 83: 289–293.