

Zeitschrift: Commentarii Mathematici Helvetici
Band: 53 (1978)

Artikel: Cohomology eigenvalues of equivariant mappings.
Autor: Skjelbred, Tor
DOI: <https://doi.org/10.5169/seals-40792>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

Download PDF: 01.11.2024

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Cohomology eigenvalues of equivariant mappings

TOR SKJELBRED

Let X be a topological space which is paracompact Hausdorff and of finite cohomology dimension over a fixed field k . Let G be a compact Lie group acting continuously on X such that there is a finite number of conjugacy classes of isotropy groups G_x , $x \in X$. Conner conjectured in [2] that if $H^*(X; k)$ is acyclic, then $H^*(X/G; k)$ is also acyclic, and he proved the conjecture in case $k = \mathbb{Q}$. The conjecture was recently proven in all characteristics by Robert Oliver [8]. The problem of relating $H^*(X/G; k)$ and $H^*(X; k)$ is still largely unsolved even in case X is the unit sphere of a linear representation. In this paper we will consider equivariant mappings $f: X \rightarrow X$ and relate the eigenvalues of the induced endomorphisms of $H^*(X/G; k)$ and of $H^*(X; k)$. The result obtained should be seen as a generalization of the Conner conjecture to G -spaces which are not necessarily acyclic.

THEOREM 1. *Let f be an equivariant self-mapping of a G -space X . Then each eigenvalue of the induced endomorphism of $\tilde{H}^*(X/G; k)$ is an eigenvalue of the induced endomorphism of $\tilde{H}^*(X; k)$, provided $\dim_k H^*(X; k) < \infty$.*

More generally we consider the monoid $\text{Map}(G, X)$ of all equivariant mappings $X \rightarrow X$, and a homomorphism from a monoid \mathcal{F} into $\text{Map}(G, X)$. Then $H^*(X; k)$ and $H^*(X/G; k)$ become right \mathcal{F} -modules. Let M be an abelian group which is a right \mathcal{F} -module. A simple subquotient of the \mathcal{F} -module M is a simple \mathcal{F} -module isomorphic to M_2/M_1 where $M_1 \subset M_2 \subset M$ are \mathcal{F} -submodules. M may be a module over a field k and \mathcal{F} commuting with k . Even if M is not finitely generated, the following lemma is straightforward.

LEMMA 1. *Let*

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$$

be an exact sequence of \mathcal{F} -modules. Then a simple \mathcal{F} -module is a subquotient of M if and only if it is a subquotient of $M' \oplus M''$.

Our main result then is,

THEOREM 2. *Let X be a G -space and let \mathcal{F} be a monoid of equivariant self-mappings of X . Then every simple subquotient of the \mathcal{F} -module $\tilde{H}^*(X/G; k)$ is a simple subquotient of the \mathcal{F} -module $\tilde{H}^*(X; k)$. If $Y \subset X$ is a closed subspace invariant under G and under all $f \in \mathcal{F}$, then every simple subquotient of the \mathcal{F} -module $H^*(X/G, Y/G; k)$ is a simple subquotient of the \mathcal{F} -module $H^*(X, Y; k)$.*

This result may be interpreted in terms of Serre classes of \mathcal{F} -modules. Let N be a simple \mathcal{F} -module over k . Then by Lemma 1, those \mathcal{F} -modules which do not have N as a subquotient form a Serre class, say C_N . Theorem 2 says that if $\tilde{H}^*(X; k)$ belongs to C_N , then so does $\tilde{H}^*(X/G; k)$. It is then a Conner conjecture modulo the Serre class C_N . If we forget equivariant mappings and consider the Serre class of finitely generated abelian groups, we obtain,

THEOREM 3. *Let X be a G -space, and assume that X has finite cohomology dimension over \mathbf{Z} . Then if $H^*(X; \mathbf{Z})$ is finitely generated, so is $H^*(X/G; \mathbf{Z})$.*

We use Čech cohomology with closed supports. We use some results on cohomology dimension [9] and the localization theory of Borel-Segal-Hsiang-Quillen [1, 6, 9, 10] without further comments. When G is finite or abelian, the proof of Theorems 1–3 is based on the localization theory. When G is connected simple, the proof is based on the Conner conjecture and on the existence of the spheres of Floyd-Hsiang [3, 5]. We first simplify the group G .

LEMMA 2. (i) *Let $N \subset G$ be a closed normal subgroup such that Theorem 2 holds for actions of N and of G/N . Then Theorem 2 holds for actions of G .*

(ii) *It suffices to prove Theorem 2 when G is either a finite group of prime order, a circle group acting semifreely, or a simple connected Lie group.*

Proof. (i) Let \mathcal{F} be a monoid of equivariant self-mappings of the G -space X . There is a natural homomorphism $\mathcal{F} \rightarrow \text{Map}(G/N, X/N)$, and hence every simple subquotient M of the \mathcal{F} -module $\tilde{H}^*(X/G; k) = \tilde{H}^*((X/N)/(G/N); k)$ is a subquotient of $\tilde{H}^*(X/N; k)$. Because $\mathcal{F} \subset \text{Map}(G, X) \subset \text{Map}(N, X)$, and Theorem 2 holds for actions of N , the simple module M must be a subquotient of $\tilde{H}^*(X; k)$. Hence Theorem 2 holds for the G -action on X .

(ii) By (i) we may assume that G is a finite group, a circle group, or a connected simple group. If $G = SO(2)$, let $Z \subset G$ be a finite subgroup containing all finite isotropy groups. Then the action of G/Z on X/Z is semifree. By (i), it suffices to prove Theorem 2 for actions of cyclic groups and for semifree circle

actions to give a proof for all circle actions. If G is finite, let S be the p -Sylow subgroup of G , where $p = \text{char}(k)$, and where $S = \{1\}$ if $p = 0$. Then by [1] (p. 38), we have $H^*(X/G; k) \subset H^*(X/S; k)$. Therefore, it suffices to prove Theorem 2 for the group S . Because S is solvable, it follows from (i) that we can reduce the problem to finite groups of prime order.

Proof of Theorem 2 for G connected simple.

We shall construct a compact G -space Z such that for each closed subgroup H of G the orbit mapping $Z \rightarrow Z/H$ induces an isomorphism

$$H^*(Z/H; \mathbf{Z}) \xrightarrow{\cong} H^*(Z; \mathbf{Z}).$$

Z is a compact G -CW complex in the sense of Matumoto [7], and G has no fixed points in Z . We construct Z by using,

THEOREM (Floyd-Hsiang [3, 5]) *Each simple connected compact Lie group G admits a real linear representation without one-dimensional direct summands such that the unit sphere admits an equivariant self-mapping of degree 0.*

Let S be the unit sphere, and $n : S \rightarrow S$ an equivariant self-mapping of degree 0. Let $Z = T(n)$ be the mapping torus of n , that is the space obtained from $S \times [0, 1]$ by identifying $(x, 1)$ with $(n(x), 0)$ for $x \in S$. Let $\pi : T(n) \rightarrow S^1$ be the projection on the second factor where $S^1 = [0, 1]/\{0, 1\}$. $T(n)$ is a G -CW complex because n is constructed by extending a piecewise linear map of a fundamental domain into the fixed point set of a principal isotropy group, where the simplicial structure is compatible with the orbit type stratification. (This is actually done for an action of some $SO(2r+1)$ on S , and the action is restricted to G by a representation of G of degree $2r+1$. This construction is found in [3, 5] and with more details in [11].) $T(n)$ is a G -space in a natural way such that the fibres $\pi^{-1}(z)$, $z \in S^1$, are canonically G -homeomorphic to S . Since n is nullhomotopic, it follows that π is a homotopy equivalence, and hence that the mapping cone $C(\pi)$ of π is contractible. Since $C(\pi)$ is a finite CW complex, the Conner conjecture, proved by Oliver, implies that $H^*(C(\pi)/H; \mathbf{Z}) = \mathbf{Z}$ for each closed subgroup H of G . Clearly $C(\pi)/H$ is the mapping cone of $T(n)/H \rightarrow S^1$, and hence

$$H^*(T(n)/H; \mathbf{Z}) \simeq H^*(S^1; \mathbf{Z}) \simeq H^*(T(n); \mathbf{Z}).$$

The G -CW structure on Z defines a finite cell complex structure on Z/G ([7]).

For each cell c of Z/G , choose $x \in Z$ such that $G(x)$ is in the interior of c , and set $G_c = G_x$. The cellular system (G_c) will be used in the Borel construction. Given two G -spaces X and Z , we consider $Z \times X$ as a G -space with the diagonal (joint) action, and there are projections of orbit spaces,

$$pr_1 : (Z \times X)/G \rightarrow Z/G, \quad pr_2 : (Z \times X)/G \rightarrow X/G.$$

The fibres of pr_1 and pr_2 are, for $x \in X, z \in Z$,

$$pr_1^{-1}(G(z)) = (G(z) \times X)/G = X/G_z$$

and

$$pr_2^{-1}(G(x)) = (Z \times G(x))/G = Z/G_x.$$

We apply the Leray spectral sequence to the mappings pr_2 and p_2 of the following commutative diagram where the vertical arrows are induced by π .

$$\begin{array}{ccccc} Z/G_x & \rightarrow & (Z \times X)/G & \xrightarrow{pr_2} & X/G \\ \downarrow \pi & & \downarrow & & \downarrow 1 \\ S^1 & \rightarrow & S^1 \times (X/G) & \xrightarrow{p_2} & X/G \end{array}$$

Here pr_2 and p_2 are proper mappings. Since π induces cohomology isomorphisms of the fibres, we have

$$H^*(S^1) \otimes H^*(X/G) \cong H^*((Z \times X)/G)$$

for any coefficient ring. This clearly is an isomorphism of \mathcal{F} -modules. For the mapping

$$pr_1 : (Z \times X)/G \rightarrow Z/G$$

we obtain a spectral sequence defined by the skeleton filtration of the cell complex Z/G , with

$$E_1 = C_{\text{cell}}^*(Z/G; \mathcal{H}^*(X/G_c))$$

and converging to $H^*((Z \times X)/G) \simeq H^*(S^1) \otimes H^*(X/G)$. For reduced cohomology, there is the spectral sequence \tilde{E} with $\tilde{E}_1 = C_{\text{cell}}^*(Z/G; \tilde{\mathcal{H}}^*(X/G_c; k))$ converging to $H^*(S^1) \otimes \tilde{H}^*(X/G; k)$. This is a spectral sequence of \mathcal{F} -modules. A simple subquotient of the \mathcal{F} -module $\tilde{H}^*(X/G; k)$ must be a simple subquotient of \tilde{E}_1 and hence of some $\tilde{H}^*(X/G_c; k)$. Because Z is without fixed points, $G_c < G$ for each c . By induction on $\dim G$, we may assume that Theorem 2 holds for actions of G_c . Hence each simple subquotient of $\tilde{H}^*(X/G_c; k)$ is a subquotient of $\tilde{H}^*(X; k)$, and this proves Theorem 2 for the given action of G . The proof for a closed pair (X, Y) of G -spaces is similar, using a spectral sequence converging to

$$H^*(S^1) \otimes H^*(X/G, Y/G; k)$$

with

$$E_1 = C_{\text{cell}}^*(Z/G; \mathcal{H}^*(X/G_c, Y/G_c; k)).$$

Proof of Theorem 2 for $G = \mathbf{Z}/p$ and $G = S^1$.

By Lemma 2, we may assume that G is acting semifreely. Let X_G be the Borel space of the G -action; it is the total space of a fibre bundle $X \rightarrow X_G \rightarrow B_G$ where B_G is the classifying space of principal G -bundles. We set $H_G^*(X) = H^*(X_G)$ and refer to [1, 6, 9] for the basic properties of this functor.

PROPOSITION 1. *Let G be a compact Lie group acting semifreely on a space X with fixed point set F . Then there is a long exact Mayer-Vietoris sequence of the form*

$$\dots \xrightarrow{\delta} H^q(X/G) \rightarrow H^q(F) \oplus H_G^q(X) \rightarrow H_G^q(F) \xrightarrow{\delta} \dots$$

Proof. Because the action is semifree and X is paracompact, there is an isomorphism

$$H^*(X/G, F) \rightarrow H_G^*(X, F)$$

induced by the projection $\pi: X_G \rightarrow X/G$, for any coefficient group. π induces, with its restriction to F_G , a homomorphism of long exact cohomology sequences,

$$\begin{array}{ccccccc}
 \dots & \xrightarrow{\delta} & H_G^*(X, F) & \rightarrow & H_G^*(X) & \rightarrow & H_G^*(F) \xrightarrow{\delta} \dots \\
 (*) & & \uparrow \pi^* & & \uparrow \pi^* & & \uparrow \pi^* \\
 \dots & \xrightarrow{\delta} & H^*(X/G, F) & \rightarrow & H^*(X/G) & \rightarrow & H^*(F) \xrightarrow{\delta} \dots
 \end{array}$$

The Mayer-Vietoris sequence is deduced from (*) by a standard argument, see p. 3 of [4]. Let P be a one-point space with its unique G -action. We set $\tilde{H}_G^*(X) = \text{coker}(H_G^*(P) \rightarrow H_G^*(X))$. There is then a reduced Mayer-Vietoris sequence if $F \neq \emptyset$,

$$(RMV) \cdots \xrightarrow{\delta} \tilde{H}^*(X/G) \rightarrow \tilde{H}^*(F) \oplus \tilde{H}_G^*(X) \rightarrow \tilde{H}_G^*(F) \xrightarrow{\delta} \cdots$$

LEMMA 3. Let $G = \mathbf{Z}/p$ or S^1 be acting semifreely on X with fixed point set $F \neq \emptyset$. Let \mathcal{F} be a monoid of equivariant self-mappings of X . Then every simple subquotient of any of the three \mathcal{F} -modules $\tilde{H}_G^*(X; k)$, $\tilde{H}_G^*(F; k)$, and $\tilde{H}^*(F; k)$ is a subquotient of the \mathcal{F} -module $\tilde{H}^*(X; k)$.

Proof. If k is of characteristic p , then $G = \mathbf{Z}/p$ or S^1 . Because $\tilde{H}_G^*(F; k) = \tilde{H}^*(F; k) \otimes H^*(B_G; k)$ and the restriction homomorphism $\tilde{H}_G^*(X; k) \rightarrow \tilde{H}_G^*(F; k)$ is surjective in high degrees, it follows that every simple subquotient of the \mathcal{F} -modules $\tilde{H}^*(F; k)$ and $\tilde{H}_G^*(F; k)$ is a subquotient of $\tilde{H}_G^*(X; k)$. The fibre bundle $X \rightarrow X_G \rightarrow B_G$ gives a spectral sequence converging to $\tilde{H}_G^*(X; k)$ with

$$E_1 = C_{\text{cell}}^*(B_G; \tilde{\mathcal{H}}^*(X; k)).$$

Hence every simple subquotient of the \mathcal{F} -module $\tilde{H}_G^*(X; k)$ is a simple subquotient of the \mathcal{F} -module $\tilde{H}^*(X; k)$.

COROLLARY 1. If $F \neq \emptyset$, then Theorem 2 holds for $G = \mathbf{Z}/p, S^1$.

Proof. The reduced Mayer-Vietoris sequence (RMV) shows that every simple subquotient of $\tilde{H}^*(X/G; k)$ is a subquotient of $\tilde{H}_G^*(F; k) \oplus \tilde{H}_G^*(X; k) \oplus \tilde{H}^*(F; k)$. By Lemma 3, it is a subquotient of the \mathcal{F} -module $\tilde{H}^*(X; k)$.

When $F = \emptyset$, $G = \mathbf{Z}/p$ or S^1 is acting freely, and there is an isomorphism $H^*(X/G; k) \simeq H_G^*(X; k)$. There is the spectral sequence of the fibring $X_G \rightarrow B_G$ with

$$\begin{aligned} E_1 &= C_{\text{cell}}^*(B_G; \mathcal{H}^*(X; k)), \\ E_2^{ab} &= H^a(\mathbf{Z}/p; H^b(X; k)) \quad \text{for } G = \mathbf{Z}/p, \text{ and} \\ E_2^{ab} &= H^a(\mathbf{CP}^\infty) \otimes H^b(X; k) \quad \text{for } G = S^1, \end{aligned}$$

and converging to $H^*(X/G; k)$. To prove Theorem 2 in this case, it suffices to show that every simple subquotient of the \mathcal{F} -module E_∞/k (where $k \subset E_\infty^{00}$ is the field of coefficients) is a subquotient of $\tilde{H}^*(X; k)$. Clearly, for $r \geq 1, b > 0$, every

simple subquotient of E_r^{ab} is a subquotient of $H^b(X; k)$. Hence, for $r \geq 2$, every simple subquotient of $d_r(E_r)$ is a subquotient of $H^+(X; k) = \sum_{b>0} H^b(X; k)$. For $a > c$, $c =$ the cohomology dimension of X over k , $E_\infty^{a0} = 0$. It follows that for $a > c$, each simple subquotient of E_2^{a0} is a subquotient of $H^+(X; k)$. As \mathcal{F} -modules, $E_2^{a0} \simeq E_2^{a+2o}$ for $a > 0$, and hence the last statement is valid for all $a > 0$. It remains only the module E_∞^{00}/k which is contained in $\tilde{H}^0(X; k)$, and the proof is complete for the case $F = \emptyset$.

The proof of Theorem 2 for a closed pair (X, Y) of G -spaces is quite similar to the proof in the absolute case with $F \neq \emptyset$. There is a Mayer-Vietoris sequence of a semifree group action,

$$\cdots \xrightarrow{\delta} H^*(X/G, Y/G) \rightarrow H^*(F, F \cap Y) \oplus H_G^*(X, Y) \rightarrow H_G^*(F, F \cap Y) \xrightarrow{\delta} \cdots$$

and there is a spectral sequence with

$$E_1 = C_{\text{cell}}^*(B_G; \mathcal{H}^*(X, Y; k)) \text{ converging to } H_G^*(X, Y; k).$$

This completes the proof of Theorem 2.

Next we give a proof of Theorem 3 which states that $H^*(X/G; \mathbf{Z})$ is finitely generated when $H^*(X; \mathbf{Z})$ is finitely generated. A preliminary result is,

PROPOSITION 2. *Let X be a G -space with a closed invariant subspace Y . Assume that X has finite cohomology dimension over a field k . Then if $H^*(X, Y; k)$ is finite dimensional over k , so is $H^*(X/G, Y/G; k)$.*

Proof. The proof is basically the same as the proof of Theorem 2, but with simplifications. Lemma 2 is valid for the present proof. If $G = \mathbf{Z}/p$ or S^1 acting semifreely, the proof is a direct consequence of the Mayer-Vietoris sequence of a semifree group action and the fact that the restriction homomorphism $H_G^*(X, Y; k) \rightarrow H_G^*(F, F \cap Y; k)$ is an isomorphism in high degrees. The exact sequence

$$\begin{aligned} \cdots \xrightarrow{\delta} H^*(X/G; Y/G; k) \rightarrow H^*(F, F \cap Y; k) \oplus H_G^*(X, Y; k) \\ \longrightarrow H_G^*(F, F \cap Y; k) \xrightarrow{\delta} \cdots \end{aligned}$$

then implies that $H^*(X/G; Y/G; k) \rightarrow H^*(F, F \cap Y; k)$ has finite dimensional kernel and cokernel. But $\dim_k H^*(F, F \cap Y; k) \leq \dim_k H^*(X, Y; k) < \infty$, and it

follows that $\dim_k H^*(X/G, Y/G; k) < \infty$. In case G is connected simple, we use the spectral sequence of the first part of the proof of Theorem 2 with

$$E_1 = C_{\text{cell}}^*(Z/G; \mathcal{H}^*(X/G_c, Y/G_c; k))$$

and converging to $H^*(S^1) \otimes H^*(X/G; Y/G; k)$. By induction on $\dim G$, we may assume that $\dim_k H^*(X/G_c, Y/G_c; k) < \infty$ for each cell c of Z/G . Since Z/G is a finite cell complex, it follows that $\dim_k E_1 < \infty$, and hence that $\dim_k H^*(X/G, Y/G; k) < \infty$. This completes the proof of Proposition 2.

THEOREM 3'. *Assume that a compact Lie group G is acting on a space X which is paracompact Hausdorff and has finite cohomology dimension (over \mathbf{Z}). Assume that there is a finite number of conjugacy classes of isotropy groups. Let Y be a closed invariant subspace. Then if $H^*(X, Y; \mathbf{Z})$ is finitely generated, so is $H^*(X/G, Y/G; \mathbf{Z})$.*

Proof. Again, the proof is basically the same as that of Theorem 2, with some changes for finite G . Let G be finite. Let $q: (X, Y) \rightarrow (X/G, Y/G)$ be the orbit mapping, and let $t: H^*(X, Y; \mathbf{Z}) \rightarrow H^*(X/G, Y/G; \mathbf{Z})$ be the transfer mapping ([1] p. 38). Then tq^* is multiplication by $m = |G|$ in $H^*(X/G, Y/G; \mathbf{Z})$, and hence, $\text{coker}(tq^*) \subset H^*(X/G, Y/G; \mathbf{Z}/m)$. Since tq^* factors through the finitely generated group $H^*(X, Y; \mathbf{Z})$, it suffices to show that $H^*(X/G, Y/G; \mathbf{Z}/m)$ is finitely generated. This is the case because, by Proposition 2, $H^*(X/G, Y/G; \mathbf{Z}/p)$ is finitely generated for each prime p . Now let G be a circle group. We may assume that G is acting semifreely, in which case the localization theory for circle actions is valid for cohomology with arbitrary coefficient group. Hence the argument in the proof of Proposition 2 is valid with integral coefficients. To prove Theorem 3' for general G , we may assume that G is connected, and that the theorem holds for all H with $\dim H < \dim G$, and hence that G is a connected simple group. Using the spectral sequence converging to $H^*(S^1) \otimes H^*(X/G, Y/G; \mathbf{Z})$, with $E_1 = C_{\text{cell}}^*(Z/G; \mathcal{H}^*(X/G_c, Y/G_c; \mathbf{Z}))$ where $\dim G_c < \dim G$, it follows that $H^*(X/G, Y/G; \mathbf{Z})$ is finitely generated.

Example. There is a pair (X, Y) of G -spaces and an equivariant mapping $f: (X, Y) \rightarrow (X, Y)$ such that a certain eigenvalue $\neq 1$ is of multiplicity one in $H^*(X/G, Y/G; k)$, and of multiplicity at least two in $H^*(X, Y; k)$. Let V be the linear space of all real n by n symmetric matrices of trace 0, and let X be the unit sphere in V . The group $SO(n)$ acts on X by conjugation with principal isotropy group $H \simeq (\mathbf{Z}/2)^{n-1}$. Let Y be the subspace consisting of all $x \in X$ such that G_x is not principal, equivalently such that $\dim G_x > 0$. In the author's paper

[11] there is constructed equivariant mappings $f_s : X \rightarrow X$ for $0 < 2s < n$, $n \geq 3$ of degrees $\deg f_s = 1 - \binom{m}{s}$ where $2m \leq n \leq 2m + 1$. Those mappings generalize the mapping f_m of Floyd-Hsiang, which is of degree 0 when $n = 2m + 1$. The mapping f'_s in the orbit space $\Delta = X/G$ is a self mapping of the orientable manifold-with-boundary Δ which is a simplex of dimension $n - 2$. In $H^*(\Delta, \partial\Delta; \mathbf{Z}) = \mathbf{Z}$ f'_s induces multiplication by $\deg f'_s$, and by Theorem (2.1) of [11], $\deg f'_s = \deg f_s = 1 - \binom{m}{s}$. It follows that in $\tilde{H}^*(\partial\Delta; \mathbf{Z}) \simeq \mathbf{Z}$, f'_s induces multiplication by $1 - \binom{m}{s}$. Because $\partial\Delta = Y/G$, Theorem 2 implies that, for each field k , $1 - \binom{m}{s}$ is an eigenvalue of $(f_s | Y)^*$ in $\tilde{H}^*(Y; k)$. From the exact sequence

$$0 \rightarrow \tilde{H}^*(Y; k) \xrightarrow{\delta} H^*(X, Y; k) \rightarrow \tilde{H}^*(X; k) \rightarrow 0$$

it follows that the eigenvalue $1 - \binom{m}{s}$ has multiplicity at least two in $H^*(X, Y; k)$, while it has multiplicity one in $H^*(X/G, Y/G; k) \simeq H^*(\Delta, \partial\Delta; k) \simeq k$.

REFERENCES

- [1] A. BOREL, *Seminar on transformation groups*, Princeton University press 1960.
- [2] P. E. CONNER, *Retraction properties of the orbit space of a compact transformation group*. Duke Math. J. 27 (1960) pp 341-357.
- [3] — and D. MONTGOMERY, *An example for $SO(3)$* . Proc. Nat. Acad. Sci. USA 48 (1962) pp. 1918-22.
- [4] E. DYER, *Cohomology Theories*, Benjamin 1969.
- [5] W. C. HSIANG and W. Y. HSIANG, *Differentiable actions of compact connected classical groups, I*. Amer. J. Math. 89 (1967) pp. 705-786.
- [6] W. Y. HSIANG, *Cohomology Theory of Topological Transformation Groups*. Ergebnisse der Mathematik, Band 85, Springer-Verlag 1975.
- [7] T. MATUMOTO, *On G -CW complexes and a theorem of J. H. C. Whitehead*. J. Fac. Sci. Univ. of Tokyo, 18 (1971) pp. 363-374.
- [8] R. A. OLIVER, *A proof of the Conner conjecture*. Ann. of Math. 103 (1976) pp 637-644.
- [9] D. QUILLEN, *The spectrum of an equivariant cohomology ring I*. Ann. of Math. 94 (1971) pp. 549-572.
- [10] G. B. SEGAL, *Equivariant K-theory*. Publ. Math. I. H. E. S. Paris 34 (1968) pp. 129-151.
- [11] T. SKJELBRED, *Acyclic orbit spaces and cohomology eigenvalues of equivariant maps*. Preprint No 8, Oslo (1975).

Matematisk institutt
Universitetet i Oslo,
Blindern, Oslo 3,
Norway.

Received August 30, 1977