

# Rings of $SL_2(\mathbb{C})$ -characters and the Kauffman bracket skein module

Autor(en): **Bullock, Doug**

Objektyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **72 (1997)**

PDF erstellt am: **29.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-54603>

## Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

## Rings of $SL_2(\mathbb{C})$ -characters and the Kauffman bracket skein module

Doug Bullock

**Abstract.** Let  $M$  be a compact orientable 3-manifold. The set of characters of  $SL_2(\mathbb{C})$ -representations of  $\pi_1(M)$  forms a closed affine algebraic set. We show that its coordinate ring is isomorphic to a specialization of the Kauffman bracket skein module, modulo its nilradical. This is accomplished by realizing the module as a combinatorial analog of the ring in which tools of skein theory are exploited to illuminate relations among characters. We conclude with an application, proving that a small manifold's specialized module is necessarily finite dimensional.

**Mathematics Subject Classification (1991).** 57M99.

**Keywords.** Knot, link, skein theory, representation theory, 3-manifold.

### 1. Introduction

The Kauffman bracket skein module is an invariant of 3-manifolds which, until recently, was both difficult to compute and topologically mysterious. A suggestion of its significance came with the discovery that it dominates the ring of  $SL_2(\mathbb{C})$ -characters of the fundamental group [3]. The relationship also provided computational tools [4] and estimates of the module's size [2, 3]. The central result of this paper sharpens the focus considerably, for we show that a specialization, modulo its nilradical, is exactly the ring of characters.

An oriented knot determines a conjugacy class in the fundamental group of a 3-manifold  $M$ . It also determines a function on any set of characters of the group: simply evaluate each character on the conjugacy class. Now suppose that  $\mathbb{C}\mathcal{L}_M$  is the vector space spanned by links in  $M$ , that  $X(M)$  is the set of characters obtained from representations in  $SL_2(\mathbb{C})$ , and that  $\mathbb{C}^{X(M)}$  is the algebra of functions on  $X(M)$ . Let

$$\tilde{\Phi} : \mathbb{C}\mathcal{L}_M \rightarrow \mathbb{C}^{X(M)}$$

be the linear map determined by sending each knot to the negative of its naturally induced function, and each link to the product of the images of its components.

The characters form an affine algebraic set [5] whose coordinate ring  $\mathcal{R}(M)$  lies

in  $\mathbb{C}^{X(M)}$ . The Kauffman bracket skein module of  $M$  acquires a ring structure when its parameter is specialized to negative one. This algebra, denoted  $V(M)$ , is a quotient of  $\mathbb{C}\mathcal{L}_M$ . Our main result and its immediate topological consequence can be summarized as:

**Theorem.** *The map  $\tilde{\Phi}$  descends to  $V(M)$ . Its image is the coordinate ring of  $X(M)$  and its kernel is the nilradical of  $V(M)$ .*

**Theorem.** *If  $M$  contains no non-boundary parallel, incompressible surfaces, then  $V(M)$  is finite dimensional.*

In the next section (which recapitulates parts of [3] and [5]) we cover the necessary background and definitions. The proof that  $\tilde{\Phi}$  descends to a map  $\Phi$  on  $V(M)$  is quite simple, depending primarily on the following observation: the Kauffman bracket skein relation maps to the fundamental  $SL_2(\mathbb{C})$  trace identity,

$$\mathrm{tr}(AB) + \mathrm{tr}(AB^{-1}) = \mathrm{tr}(A)\mathrm{tr}(B).$$

That  $\Phi$  maps onto  $\mathcal{R}(M)$  is also fairly elementary. The bulk of the paper is therefore devoted to characterizing  $\ker \Phi$ .

This begins in Section 3 with an investigation of trace identities on the  $SL_2(\mathbb{C})$ -representations of a free group. There is a map,  $\Psi$ , carrying polynomials in traces to elements of the skein module of a handlebody. Together,  $\Phi$  and  $\Psi$  make skein theory into a graphical calculus for manipulating relations among characters. It turns out that  $\ker \Phi$  contains exactly those trace identities which  $\Psi$  does not send to zero. The central result is that any homogeneous trace identity on  $2 \times 2$  matrices—restricted to  $SL_2(\mathbb{C})$ —lies in the kernel of  $\Psi$ . The proof turns on a classification theorem due independently to Procesi [14] and Razmyslov [16].

Section 3 attains sufficient conditions for a trace identity to vanish in the skein module of a handlebody. Section 4 provides the finishing touch. We rely on a parameterization of the character set given in [8]. Most of the defining polynomials turn out to be specialized Procesi identities, while the remaining few succumb to other tools from Section 3. It follows from a standard result of algebraic geometry that the only trace identities not vanishing in the skein module are nilpotent. It is then a small step to extend the result to arbitrary compact 3-manifolds.

The author would like to thank Professors Charles Frohman, Xiaio-Song Lin, Józef Przytycki and Bruce Westbury for many helpful conversations; the referee for suggesting stylistic improvements; Adam Sikora in particular for his insight into the importance of nilpotents; and the organizers and participants of the Banach Center's 1995 Mini Semester on Knot Theory, where the ideas in this paper first began to coalesce.




Figure 1.

## 2. Definitions and background

Let  $M$  be a compact, orientable 3-manifold. Its Kauffman bracket skein module,  $K(M)$ , is built from the set  $\mathcal{L}_M$  of unoriented, framed links in  $M$ . By a framed link we mean an embedded collection of annuli considered up to isotopy in  $M$ , and we include the empty collection  $\emptyset$ . Three links  $L$ ,  $L_0$  and  $L_\infty$  are said to be *Kauffman skein related* if they can be embedded identically except in a ball where they appear as shown in Figure 1. (Framings are vertical with respect to the page). The notation  $L \amalg \bigcirc$  indicates the union of  $L$  with an unlinked, 0-framed unknot.

Let  $R$  denote the ring of Laurent polynomials  $\mathbb{C}[A^{\pm 1}]$  and  $R\mathcal{L}_M$  the free  $R$ -module with basis  $\mathcal{L}_M$ . If  $L$ ,  $L_0$  and  $L_\infty$  are Kauffman skein related then  $L - AL_0 - A^{-1}L_\infty$  is called a *skein relation*. For any  $L$  in  $\mathcal{L}_M$  the expression  $L \amalg \bigcirc + (A^2 + A^{-2})L$  is called a *framing relation*. Let  $S(M)$  be the smallest submodule of  $R\mathcal{L}_M$  containing all possible skein and framing relations. We define  $K(M)$  to be the quotient  $R\mathcal{L}_M/S(M)$ .

The indeterminate  $A$  is often interpreted as a complex number so that  $K(M)$  becomes a vector space. It seems that the simplest value is  $A = -1$ , and we let  $V(M)$  denote this specialization. Notice that the specialized skein relations imply

$$\begin{array}{c} \diagup \diagdown \\ \diagdown \diagup \end{array} = \begin{array}{c} \diagdown \diagup \\ \diagup \diagdown \end{array}$$

in  $V(M)$ . There is a product on links,  $L_1 L_2 = L_1 \cup L_2$ , which makes  $V(M)$  into a commutative algebra with  $\emptyset$  serving as the identity. It follows from [1, Theorem 1] that  $V(M)$  is generated by a finite set of knots.

By a *representation* we mean a homomorphism of groups

$$\rho : \pi_1(M) \rightarrow SL_2(\mathbb{C}).$$

The *character* of a representation is the composition

$$\chi_\rho = \text{trace} \circ \rho,$$

and  $X(M)$  denotes the set of all characters. For each  $\gamma \in \pi_1(M)$  there is a function  $t_\gamma : X(M) \rightarrow \mathbb{C}$  given by  $\chi_\rho \mapsto \chi_\rho(\gamma)$ .



The following theorem appears to have been discovered independently by Vogt [17] and Fricke [6], first proved by Horowitz [9], and then rediscovered by Culler and Shalen [5].

**Theorem 1.** (Vogt, Fricke, Horowitz, Culler–Shalen). *There exists a finite set of elements  $\{\gamma_1, \dots, \gamma_m\}$  in  $\pi_1(M)$  such that every  $t_\gamma$  is an element of the polynomial ring  $\mathbb{C}[t_{\gamma_1}, \dots, t_{\gamma_m}]$ .*

For Culler and Shalen, Theorem 1 was an initial step in a much deeper result.

**Theorem 2.** (Culler–Shalen). *If every  $t_\gamma$  is an element of  $\mathbb{C}[t_{\gamma_1}, \dots, t_{\gamma_m}]$ , then  $X(M)$  is the zero set of an ideal in  $\mathbb{C}[t_{\gamma_1}, \dots, t_{\gamma_m}]$ .*

Recall that a *closed algebraic set*  $X$  in  $\mathbb{C}^m$  is the common zero set of an ideal of polynomials in  $\mathbb{C}[x_1, \dots, x_m]$ . The elements of  $\mathbb{C}[x_1, \dots, x_m]$  are *polynomial functions* on  $X$ , and the functions  $x_i$  are *coordinates* on  $X$ . The quotient of  $\mathbb{C}[x_1, \dots, x_m]$  by the ideal of polynomials vanishing on  $X$  is called the *coordinate ring* of  $X$ . Different choices of coordinates would clearly lead to different parameterizations of  $X$ , but it follows from [5] that any two parameterizations of  $X(M)$  are equivalent via polynomial maps. Hence their coordinate rings are isomorphic and we may identify them as one object: the *ring of characters* of  $\pi_1(M)$ , which we denote by  $\mathcal{R}(M)$ .

Each knot  $K$  determines a unique  $t_\gamma$  as follows. Let  $\vec{K}$  denote an unspecified orientation on  $K$ . Choose any  $\gamma \in \pi_1(M)$  such that  $\gamma \simeq \vec{K}$  (meaning the loop  $\gamma$  is freely homotopic to an embedding of  $\vec{K}$ ). Since trace is invariant under conjugation it makes sense to define  $\chi_\rho(\vec{K}) = \chi_\rho(\gamma)$ . Since  $\text{tr}(A) = \text{tr}(A^{-1})$  in  $SL_2(\mathbb{C})$  we can also define  $\chi_\rho(K) = \chi_\rho(\gamma)$ . Thus  $K$  determines the map  $\chi_\rho \mapsto \chi_\rho(\gamma)$ . Conversely, any  $t_\gamma$  is determined by some (non-unique)  $K$ . The main theorem of [3] is that this correspondence is well defined at the level of  $V(M)$ .

**Theorem 3.** *The map  $\Phi : V(M) \rightarrow \mathcal{R}(M)$  given by*

$$\Phi(K)(\chi_\rho) = -\chi_\rho(K)$$

*is a well defined surjective map of algebras. If  $V(M)$  is generated by the knots  $K_1, \dots, K_m$  then  $\Phi(K_1), \dots, \Phi(K_m)$  are coordinates on  $X(M)$ .*

*Proof.* Let  $\mathbb{C}^{X(M)}$  denote the algebra of functions from  $X(M)$  to  $\mathbb{C}$ . Define a map

$$\tilde{\Phi} : \mathbb{C}\mathcal{L}_M \rightarrow \mathbb{C}^{X(M)}$$

as follows. If  $K$  is a knot set

$$\tilde{\Phi}(K)(\chi_\rho) = -\chi_\rho(K).$$

If  $L$  is a link with components  $K_1, \dots, K_n$  set

$$\tilde{\Phi}(L) = \prod_{i=1}^n \tilde{\Phi}(K_i).$$

Let  $\tilde{\Phi}(\emptyset) = 1$  and extend linearly.

Consider the image of  $S(M)$  under  $\tilde{\Phi}$ . For a framing relation,  $L \amalg \bigcirc + 2L$ , we have

$$\begin{aligned} \tilde{\Phi}(L \amalg \bigcirc + 2L)(\chi_\rho) &= \tilde{\Phi}(L)\tilde{\Phi}(\bigcirc + 2\emptyset) \\ &= \tilde{\Phi}(L)(-\chi_\rho(\bigcirc) + 2) \\ &= \tilde{\Phi}(L)(-\text{tr}(\text{Id}) + 2) \\ &= 0. \end{aligned}$$

Next, let  $L + L_0 + L_\infty$  be a skein relation in which  $L$  and  $L_0$  are knots. It follows that  $L_\infty$  has two components,  $K_1$  and  $K_2$ . Assume embeddings as in Figure 1 and choose a base point  $*$  in the neighborhood where  $L$ ,  $L_0$  and  $L_\infty$  differ. It is now possible to find loops  $a$  and  $b$  in  $\pi_1(M, *)$  so that a slight perturbation of  $ab$  is  $\vec{L}$ . With favorable orientations on the other knots we have  $ab^{-1} \simeq \vec{L}_0$ ,  $a \simeq \vec{K}_1$ , and  $b \simeq \vec{K}_2$ . Given any  $\chi_\rho$ , set  $A = \rho(a)$  and  $B = \rho(b)$  so that

$$\begin{aligned} \tilde{\Phi}(L + L_0 + L_\infty)(\chi_\rho) &= -\chi_\rho(L) - \chi_\rho(L_0) + \chi_\rho(K_1)\chi_\rho(K_2) \\ &= -\text{tr}(AB) - \text{tr}(AB^{-1}) + \text{tr}(A)\text{tr}(B) \\ &= 0. \end{aligned}$$

Finally, note that every skein relation can be written as  $L' \cup L + L' \cup L_0 + L' \cup L_\infty$  where  $L$  and  $L_0$  are knots. Hence  $\tilde{\Phi}$  descends to a well defined map of algebras,

$$\Phi : V(M) \rightarrow \mathbb{C}^{X(M)},$$

which is determined by its values on knots.

Let  $K_1, \dots, K_m$  be generators of  $V(M)$ . Every element of  $V(M)$  can be written as a polynomial in these knots, so the image of  $\Phi$  lies in  $\mathbb{C}[\Phi(K_1), \dots, \Phi(K_m)]$ . Since each  $t_\gamma$  is equal to  $-\Phi(K)$  for some knot  $K$ , Theorems 1 and 2 imply that the functions  $\Phi(K_i)$  are coordinates on  $X(M)$ . It follows that  $\Phi$  maps onto  $\mathcal{R}(M)$ .  $\square$




Figure 2.

### 3. Trace identities

In the previous section we obtained a surjection  $\Phi : V(M) \rightarrow \mathcal{R}(M)$  based on a natural correspondence between knots and functions on  $X(M)$ . Under this correspondence elements of  $S(M)$  were sent to polynomials that vanish on  $X(M)$ , making  $\Phi$  well defined. Our ultimate goal is to show that  $\ker \Phi$  is the set of nilpotent elements in  $V(M)$ . To this end we reverse the correspondence, mapping polynomials on  $X(M)$  to elements of  $V(M)$ . For now, we will treat only the case where  $M$  is a handlebody. In this setting the kernel of  $\Phi$  consists of polynomials that vanish on  $X(M)$  but not in  $V(M)$ .

For the time being we will be concerned only with free groups, so throughout this section  $H$  will denote the manifold  $P \times I$  where  $P$  is the planar surface in Figure 2. We also fix a base point  $*$  in  $P$  and a set of generators  $\{a_1, \dots, a_n\}$  for  $\pi_1(H, *)$ . Each loop  $a_i$  travels once across the  $i$ -th handle in the direction shown in Figure 2. Let  $\mathcal{W}$  denote  $\pi_1(H, *)$  modulo the equivalence generated by

$$w \sim w' \iff w' = w^{-1}, \text{ or } w' = gw g^{-1} \text{ for some } g \in \pi_1(H, *).$$

Consider the ring of polynomials  $\mathbb{C}[\mathcal{W}]$ .

**Example 1.**

$$p = (a_1)(a_2)(a_3) - (a_1a_2)(a_3) - (a_1a_3)(a_2) - (a_2a_3)(a_1) + (a_1a_2a_3) + (a_1a_3a_2)$$

**Example 2.**

$$\begin{aligned} q = & (a_1)^2 + (a_2)^2 + (a_3)^2 + (a_1a_2)^2 + (a_1a_3)^2 + (a_2a_3)^2 \\ & + (a_1a_2a_3)^2 + (a_1a_2)(a_1a_3)(a_2a_3) + (a_1a_2a_3)(a_1)(a_2)(a_3) \\ & - (a_1a_2a_3)(a_1)(a_2a_3) - (a_1a_2a_3)(a_2)(a_1a_3) - (a_1a_2a_3)(a_3)(a_1a_2) \\ & - (a_1)(a_2)(a_1a_2) - (a_1)(a_3)(a_1a_3) - (a_2)(a_3)(a_2a_3) - 4 \end{aligned}$$

The parentheses are necessary to distinguish multiplication in  $\pi_1(H)$  from multiplication in  $\mathbb{C}[\mathcal{W}]$ . Note that there is some ambiguity in the notation for an individual element of  $\mathbb{C}[\mathcal{W}]$ . For instance  $(w^2) + (1) - (w)^2$  is the same as  $(ww) + (ww^{-1}) - (w)(w^{-1})$ . Occasionally it will be convenient to write a polynomial using non-reduced words.

A representation of  $\pi_1(H, *)$  in  $SL_2(\mathbb{C})$  is any assignment of matrices to each  $a_i$ . Letting parentheses denote the operation of trace, each element of  $\mathbb{C}[\mathcal{W}]$  becomes a function from the representation space to  $\mathbb{C}$ . The elements of  $\mathbb{C}[\mathcal{W}]$  that vanish as functions on the set of representations are called  $SL_2(\mathbb{C})$ -trace identities. They form an ideal  $\mathcal{I} \subset \mathbb{C}[\mathcal{W}]$ .

Each  $w \in \mathcal{W}$  corresponds, up to free homotopy, to a unique unoriented curve in  $H$ . We will use  $K_w$  to denote any knot in this homotopy class. Since crossings are irrelevant,  $K_w$  represents a unique element of  $V(H)$ . The assignment  $w \mapsto -K_w$  defines a surjection of algebras,

$$\Psi : \mathbb{C}[\mathcal{W}] \rightarrow V(H).$$

The map  $\Psi$  turns an element of  $\mathbb{C}[\mathcal{W}]$  into a linear combination of links in  $V(H)$ . The basic tool for calculating in the skein module is a resolving tree. Let  $T$  be a finite, connected, contractible graph in which no vertex has valence greater than three. Suppose each vertex is labeled  $cL$  for some  $c \in \mathbb{C}$  and some  $L \in \mathcal{L}_H$ . Assume further that there is a distinguished vertex  $c_0L_0$  called the *root*. Define the potential of a vertex to be the number of edges in a path to the root. A (necessarily univalent) vertex that is not adjacent to one of higher potential is called a *leaf*. We say  $T$  is a *resolving tree* for  $c_0L_0$  if each vertex  $cL$  satisfies exactly one of the following:

- (1)  $cL$  is a leaf.
- (2)  $cL$  is adjacent to exactly one higher potential vertex,  $-2cL$ .
- (3)  $cL$  is adjacent to exactly two higher potential vertices,  $c'L'$  and  $c''L''$ , in which case  $cL - c'L' - c''L''$  is a skein relation.

Figure 3, in which the dots represent a thrice punctured plane, is a resolving tree for any knot that projects to the leftmost diagram. It is also an example of the most common way to produce a resolving tree. Beginning with a projection of the root, the tree grows by smoothing one crossing at a time. Once all crossings have been eliminated, trivial circles are removed via framing relations. The sum of the leaves is the *standard resolution* of the root—an element of  $\mathbb{C}\mathcal{L}_H$  which is equal to the root in  $V(H)$ . Although the procedure given here does not result in a unique tree, the following theorem [15] implies uniqueness of the standard resolution in  $\mathbb{C}\mathcal{L}_H$ .

**Theorem 4.** (Przytycki). *The links in  $H$  represented by diagrams in  $P$  with no crossings and no trivial circles are a basis for  $V(H)$ .*

A *resolving forest* for an element of  $\mathbb{C}\mathcal{L}_H$  is simply a collection of trees, one for each term in the linear combination. As with individual links, there is a standard resolution of each element of  $\mathbb{C}\mathcal{L}_H$ . Summing the potential function over all vertices assigns a useful complexity to a forest, the *total potential*.

The remainder of this section is devoted to the establishment of conditions under which  $\Psi$  maps a polynomial to zero.

**Lemma 1.** *In Examples 1 and 2 we have  $\Psi(p) = \Psi(q) = 0$ .*

*Proof.* For Example 1, suppose that  $P$  has been deformed to look like Figure 4 and that its fundamental group is generated by the indicated loops. The map  $\Psi$






Figure 4.




Figure 5.

applied to  $-(a_1a_2a_3)$  is the root of Figure 3. The leaves of the tree, reading down the page, are

$$\begin{aligned} & -2\Psi((a_1a_3a_2)) + \Psi((a_1a_3a_2)) - \Psi((a_1a_2)(a_3)) + \Psi((a_1a_3a_2)) \\ & + \Psi((a_1)(a_2)(a_3)) - \Psi((a_2a_3)(a_1)) - \Psi((a_1a_3)(a_2)) + \Psi((a_1a_3a_2)). \end{aligned}$$

Since the sum of the leaves of a resolving tree is equal to its root,  $\Psi(p) = 0$ .

To see that  $\Psi(q) = 0$ , resolve the diagram of  $-\Psi((a_1a_2)(a_1a_3)(a_2a_3))$  shown in Figure 5.  $\square$

If  $p$  is a trace identity then a natural way to produce a new trace identity,  $q$ , is to substitute new words for each  $a_i$  in  $p$ . If  $\Psi(p) = 0$  then one would hope  $\Psi(q) = 0$  as well. Although this is true, the proof requires some effort.

**Lemma 2.** *Let  $p \in \mathbb{C}[\mathcal{W}]$ . If there exist words  $w_1$  and  $w_2$  such that  $(w_1w_2) + (w_1w_2^{-1}) - (w_1)(w_2)$  divides  $p$ , then  $\Psi(p) = 0$ . Also, if  $p$  is divisible by  $(1) - 2$  then  $\Psi(p) = 0$ .*

*Proof.* In the first case consider the loop  $w_1w_2$ , but perturbed slightly so as to become an embedding. The resulting knot is, by definition, some  $K_{w_1w_2}$ . Similarly, perturb  $w_1w_2^{-1}$ ,  $w_1$ ,  $w_2$  to obtain  $K_{w_1w_2^{-1}}$ ,  $K_{w_1}$ ,  $K_{w_2}$ . The perturbations may be chosen to create embeddings of  $K_{w_1w_2}$ ,  $K_{w_1w_2^{-1}}$  and  $K_{w_1}K_{w_2}$  that coincide outside of a small neighborhood of the base point. Within that neighborhood they can be made to look like Figure 1, so they form a Kauffman skein triple. We now have

$$\begin{aligned} 0 &= -K_{w_1w_2} - K_{w_1w_2^{-1}} - K_{w_1}K_{w_2} \\ &= \Psi((w_1w_2) + (w_1w_2^{-1}) - (w_1)(w_2)), \end{aligned}$$

which implies  $\Psi(p) = 0$ .

In the second case  $\Psi(p)$  contains a factor of  $\bigcirc + 2 \emptyset$ , which also implies  $\Psi(p) = 0$ .  $\square$

**Proposition 1.** *Let  $p \in \mathbb{C}[W]$ . Choose words  $w_1, \dots, w_n$ , and form a new polynomial  $q$  by substituting  $w_i$  for  $a_i$  in  $p$ . If  $\Psi(p) = 0$  then  $\Psi(q) = 0$ .*

*Proof.* The proof is by induction on a complexity,  $\kappa(p)$ , which we define as follows. For each  $w \in W$  choose a diagram in  $P$  representing  $K_w$ . Express  $\Psi(p)$  as an element of  $\mathbb{C}\mathcal{L}_H$  using these diagrams, and then choose a forest for its standard resolution. Define  $\kappa(p)$  to be the minimum total potential over all choices of diagrams and forests.

Assume first that  $\kappa(p) = 0$ , implying diagrams in which  $\Psi(p)$  is expressed as its own standard resolution. If  $\Psi(p) = 0$ , we can invoke Theorem 4 to conclude that this particular expression of  $\Psi(p)$  is formally zero in  $\mathbb{C}\mathcal{L}_H$ . It is not possible for a diagram to represent more than one  $w$ , so  $p$  (and hence  $q$ ) must be identically zero.

Now assume that  $\kappa(p) > 0$ . Choose diagrams and a forest realizing  $\kappa(p)$ ; also select a root  $cL$  which is not a leaf. There are three cases depending on the first resolution of  $cL$ .

*Case 1: The resolution removes a self crossing of some component.* Letting  $K$  denote that component, we construct loops in  $\pi_1(H, *)$ . Begin by choosing a point  $x$  near the crossing in question. Let  $\alpha_0$  be an arc running from  $*$  to  $x$ ; let  $\alpha_1$  be an arc running parallel to  $\vec{K}$  until it returns to  $x$ ; and let  $\alpha_2$  be an arc parallel to the remaining portion of  $\vec{K}$ . Set  $\gamma_1 = \alpha_0 \alpha_1 \alpha_0^{-1}$  and  $\gamma_2 = \alpha_0 \alpha_2 \alpha_0^{-1}$ . We now have  $K = K_{\gamma_1 \gamma_2}$ . Furthermore, the resolution changes  $K$  into  $K_{\gamma_1 \gamma_2^{-1}}$  and  $K_{\gamma_1} K_{\gamma_2}$ .

The term of  $p$  represented by  $cL$  must contain the indeterminate  $(\gamma_1 \gamma_2)$ . Replace that appearance of  $(\gamma_1 \gamma_2)$  with  $(\gamma_1)(\gamma_2) - (\gamma_1 \gamma_2^{-1})$ , creating a new polynomial  $p'$ . Since  $p - p'$  is divisible by  $r = (\gamma_1 \gamma_2) + (\gamma_1 \gamma_2^{-1}) - (\gamma_1)(\gamma_2)$ , Lemma 2 implies  $\Psi(p') = 0$ . Let  $q'$  and  $r'$  be the results of substituting  $w_i$  for  $a_i$  in  $p'$  and  $r$  (respectively). Removing the root  $cL$  from the forest for  $\Psi(p)$  produces a forest for  $\Psi(p')$  with lower total potential. Hence  $\kappa(p') < \kappa(p)$  and, by induction,  $\Psi(q') = 0$ . Furthermore,  $r'$  has the form  $(\gamma'_1 \gamma'_2) + (\gamma'_1 \gamma'_2^{-1}) - (\gamma'_1)(\gamma'_2)$ . Since  $r'$  divides  $q - q'$  we have  $\Psi(q) = 0$ .

*Case 2: The resolution removes a crossing between two components.* In this case the components involved in the crossing correspond to loops  $\gamma_1$  and  $\gamma_2$ , for which the resolution produces  $K_{\gamma_1 \gamma_2}$  and  $K_{\gamma_1 \gamma_2^{-1}}$ . As in Case 1 we create  $p'$  by replacing  $(\gamma_1)(\gamma_2)$  in  $p$  with  $(\gamma_1 \gamma_2) + (\gamma_1 \gamma_2^{-1})$ . The proof then proceeds by induction as before.

*Case 3: The resolution removes a trivial circle.* The trivial circle corresponds to an appearance of  $(1)$  in  $p$ . Form  $p'$  by replacing that  $(1)$  with the scalar 2. Then create  $q$  and  $q'$  as above, noting that  $(1) - 2$  divides both  $p - p'$  and  $q - q'$ . As

above,  $\kappa(p') < \kappa(p)$ , and it follows that  $\Psi(q) = 0$ .  $\square$

We would now like to consider a more general sort of trace identity. Let  $S_n$  denote the group of permutations of the set  $\{a_1, \dots, a_n\}$ . Let  $S_m$  denote a group of permutations of some subset  $\{a_{i_1}, \dots, a_{i_m}\}$ . Consider the group algebra  $\mathbb{C}S_m$ . By writing the elements of  $S_m$  in cycle notation, including trivial cycles, we obtain expressions in  $\mathbb{C}[\mathcal{W}]$ . (Example 1, for instance). In fact, since no inverses appear in these expressions, they can be regarded as functions on the set of  $m$ -tuples of  $2 \times 2$  matrices. If an element of  $\mathbb{C}S_m$ , regarded as such a function, vanishes for every assignment of  $2 \times 2$  matrices we call it a *Procesi identity* on  $S_m$ . Note that a Procesi identity is clearly an  $SL_2(\mathbb{C})$  trace identity, but that the converse is just as clearly false.

Using the group algebra to encode Procesi identities is useful for the theorem we are about to prove, but there is a drawback as well. Multiplication in  $\mathbb{C}S_m$  is not the same as multiplication in  $\mathbb{C}[\mathcal{W}]$ . If  $p$  and  $q$  are elements of  $\mathbb{C}S_m$  we denote their product in the group algebra as  $p \cdot q$ , always assuming that  $p$ ,  $q$  and  $p \cdot q$  are written in cycle notation. Note that  $pq$  need not lie in  $\mathbb{C}S_m$ , and that  $p \cdot q$  may involve elements of  $\mathcal{W}$  which do not appear in either  $p$  or  $q$ . Fortunately, the skein module keeps track of how multiplication in  $S_m$  rearranges the elements of  $\mathcal{W}$ .

**Proposition 2.** *Let  $p \in \mathbb{C}S_m$ . If  $\Psi(p) = 0$  then  $\Psi(\tau \cdot p) = 0$  for every  $\tau \in S_n$ .*

*Proof.* As an initial simplification assume that  $\tau = (a_i a_j)$  with  $i < j$ , and that  $S_m$  permutes the set  $\{a_1, \dots, a_m\}$ . There are three cases, depending on the intersection of  $\{a_1, \dots, a_m\}$  and  $\{a_i, a_j\}$ .

*Case 1:*  $m < i$ . As an element of  $\mathbb{C}[\mathcal{W}]$ ,  $\tau \cdot p$  factors into  $(a_i a_j)p$ . Hence  $\Psi(p) = 0$  implies  $\Psi(\tau \cdot p) = 0$ .

*Case 2:*  $i \leq m < j$ . Each term of  $p$  contains a cycle in which  $a_i$  appears. Assume that it is written  $(a_i \alpha)$  and write  $\tau \cdot (a_i \alpha)$  as  $(a_j a_i \alpha)$ . Fix a diagram for each term of  $\Psi(p)$  with the property that it traverses handles 1 through  $m$  exactly once and misses the others. In a resolving forest for the standard resolution of  $\Psi(p)$  the skein relations take place in neighborhoods away from the handles, and no trivial circle runs once over a handle. Therefore every diagram in the forest meets the handles in precisely the same set of arcs, and we can apply the operation shown in Figure 6 to the entire forest. Note that this changes the diagram for  $\Psi((a_i \alpha))$  into a diagram for  $\Psi((a_j a_i \alpha))$ , producing a resolution of  $\Psi(\tau \cdot p)$ . By Theorem 4, the resolution of  $\Psi(p)$  is formally zero in  $\mathbb{C}\mathcal{L}_H$ . Since the resolution of  $\Psi(\tau \cdot p)$  is obtained by applying Figure 6 to each term, it must also be zero.

*Case 3:*  $j \leq m$ . Each term of  $p$  contains either a cycle  $(a_i \alpha a_j \beta)$  or a product of cycles  $(a_i \alpha)(a_j \beta)$ . The action of  $\tau$  interchanges the two possibilities. Notice that the operation in Figure 7 interchanges the diagrams for  $\Psi((a_i \alpha)(a_j \beta))$  and  $\Psi((a_i \alpha a_j \beta))$ . The proof then follows the resolving argument of Case 2.

Subject to our initial simplification, we now have  $\Psi(\tau \cdot p) = 0$ . Retaining the






Figure 6.  
Band sum of  $\Psi(a_i\alpha)$  and  $\Psi(a_j)$ .




Figure 7.  
Band relating  $\Psi((a_i\alpha)(a_j\beta))$  and  $\Psi(a_i\alpha a_j\beta)$ .

assumption that  $\tau = (a_i a_j)$ , we next allow  $S_m$  to permute any set  $\{a_{i_1}, \dots, a_{i_m}\}$ . A substitution converts this set into  $\{a_1, \dots, a_m\}$ , but preserves that fact that  $\tau$  is a transposition. Hence, by Proposition 1, we again have  $\Psi(\tau \cdot p) = 0$ . Finally, since any element of  $S_n$  is a product of transpositions,  $\Psi(\tau \cdot p) = 0$  for all  $\tau \in S_n$ .  $\square$

Our interest in Procesi identities stems from a classification theorem due independently to Procesi [14] and Razmyslov [16]. Leron [12] is an excellent reference for the proof. For the sake of completeness we include some definitions taken from [7, Chapter 4]. A *Young diagram* for  $S_m$  is a collection of  $m$  boxes arranged in left justified rows of decreasing length. A *Young tableau* is an assignment of  $a_{i_1}, \dots, a_{i_m}$  to the boxes. Figure 8 is an example of a Young diagram for  $S_{10}$  and a tableau using  $\{a_1, \dots, a_{10}\}$ . Given a tableau  $Y$  define  $P_Y$  to be the subgroup of  $S_m$  stabilizing the rows. For the tableau in Figure 8

$$P_Y \cong S_3 \times S_3 \times S_2.$$

Similarly, define  $Q_Y$  to be the column stabilizer. The Young symmetrizer corresponding to  $Y$  is the element

$$\left( \sum_{\sigma \in P_Y} \sigma \right) \cdot \left( \sum_{\tau \in Q_Y} \text{sgn}(\tau) \tau \right) \in \mathbb{C}S_m.$$




Figure 8.

**Theorem 5.** (Procesi, Razmyslov). *Procesi identities on a fixed  $S_m$  constitute an ideal in  $\mathbb{C}S_m$ . The ideal is generated by Young symmetrizers corresponding to diagrams with at least three rows.*

**Lemma 3.** *Let  $Y$  be a Young tableau on  $\{a_1, \dots, a_m\}$ , and assume that  $a_m$  occupies the last box of a row and column as shown schematically in Figure 9. Let  $Y'$  be the tableau obtained from  $Y$  by removing the box containing  $a_m$ . Using the notation  $r_{s+1} = c_{t+1} = a_m$ , we can express  $P_Y$  and  $Q_Y$  as the following disjoint unions:*

$$(1) \quad P_Y = \bigcup_{i=1}^{s+1} (r_i a_m) \cdot P_{Y'}, \text{ and}$$

$$(2) \quad Q_Y = \bigcup_{i=1}^{t+1} (c_i a_m) \cdot Q_{Y'}.$$

*Proof.* Let  $\lambda_1, \dots, \lambda_x$  be the lengths of the rows of  $Y'$ . A row stabilizer is a product of symmetric groups, so

$$|P_{Y'}| = (\lambda_1!) \cdots (\lambda_x!), \quad \text{and}$$

$$|P_Y| = (s+1)(\lambda_1!) \cdots (\lambda_x!).$$

Each coset  $(r_i a_m) \cdot P_{Y'}$  stabilizes the rows of  $Y$  and, since each contains the element  $(r_i a_m)$ , they are disjoint. Counting elements finishes the proof for  $P_Y$ . The proof for  $Q_Y$  is similar.  $\square$

**Theorem 6.** *If  $p$  is a Procesi identity then  $\Psi(p) = 0$ .*

*Proof.* Implicit in the statement is the fact that  $p$  is a Procesi identity on some  $S_m$ , so we proceed by induction on  $m$ . By Theorem 5 and Proposition 2 we may assume




Figure 9.

that  $p$  is a Young symmetrizer corresponding to a digram with at least three rows. If  $m = 3$  there is only one such diagram and  $p$  is the result of substituting  $a_{i_1}$ ,  $a_{i_2}$  and  $a_{i_3}$  into Example 1. By Lemma 1 and Proposition 1,  $\Psi(p) = 0$ .

Now assume  $m > 3$ . Choose a diagram with at least three rows and a tableau  $Y$  satisfying the hypotheses of Lemma 3. A symmetrizer corresponding to any other tableau with the same diagram is obtained from this one by a substitution. Therefore, by Proposition 1, it suffices to consider only  $Y$ . With notation as in Lemma 3, let  $p'$  be the symmetrizer corresponding to the tableau  $Y'$ . We now have

$$\begin{aligned}
 p &= \left( \sum_{\sigma \in P_Y} \sigma \right) \cdot \left( \sum_{\tau \in Q_Y} \text{sgn}(\tau) \tau \right) \\
 &= \left[ \sum_{i=1}^{s+1} \left( \sum_{\sigma \in P_{Y'}} (r_i a_m) \cdot \sigma \right) \right] \cdot \left[ \sum_{j=1}^{t+1} \left( \sum_{\tau \in Q_{Y'}} \text{sgn}((c_j a_m) \cdot \tau) (c_j a_m) \cdot \tau \right) \right] \\
 &= \sum_{i,j} \text{sgn}(c_j a_m) (r_i a_m) \cdot (c_j a_m) \cdot \left( \sum_{\sigma \in P_{Y'}} \sigma \right) \cdot \left( \sum_{\tau \in Q_{Y'}} \text{sgn}(\tau) \tau \right) \\
 &= \sum_{i,j} \text{sgn}(c_j a_m) (r_i a_m) \cdot (c_j a_m) \cdot p'.
 \end{aligned}$$

By induction and Proposition 2,  $\Psi(p) = 0$ . □

#### 4. The coordinate ring

Consider the following diagram of algebra maps:

$$\begin{array}{ccc} V(H) & \xleftarrow{\Psi} & \mathbb{C}[\mathcal{W}] \\ \downarrow \Phi & & \downarrow \\ \mathcal{R}(H) & \xleftarrow{?} & \mathbb{C}[\mathcal{W}]/\mathcal{I} \end{array}$$

We know that  $\Psi$  and  $\Phi$  are surjections. If we can establish an identification along the bottom row, then the kernel of  $\Phi$  will be exactly those elements of  $\mathcal{I}$  that do not lie in  $\ker \Psi$ . In Section 3 we developed conditions under which a trace identity does map to zero; what we lack is a list of generators for  $\mathcal{I}$ . In order to complete the picture we will have to choose coordinates on  $X(H)$ . In other words, we need a finitely generated subalgebra of  $\mathbb{C}[\mathcal{W}]$  and an explicitly described ideal so that the quotient is  $\mathcal{R}(H)$ . The subalgebra and the ideal (up to nilpotents) are taken from [8]. Our tasks are to show that the restricted  $\Psi$  is still onto, to prove that it carries the ideal to zero, and to clarify the bottom row of the diagram. This will establish the main theorem for handlebodies, after which it extends easily to compact, oriented 3-manifolds.

Not only did Vogt [17], Fricke [6], and Culler and Shalen [5] apparently discover Theorem 1 independently, they all arrived at the same set of generators. Let  $\gamma = a_{i_1} \cdots a_{i_m}$  be an element of  $G$  in which each  $a_{i_j}$  is distinct. Following [8] we adopt the shorthand notation  $t_{i_1 \dots i_m}$  for the map  $t_\gamma$ . The generating set in all versions of Theorem 1 is  $\mathcal{T} = \{t_{i_1 \dots i_m} \mid i_1 < i_2 < \cdots < i_m\}$ .

Note that  $\mathbb{C}[\mathcal{T}]$  becomes a subring of  $\mathbb{C}[\mathcal{W}]$  by replacing  $t_{i_1 \dots i_m}$  with  $(a_{i_1} \cdots a_{i_m})$ , so  $\Psi$  is well defined on  $\mathbb{C}[\mathcal{T}]$ . Theorem 1 says that for every  $p \in \mathbb{C}[\mathcal{W}]$  there exists  $q \in \mathbb{C}[\mathcal{T}]$  such that  $p$  and  $q$  represent the same element of  $\mathcal{R}(H)$ . To maintain surjectivity of  $\Psi$  we need a stronger result, for which we turn to the combinatorial construction of  $\mathcal{T}$  in [1].

**Theorem 7.** *Let  $\mathcal{K}$  be a set of knots containing exactly one  $K_\gamma$  for each  $t_\gamma \in \mathcal{T}$ . Any link  $L \in \mathcal{L}_H$  has a resolving tree whose leaves are monomials in  $\mathbb{C}[\mathcal{K}]$ .*

**Corollary 1.** *For every  $p \in \mathbb{C}[\mathcal{W}]$  there exists  $q \in \mathbb{C}[\mathcal{T}]$  such that  $\Psi(p) = \Psi(q)$ .*

The main result of [8] is the construction of an ideal,  $\mathcal{J}_H$ , which defines  $X(H)$  in the coordinates  $\mathcal{T}_0 = \{t_{i_1 \dots i_m} \in \mathcal{T} \mid m \leq 3\}$ . The radical of this ideal,  $\sqrt{\mathcal{J}_H}$ , is the ideal of trace identities in  $\mathbb{C}[\mathcal{T}_0]$ . The authors of [8] prove that the trace identities in  $\mathbb{C}[\mathcal{T}_0]$  generate those in  $\mathbb{C}[\mathcal{T}]$ , but once again we need a slightly stronger statement.

**Lemma 4.** (Compare [8, Lemma 4.1.1]). *Choose distinct indices  $i, j, k, m_1, \dots, m_l$*

and let  $\alpha = m_1 \cdots m_l$ . If

$$\begin{aligned} q = & -2t_{ijk\alpha} + t_{ik}t_jt_\alpha - t_it_jt_{k\alpha} - t_jt_kt_{\alpha i} - t_{ik}t_{j\alpha} \\ & + t_{ij}t_{k\alpha} + t_{jk}t_{\alpha i} - t_{ikj}t_\alpha + t_it_jt_{k\alpha} + t_jt_{k\alpha i} + t_kt_{\alpha ij} \end{aligned}$$

then  $\Psi(q) = 0$ .

*Proof.* Let  $p_{xyz}$  denote the Procesi identity of Example 1 with the substitutions  $a_1 = a_x$ ,  $a_2 = a_y$  and  $a_3 = a_z$ . Consider the polynomial

$$p' = (a_1a_3) \cdot p_{234} - (a_3a_4) \cdot p_{124} - (a_1a_4) \cdot p_{123}$$

as an element of  $\mathbb{C}[\mathcal{W}]$ . By Theorem 5,  $p'$  is a Procesi identity, implying  $\Psi(p') = 0$ . Substituting  $a_1 = a_i$ ,  $a_2 = a_j$ ,  $a_3 = a_k$  and  $a_4 = a_{m_1} \cdots a_{m_l}$  in  $p'$  gives  $q$ , so Proposition 1 implies  $\Psi(q) = 0$ .  $\square$

**Proposition 3.** *For every  $q \in \mathbb{C}[T]$  there exists  $q_0 \in \mathbb{C}[T_0]$  such that  $\Psi(q) = \Psi(q_0)$ .*

*Proof.* Let  $q \in \mathbb{C}[T]$ . Define  $l$  to be the maximum length of a subscript appearing in  $q$  and let  $m$  be the number of maximum length subscripts. We say that the complexity of  $q$  is the ordered pair  $(l, m)$ . The proof is by induction on complexity, ordered lexicographically. If  $l \leq 3$  then  $q$  can be converted to  $q_0 \in \mathbb{C}[T_0]$  by repeated application of the identity in Example 1 (perhaps with a substitution of indices). The difference between any pair of successive stages is divisible by a Procesi identity, so Theorem 6 implies  $\Psi(q) = \Psi(q_0)$ .

If  $l > 3$  then  $q$  contains some  $t_{ijk\alpha}$  in which  $ijk\alpha$  is a maximum length subscript. The identity of Lemma 4 allows us to replace  $t_{ijk\alpha}$  with an expression involving only shorter subscripts. The result is a new polynomial,  $q'$ , with lower complexity and  $\Psi(q) = \Psi(q')$ .  $\square$

We now state the main result of [8]. Let

$$\begin{aligned} M_{ii} &= t_i^2 - 4, & \text{and} \\ M_{ij} &= M_{ji} = 2t_{ij} - t_it_j, & \text{if } i < j. \end{aligned}$$

**Theorem 8.** (González-Acuña–Montesinos).  *$X(M)$  is the zero set of the ideal  $\mathcal{J}_H$  in  $\mathbb{C}[T_0]$  generated by the following polynomials.*

$$\begin{aligned} q_1 = & t_i^2 + t_j^2 + t_k^2 + t_{ij}^2 + t_{ik}^2 + t_{jk}^2 + t_{ijk}^2 + t_{ij}t_{ik}t_{jk} + t_{ijk}t_it_jt_k \\ & - t_{ijk}t_it_{jk} - t_{ijk}t_jt_{ik} - t_{ijk}t_kt_{ij} - t_it_jt_{ij} - t_it_kt_{ik} - t_jt_kt_{jk} - 4, \end{aligned}$$

in which  $i, j$  and  $k$  are distinct.

$$\begin{aligned}
q_2 &= \begin{vmatrix} M_{11} & M_{12} & M_{1i} & M_{1j} \\ M_{21} & M_{22} & M_{2i} & M_{2j} \\ M_{i1} & M_{i2} & M_{ii} & M_{ij} \\ M_{j1} & M_{j2} & M_{ji} & M_{jj} \end{vmatrix}, \quad \text{with } 2 < i < j \leq n. \\
q_3 &= \begin{vmatrix} M_{11} & M_{12} & M_{13} & M_{1i} \\ M_{21} & M_{22} & M_{23} & M_{2i} \\ M_{31} & M_{32} & M_{33} & M_{3i} \\ M_{j1} & M_{j2} & M_{j3} & M_{ji} \end{vmatrix}, \quad \text{with } 3 < i < j \leq n. \\
q_4 &= (t_{123} - t_{132})(2t_{ijk} + t_i t_j t_k - t_i t_{jk} - t_j t_{ik} - t_k t_{ij}) - \begin{vmatrix} t_1 & t_{1i} & t_{1j} & t_{1k} \\ t_2 & t_{2i} & t_{2j} & t_{2k} \\ t_3 & t_{3i} & t_{3j} & t_{3k} \\ 2 & t_i & t_j & t_k \end{vmatrix}, \\
&\quad \text{in which } 1 \leq i < j < k \leq n \text{ and } t_{mm} \text{ denotes } t_m^2 - 2.
\end{aligned}$$

The next step is to show that all of these polynomials lie in  $\ker \Psi$ . The proofs involved in this are closely modeled on those in [8]. Our contribution is the observation that  $q_2, q_3$  and  $q_4$  lift to Procesi identities, and that  $q_1$  follows directly from a resolving tree. We will introduce further notation from [8] as it becomes necessary.

**Lemma 5.**  $\Psi(q_1) = 0$ .

*Proof.* Example 2, Lemma 1 and Proposition 1. □

Let  $A_1, \dots, A_4$  be  $2 \times 2$  matrices with

$$A_i = \begin{pmatrix} \alpha_i & \beta_i \\ \gamma_i & \delta_i \end{pmatrix}.$$

Define

$$M(A_i) = \begin{pmatrix} \alpha_1 & \beta_1 & \gamma_1 & \delta_1 \\ \alpha_2 & \beta_2 & \gamma_2 & \delta_2 \\ \alpha_3 & \beta_3 & \gamma_3 & \delta_3 \\ \alpha_4 & \beta_4 & \gamma_4 & \delta_4 \end{pmatrix}, \quad J = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \text{and } J^* = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}.$$

In general,  $(c_{ij})$  will denote a  $4 \times 4$  matrix and  $|c_{ij}|$  its determinant. We use  $A^t$  to denote the transpose of  $A$ .

**Lemma 6.** (Compare [8, Lemma 4.6]). *The polynomial  $p = |2(x_i y_j) - (x_i)(y_j)|$  is a Procesi identity on the symmetric group permuting  $\{x_1, x_2, x_3, x_4, y_1, y_2, y_3, y_4\}$ .*

*Proof.* Clearly  $p$  is an element of the group algebra over the permutations of  $\{x_1, x_2, x_3, x_4, y_1, y_2, y_3, y_4\}$ . To see that  $p$  is an identity, assign matrices  $A_i$  and

$B_j$  to each  $x_i$  and  $y_j$ . Direct calculation shows that

$$(\text{tr}(A_i B_j)) = M(A_i) J M(B_j)^t,$$

and

$$(\text{tr}(A_i B_j) - \text{tr}(A_i) \text{tr}(B_j)) = -M(A_i) J J^* M(B_j)^t.$$

Since  $|I - J^*| = 0$  we have

$$|2 \text{tr}(A_i B_j) - \text{tr}(A_i) \text{tr}(B_j)| = |M(A_i) J (I - J^*) M(B_j)^t| = 0.$$

□

**Lemma 7.** (Compare [8, Corollary 4.12]). *The polynomial*

$$\begin{aligned} p = & [(x_1 x_2 x_3) - (x_1 x_3 x_2)] \\ & \times [2(y_1 y_2 y_3) + (y_1)(y_2)(y_3) - (y_1)(y_2 y_3) - (y_2)(y_1 y_3) - (y_3)(y_1 y_2)] \\ & - \begin{vmatrix} (x_1) & (x_1 y_1) & (x_1 y_2) & (x_1 y_3) \\ (x_2) & (x_2 y_1) & (x_2 y_2) & (x_2 y_3) \\ (x_3) & (x_3 y_1) & (x_3 y_2) & (x_3 y_3) \\ 2 & (y_1) & (y_2) & (y_3) \end{vmatrix} \end{aligned}$$

is a Procesi identity on  $\{x_1, x_2, x_3, y_1, y_2, y_3\}$ .

*Proof.* (Compare [8, Lemma 4.10 and Proposition 4.11]). For any  $2 \times 2$  matrices  $A_1, \dots, A_4, B_1, \dots, B_4$  we have

$$\begin{aligned} |\text{tr}(A_i B_j)| &= |M(A_i) J M(B_j)| \\ &= |J| |M(A_i)| |M(B_j)| \\ &= -|M(A_i)| |M(B_j)|. \end{aligned}$$

If  $A_4 = I$  then, by direct calculation,

$$|M(A_i)| = \text{tr}(A_1 A_2 A_3) - \text{tr}(A_1 A_3 A_2).$$

If  $B_4 = I$  as well then

$$|\text{tr}(A_i B_j)| = -[\text{tr}(A_1 A_2 A_3) - \text{tr}(A_1 A_3 A_2)][\text{tr}(B_1 B_2 B_3) - \text{tr}(B_1 B_3 B_2)].$$

Changing columns in  $|\text{tr}(A_i B_j)|$  and applying the identity of Example 1, we see that  $p$  vanishes for an arbitrary assignment of  $2 \times 2$  matrices. As in Lemma 6, it is clearly an element of the appropriate group algebra. □

**Lemma 8.** (Compare [8, Proposition 4.8]).  $\Psi(q_2) = 0$ .

*Proof.* Create  $p \in \mathbb{C}[\mathcal{W}]$  by specializing the Procesi identity of Lemma 6 at  $x_1 = y_1 = a_1$ ,  $x_2 = y_2 = a_2$ ,  $x_3 = y_3 = a_i$  and  $x_4 = y_4 = a_j$ . Theorem 6 and Proposition 1 imply  $\Psi(p) = 0$ . Note that  $p$  and  $q_2$  are determinants of matrices that differ only along their diagonals. The differences between diagonal terms are of the form

$$2(a_m^2) - (a_m)^2 - t_m^2 + 4,$$

which can be rewritten as

$$2(a_m^2) - 2(a_m)^2 + 2(1) - 2(1) + 4.$$

Hence,  $q_2$  may be obtained from  $p$  by a finite sequence of substitutions of the form  $(a_m^2) = (a_m)^2 - (1)$  or  $(1) = 2$ . Each step involves a pair of polynomials whose difference is divisible either by  $(a_m a_m) + (a_m a_m^{-1}) - (a_m)(a_m^{-1})$  or by  $(1) - 2$ , so Lemma 2 implies  $\Psi(q_2) = \Psi(p) = 0$ .  $\square$

**Lemma 9.** (Compare [8, Proposition 4.9]).  $\Psi(q_3) = 0$ .

*Proof.* Specialize the identity of Lemma 6 at  $x_1 = y_1 = a_1$ ,  $x_2 = y_2 = a_2$ ,  $x_3 = y_3 = a_3$ ,  $x_4 = a_i$  and  $y_4 = a_j$ . Then proceed as in Lemma 8.  $\square$

**Lemma 10.** (Compare [8, Corollary 4.12]).  $\Psi(q_4) = 0$ .

*Proof.* Specialize the Procesi identity of Lemma 7 at  $y_1 = a_i$ ,  $y_2 = a_j$ ,  $y_3 = a_k$  and  $x_m = a_m$  for  $m = 1, 2, 3$ . If  $i > 3$  this is precisely  $q$ . If not, then proceed as in Lemmas 8 and 9, using the fact that  $t_{mm}$  denotes  $t_m^2 - 2$ .  $\square$

This is enough to prove the main theorem for  $H$ , but we may as well consider an arbitrary compact, orientable 3-manifold. Let  $M$  be the result of adding 2-handles to  $H$  along curves  $\{c_1, \dots, c_m\}$  in  $\partial H$ . Choose words  $w_i$  in  $\pi_1(H)$  so that, as a loop, each  $w_i$  is freely homotopic to some orientation of  $c_i$ . For each  $i$  and  $j$  form the polynomial  $p_{ij} = (w_i a_j) - (a_j) \in \mathcal{R}(H)$ . Using the obvious identification  $\mathcal{R}(H) \cong \mathbb{C}[\mathcal{T}_0]/\sqrt{\mathcal{J}_H}$ , create an ideal  $\mathcal{J}_M$  in  $\mathbb{C}[\mathcal{T}_0]$  generated by  $\mathcal{J}_H \cup \{p_{ij}\}$ .

**Theorem 9.** (González-Acuña–Montesinos).  $X(M)$  is the zero set of  $\mathcal{J}_M$  in  $\mathbb{C}[\mathcal{T}_0]$ .

It follows immediately that  $\mathcal{R}(M) = \mathbb{C}[\mathcal{T}_0]/\sqrt{\mathcal{J}_M}$ . We know that  $\Phi$  maps  $V(M)$  onto  $\mathcal{R}(M)$  and it is clear that  $\Psi$  maps  $\mathbb{C}[\mathcal{W}]$  onto  $V(H)$ , and hence onto  $V(M)$ . Using these maps, we can now see how  $V(M)$  compares to  $\mathcal{R}(M)$ . Henceforth, consider  $\Psi$  to be the restriction to  $\mathbb{C}[\mathcal{T}_0]$ .

**Proposition 4.**  $\mathcal{J}_M \subset \ker \Psi \subset \sqrt{\mathcal{J}_M}$

*Proof.* That  $\mathcal{J}_H$  lies in  $\ker \Psi$  is the content of Lemmas 5, 8, 9, and 10. To see that  $\Psi(p_{ij}) = 0$ , construct a knot  $K_{a_j}$  for each generator  $a_j$ . For each  $i$  and  $j$ , there



is a band sum  $c_i \#_b K_{a_j}$  producing a knot  $K_{w_i a_j}$ . Since  $c_i \#_b K_{a_j} \simeq K_{a_j}$  in  $M$ , we have  $\Psi(p_{ij}) = K_{a_j} - K_{w_i a_j} = 0$ .

For the second containment, note that Corollary 1 and Proposition 3 imply that  $\Psi|_{\mathbb{C}[T_0]}$  is still onto. It should now be clear that

$$\mathbb{C}[T_0] \xrightarrow{\Psi} V(M) \xrightarrow{\Phi} \mathcal{R}(M) \cong \mathbb{C}[T_0]/\sqrt{\mathcal{J}_M}$$

is the canonical projection.  $\square$

There are various phrasings of the immediate implication of Proposition 4.

**Theorem 10.** *Let  $M$  be a compact orientable 3-manifold with  $\Phi$ ,  $\Psi$ ,  $\mathcal{J}_M$ , and  $T_0$  defined as above. Denote the ideal of nilpotents in  $V(M)$  by  $\sqrt{0}$ .*

- (1)  *$X(M)$  is the zero set of  $\ker \Psi$  in  $\mathbb{C}[T_0]$ .*
- (2)  *$\sqrt{\ker \Psi} = \sqrt{\mathcal{J}_M}$ .*
- (3)  *$\ker \Phi = \sqrt{0}$ .*
- (4)  *$\Phi$  induces an isomorphism  $\hat{\Phi} : V(M)/\sqrt{0} \rightarrow \mathcal{R}(M)$ .*
- (5)  *$\Psi$  induces an isomorphism  $\hat{\Psi} : \mathbb{C}[T_0]/\sqrt{\mathcal{J}_M} \rightarrow V(M)/\sqrt{0}$ .*
- (6) *Under the identification of  $\mathcal{R}(M)$  with  $\mathbb{C}[T_0]/\sqrt{\mathcal{J}_M}$ , the maps  $\hat{\Psi}$  and  $\hat{\Phi}$  are inverses.*

*Proof.*

- (1) This is immediate from Proposition 4 and the fact the  $X(M)$  is the zero set of both  $\mathcal{J}_M$  and  $\sqrt{\mathcal{J}_M}$ .
- (2) Nullstellensatz.
- (3) Since  $\mathcal{R}(M)$  cannot, by definition, contain a non-zero nilpotent element,  $\Phi(\sqrt{0}) = 0$ . Suppose now that  $\Phi(\alpha) = 0$ , and write  $\alpha$  as  $\Psi(\beta)$ . We have seen that

$$\mathbb{C}[T_0] \xrightarrow{\Psi} V(M) \xrightarrow{\Phi} \mathcal{R}(M) \cong \mathbb{C}[T_0]/\sqrt{\mathcal{J}_M}$$

is the canonical projection. Hence,  $\beta \in \sqrt{\mathcal{J}_M}$ . It follows from Theorem 10(2) that  $\Psi(\beta^n) = 0$  for some  $n$ , meaning  $\alpha$  is nilpotent.

- (4) Theorem 3 and Theorem 10(3).
- (5) The composition

$$\mathbb{C}[T_0] \xrightarrow{\Psi} V(M) \xrightarrow{\pi} V(M)/\sqrt{0} \xrightarrow{\hat{\Phi}} \mathcal{R}(M) \cong \mathbb{C}[T_0]/\sqrt{\mathcal{J}_M}$$

is the canonical projection. (Here  $\pi$  is also projection). Hence,  $\ker(\pi \circ \Psi) = \sqrt{\mathcal{J}_M}$ .

- (6) It is easy to see that both

$$\mathbb{C}[T_0]/\sqrt{\mathcal{J}_M} \xrightarrow{\hat{\Psi}} V(M)/\sqrt{0} \xrightarrow{\hat{\Phi}} \mathcal{R}(M) \cong \mathbb{C}[T_0]/\sqrt{\mathcal{J}_M}$$

and

$$V(M)/\sqrt{0} \xrightarrow{\widehat{\Phi}} \mathcal{R}(M) \cong \mathbb{C}[\mathcal{T}_0]/\sqrt{\mathcal{J}_M} \xrightarrow{\widehat{\Psi}} V(M)/\sqrt{0}$$

are the identity.  $\square$

We conclude with an application. The author would like to thank Charles Frohman for suggesting that this result might follow quickly, Victor Camillo for encouraging us to disregard nilpotents, and Bernadette Mullins for pointing out the result from ring theory used in the proof. Recall that a 3-manifold is *small* if it contains no incompressible, non-boundary parallel surface.

**Theorem 11.** (Compare [2, Corollary 1]). *If  $M$  is small then  $\dim V(M) < \infty$ .*

*Proof.* Suppose that  $X(M)$  has positive dimension. It follows that some component of  $X(M)$  contains a curve whose smooth projective resolution has an ideal point. From [5, 2.2.1] we then have a non-trivial splitting of  $\pi_1(M)$ , meaning  $M$  is not small. Hence,  $X(M)$  consists of a finite set of points and  $\mathcal{R}(M)$  is finite dimensional as a vector space. It is a standard result of commutative algebra that an ideal in a Noetherian ring contains some power of its radical. Thus, from Theorem 10(2), we obtain

$$\left(\sqrt{\mathcal{J}_M}\right)^n \subset \ker \Psi \subset \sqrt{\mathcal{J}_M}$$

for some  $n$ . Since  $\mathcal{R}(M) \cong \mathbb{C}[\mathcal{T}_0]/\sqrt{\mathcal{J}_M}$ , it is a simple exercise to show that  $\mathbb{C}[\mathcal{T}_0]/(\sqrt{\mathcal{J}_M})^n$  is finite dimensional. The result now follows from the fact that  $V(M) \cong \mathbb{C}[\mathcal{T}_0]/\ker \Psi$ , which in turn is the homomorphic image of  $\mathbb{C}[\mathcal{T}_0]/(\sqrt{\mathcal{J}_M})^n$ .  $\square$

## References

- [1] D. Bullock, *A finite set of generators for the Kauffman bracket skein algebra*, preprint.
- [2] D. Bullock, Estimating a skein module with  $SL_2(\mathbb{C})$  characters, *Proc. Amer. Math. Soc.*, to appear.
- [3] D. Bullock, Estimating the states of the Kauffman bracket skein module, *Proc. Banach Center Mini-Semester on Knot Theory*, to appear.
- [4] D. Bullock and J. H. Przytycki, Kauffman bracket skein module quantization of symmetric algebra and  $so(3)$ , in preparation.
- [5] M. Culler and P. Shalen, Varieties of group representations and splittings of 3-manifolds, *Ann. Math.* **117** (1983), 109–146.
- [6] R. Fricke and F. Klein, *Vorlesungen über die Theorie der automorphen Functionen*, Vol. 1, B. G. Teubner, Leipzig 1897.
- [7] W. Fulton and J. Harris, *Representation theory: a first course*, Springer Verlag, New York 1991.
- [8] F. González-Acuña and J. M. Montesinos, On the character variety of group representations in  $SL(2, \mathbb{C})$  and  $PSL(2, \mathbb{C})$ , *Math. Z.* **214** (1993), 627–652.

- [9] R. Horowitz, Characters of free groups represented in the two dimensional linear group, *Comm. Pure Appl. Math.* **25** (1972), 635–649.
- [10] J. Hoste and J. H. Przytycki, The  $(2, \infty)$ -skein module of lens spaces; a generalization of the Jones polynomial, *J. Knot Theory Ramifications* **2** no. 3 (1993), 321–333.
- [11] J. Hoste and J. H. Przytycki, The Kauffman bracket skein module of  $S^1 \times S^2$ , *Math. Z.* **220** (1995), 65–73.
- [12] U. Leron, Trace identities and polynomial identities of  $n \times n$  matrices, *J. Algebra* **42** (1976), 369–377.
- [13] W. Magnus, Rings of Fricke characters and automorphism groups of free groups, *Math. Z.* **170** (1980), 91–103.
- [14] C. Procesi, The invariant theory of  $n \times n$  matrices, *Adv. in Math.* **19** (1976), 306–381.
- [15] J. H. Przytycki, Skein modules of 3-manifolds, *Bull. Pol. Acad. Sci.* **39(1-2)** (1991), 91–100.
- [16] Ju. P. Razmyslov, Trace identities of full matrix algebras over a field of characteristic zero, *Izv. Akad. Nauk. SSSR Ser. Mat.* **38** (1974), 723–756 (Russian). Engl. Transl.: *Math. USSR-Izv.* **8** (1974), 727–760.
- [17] H. Vogt, Sur les invariants fondamentaux des équations différentielles linéaires du second ordre, *Ann. Sci. Écol. Norm. Supér., III. Sér.* **6** (1889), 3–72.

Doug Bullock  
Boise State University  
Boise, ID 83725, USA  
e-mail: bullock@math.idbsu.edu

(Received: April 18, 1996)