

A new occurrence of *Schwagerina princeps* in Sumatra

Autor(en): **Ozawa, Y.**

Objekttyp: **Article**

Zeitschrift: **Eclogae Geologicae Helvetiae**

Band (Jahr): **22 (1929)**

Heft 1

PDF erstellt am: **18.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-158834>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*

ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

A New Occurrence of *Schwagerina princeps* in Sumatra.

By Y. OZAWA (Tokyo).

By the courtesy of Dr. A. TOBLER, of the Museum of Natural History of Basle, I had an opportunity to examine a large number of thin sections of Fusulina limestones collected by him during his expedition to Sumatra. Some of these sections were described by E. LANGE¹⁾, but most of them are untouched. I have re-examined the specimens described by LANGE. And a revision of his species belonging to the Fusulinidae will be published in the near future. Among the untouched specimens there are some interesting species which are not hitherto known from Sumatra. These are *Schwagerina princeps* EHRENB., *Neoschwagerina craticulifera* SCHWAGER and *Fusulina japonica* GÜMBEL²⁾.

Schwagerina princeps is found in the Productus limestone of Telók Gedáng, laag d³⁾ on the Merangin River underlying the plant bed of Soengi Garing³⁾ containing a *Pecopteris* flora. Its occurrence there is very important because of inconsistency of opinion concerning the age of the Productus limestone and the plant bed.

The fossils in the Productus limestone were studied and described by O. E. MEYER⁴⁾ who determined its age as Upper Permian (Neodyas) by an occurrence of *Productus sumatrensis*. Later, W. J.

¹⁾ E. LANGE, Eine mittelpermische Fauna von Guguk Bulat (Padanger Oberland). Nr. 5 der „Beiträge zur Geologie und Paläontologie von Sumatra, herausgegeben von AUG. TOBLER, Basel“. Verh. v. h. Geol.-mijnbouwk. Genootschap voor Nederland en Koloniën. Geol. serie, deel 7, 1925.

²⁾ The latter two species were obtained from other localities as *Schwagerina princeps*.

³⁾ Refer to A. TOBLER, Djambi-Verslag. Jaarboek v. h. Mijnwezen in Nederl. Oost-Indië, 48 Jaarg. 1919. (1922) p. 191, 306, 541 (Telók Gedáng), p. 193, 308, 541 (Soengi Garing); bijlage VI, speciaalkaart 3; bijlage VII, profil 4, and A. TOBLER, Unsere paläontologische Kenntnis von Sumatra. Eclogae geol. Helv., Vol. 18, 1923, p. 317.

⁴⁾ O. E. MEYER, Brachiopoden des Perm und Untercarbon der Residentschaft Djambi (Sumatra). Nr. 2 der „Beiträge zur Geologie und Paläontologie von Sumatra, herausgegeben von AUG. TOBLER, Basel“. Verh. v. h. Geol.-mijnbouwk. Genootschap voor Nederland en Koloniën. Geol. serie, deel 5, 1922.

JONGMANS and W. GOTCHAN¹⁾ examined the plant fossils obtained from the bed overlying the Productus limestone and insisted that its age is the uppermost Carboniferous.

On account of an occurrence of *Schwagerina princeps* in the Productus limestone I cannot hold the opinion of MEYER, because *Schwagerina princeps* marks a definite horizon in the Permian-Carboniferous series in Asia and is generally considered as indicating the uppermost Carboniferous though some authors, such as E. HAUC and H. GERTH, consider it to be of Lower Permian age.

On the other hand, the rock succession of the Permian and Carboniferous formations in the Merangin district more or less resembles that in North China and Korea where it was surveyed in detail in several coal fields. Many important papers concerning the stratigraphy of coal bearing Permian-Carboniferous formation were published in China and Japan. The generalised succession²⁾ of coal bearing formation in North China and Korea is as follows (arranged in the descending order):

3. Coal bearing complex containing abundant plant fossils which were studied by H. YABE and others and determined to be Rotliegendes.
2. A complex of sandstone, shale and limestone. The uppermost limestone generally contains *Schwagerina princeps* associated with *Fusulina* spp., which are decidedly older than *Neoschwagerina*.
1. The fire clay (weathered substance of Ordovician limestone).

Comparing the above succession with that of the coal bearing formation in the Merangin district of Sumatra, we are struck by the similarity of their general succession. The plant bed in Sumatra as determined by JONGMANS and GOTCHAN is of the uppermost Carboniferous, therefore it is natural to correlate the Productus limestone of the Merangin district with the *Schwagerina princeps* limestone in North China and Korea. And then we can safely conclude that the Permian-Carboniferous succession in Sumatra is of the same type as that of North China and Korea.

¹⁾ W. J. JONGMANS und W. GOTCHAN, Beiträge zur Kenntnis der Flora des Obercarbons von Sumatra. Nr. 7 der „Beiträge zur Geologie und Paläontologie von Sumatra, herausgegeben von AUG. TOBLER, Basel“. Verh. v. Geol.-mijnbouwk. Genootschap voor Nederland en Koloniën. Geol. serie, deel 8, 1925.

²⁾ Refer to Y. OZAWA's paper on "Some Carboniferous Fossils collected in Manchuria and Korea", Japanese Journal of Geology and Geography, vol. 5, no. 3, 1927.