

Plates

Autor(en): **[s.n.]**

Objekttyp: **Appendix**

Zeitschrift: **Eclogae Geologicae Helvetiae**

Band (Jahr): **87 (1994)**

Heft 1

PDF erstellt am: **17.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Plate 1

- Fig. 1. Wackestone with large calcareous sponge spicules and *Lingulonodosaria* cf. *ovalis*. Sample VD 514 (Carixian). $\times 80$.
- Fig. 2. Wackestone with large recrystallized echinoid remains and *Paralingulina* cf. *pupiformis* (axial lateral view). S. VD 510 (Carixian). $\times 95$.
- Fig. 3. Echinoid remains and calcareous sponge spicules. Same sample. $\times 10$.
- Fig. 4. High-spired sculptured gastropods. S. VD 499 (Carixian). $\times 95$.
- Fig. 5. Finely agglutinated *Ammodiscidae* (*Usbekistaniinae*?) with a calcitic cement. S. VD 530 (Carixian). $\times 85$.
- Fig. 6. *Planiinvoluta* aff. *P. carinata* Leischner. S. VD 525 (Carixian). $\times 85$.
- Fig. 7. *Verneulinidae*. S. VD 511 (Carixian). $\times 90$.
- Fig. 8. *Miliolidae* (*Quinqueloculina*). S. VD 519 (Carixian). $\times 105$.
- Fig. 9. *Planiinvoluta* aff. *P. carinata*. S. VD 521 (Carixian). $\times 85$.
- Fig. 10. *Agerina martana*. S. VD 511 (Carixian). $\times 100$.
- Fig. 11. *Ophthalmidium* sp.. S. VD 521 (Carixian). $\times 85$.
- Fig. 12. *Paralingulina tenera tenera* (transverse section). S. VD 527 (Carixian). $\times 70$.
- Fig. 13 and 14. *Paralingulina* cf. *pupiformis* (lateral and frontal sections). S. VD 511 (Carixian). $\times 85$.
- Fig. 15. Thick ostracod carapace (transverse section). S. VD 495 (Lower Domerian). $\times 45$.
- Fig. 16. *Icthyolaridae*. S. VD 491 (Middle Domerian). $\times 70$.
- Fig. 17. *Repmanina charoides* corona (siliceous test). S. 492 (Middle Domerian). $\times 105$.
- Fig. 18. *Ammodiscus* cf. *siliceous* (siliceous test). S. VD 493 (Middle Domerian). $\times 80$.
- Fig. 19. Mudstone-wackestone yielding ammonite nuclei, small gastropods and echinoid remains. S. VD 498 (Carixian/Domerian transition). $\times 10$.
- Fig. 20. Bioturbated mudstone, wackestone with ostracods, lenticulinas and echinoderm remains. S. VD 494.4 (Lower/Middle Domerian). $\times 10$.




Plate 1: Microfacies 1 (Fig. 1, 2, 3, 5-14) and 2 (Fig. 4, 15-20).

Plate 2

- Fig. 1. Planar-bedded and densely packed echinoderm and peloidal calcarenite. Sample VD 399 (Lower Toarcian). $\times 25$.
- Fig. 2. Fine-grained packstone. S. VD 399.6 (Lower Toarcian). $\times 25$.
- Fig. 3. Grainstone with coated grains and *Planiinvoluta* (?). S. VD 389 (Lower Toarcian). $\times 35$.
- Fig. 4. Grainstone with coated grains and fecal pellets. S. VD 375.1 (Lower Toarcian). $\times 35$.
- Fig. 5. Recrystallized bioclastic packstone containing large *lituolidae* and echinoid fragments. S. VD 389 (Lower Toarcian). $\times 30$.
- Fig. 6. Green alga bioclast. Same sample. $\times 25$.
- Fig. 7. *Ataxophragmiidae*, *Planiinvoluta* (?) and other bioclasts. Same sample. $\times 30$.
- Fig. 8. Graded wackestone-packstone with radiolarians and flat, thin bivalves. S. VD 378.8 (Lower Toarcian). $\times 20$.
- Fig. 9. Grainstone containing ooids, peloids and skeletal fragments. S. VD 389 (Lower Toarcian). $\times 35$.
- Fig. 10. Calcified radiolarians and pyritized grains. S. VD 363 (Lower Toarcian). $\times 115$.
- Fig. 11 and 12. *Planiinvoluta* sp. S. VD 375.10 (Lower Toarcian). $\times 70$.
- Fig. 13. *Planiinvoluta* sp. (attached side of the right). S. VD 387 (Lower Toarcian). $\times 100$.
- Fig. 14. Radiolarian packstone. S. VD 368 (Lower Toarcian). $\times 25$.




Plate 2: Microfacies 3 (Fig. 1 and 2), 4 (Fig. 3–9, 11–13) and 5 (Fig. 10 and 14).

Plate 3

- Fig. 1. Bivalve and peloidal packstone. Sample VD 189 (Middle Toarcian). $\times 80$.
- Fig. 2. *Planiinvoluta* aff. *P. carinata*. S. VD 204 (Lower Toarcian). $\times 85$.
- Fig. 3. *Planiinvoluta* (?). S. VD 180 (Upper Toarcian). $\times 85$.
- Fig. 4. Peloidal grainstone with *Earlandia* sp. S. VD 189 (Middle Toarcian). $\times 100$.
- Fig. 5 and 6. Wackestone-packstone containing bivalve shells, small gastropods, echinoderm remains and ammonite nuclei. Geopetal fabrics (shelter porosity) are common below convexly embedded shells. S. VD 177 (Upper Toarcian). $\times 15$.
- Fig. 7, 8 and 9. Wackestone-packstone similar to the sample VD 177. Radiolarians are common. S. VD 178 (Upper Toarcian). $\times 15$.
- Fig. 10. Bivalve packstone containing large (1 to 3 cm) intraclast with radiolarian and bivalve remains. S. VD 174,5 (Aalenian). $\times 10$.
- Fig. 11. Densely packed flat bivalve deposits. S. VD 161,5 (Aalenian). $\times 10$.




Plate 3: Microfacies 6 (Fig. 1, 3 and 4), 8 (Fig. 5-9) and 9 (Fig. 10 and 11).

Plate 4

- Fig. 1. *Jaculella liasica*. Sample VD 496.18 (Lower Domerian). $\times 111$.
- Fig. 2. *Repmania charoides corona*. S. VD 498 (Carixian/Domerian transition). $\times 200$.
- Fig. 3. *Glomospira variabilis*. S. VD 496.18 (Lower Domerian). $\times 121$.
- Fig. 4. *Jaculella anulata* (side view). S. VD 488.9 (Upper Domerian). $\times 64$.
- Fig. 5. Same specimen (oblique apertural view).
- Fig. 6. *Glomospirella gaultina*. S. VD 496.18 (Lower Domerian). $\times 127$.
- Fig. 7. *Glomospirella* sp. 1. S. VD 488.9 (Upper Domerian). $\times 121$.
- Fig. 8. *Prodentalina* cf. *constricta* (side view). S. VD 373 (Lower Toarcian). $\times 140$.
- Fig. 9. Same specimen (apertural view).
- Fig. 10. *Paralingulina tenera praepupa* (oblique apertural view). S. VD 373 (Lower Toarcian).
- Fig. 11. Same specimen (side view) $\times 147$.
- Fig. 12. *Tristix liasina* (side view). S. VD 372.5 (Lower Toarcian). $\times 272$.
- Fig. 13. Same specimen (lateral face view).
- Fig. 14. *Prodentalina compressa* (side view). S. VD 371 (Lower Toarcian). $\times 123$.
- Fig. 15. Same specimen (front view).
- Fig. 16. Same specimen (dorso-apertural view).
- Fig. 17. *Lingulonodosaria dentaliniformis* (side view). S. VD 373 (Lower Toarcian). $\times 126$.
- Fig. 18. Same specimen (oblique apertural view).



Plate 4: Carixian, Domerian and Lower Toarcian (Tenuicostatum Zone) microforaminifera.

Plate 5

Fig. 1. *Reinholdella planiconvexa* (dorsal view). S. VD 360.9 (Lower Toarcian, Tenuicostatum Zone). $\times 131$.

Fig. 2. Same species (side view), second specimen. Same sample.

Fig. 3. Same species (ventral view), third specimen. Same sample.

Fig. 4. *Conicospirillina* sp. (dorsal view). S. VD 185.2 (Middle Toarcian, Erbaense Zone). $\times 130$.

Fig. 5. Same specimen (edge view).

Fig. 6. Same specimen (ventral view).

Fig. 7. *Ammobaculites* cfr. *vetusta*, S. VD 185.3 (Middle Toarcian, Erbaense Zone). $\times 128$.

Fig. 8. *Haplophragmoides kingakensis* (side view). S. VD 185.3 (Middle Toarcian, Erbaense Zone). $\times 61$.

Fig. 9. Same specimen (apertural view).

Fig. 10. *Lenticulina* (*Astacolus*) *d'orbignyi* (side view). S. VD 185.5 (Middle Toarcian, Erbaense Zone). $\times 55$.

Fig. 11. Same specimen (edge view).

Fig. 12. *Astacolus* cf. *stilla*. S. VD 185.5 (Middle Toarcian, Erbaense Zone). $\times 38$.

Fig. 13. *Astacolus* gr. *pennensis*. Same sample $\times 67$.

Fig. 14. *Vaginulinopsis deeckei parallela*. S. VD 185.3 (Middle Toarcian, Erbaense Zone). $\times 62$.

Fig. 15. *Vaginulinopsis* sp. (edge view). S. VD 185.5 (Middle Toarcian, Bifrons Zone).

Fig. 16. Same specimen (side view). $\times 52$.

Fig. 17. *Falsopalmula tenuistriata*. S. VD 177 (Lower Aalenian). $\times 54$.

Fig. 18. *Astacolus dictyoides* (side view). Same sample. $\times 62$.

Fig. 19. Same specimen (oblique apertural view).

Fig. 20. *Lenticulina* (*Astacolus*) *d'orbignyi*. S. VD 181.3 (Upper Toarcian, Meneghinii Zone). $\times 51$.

Fig. 21. *Astacolus similis*. S. VD 192.8 (Middle Toarcian, Bifrons Zone). $\times 107$.

Fig. 22. *Astacolus mutabilis*. S. VD 184.5 (Middle/Upper Toarcian, Erbaense). $\times 66$.




Plate 5: Toarcian microforaminifers.

Plate 6

Fig. 1. “Cigar-shaped” *Lenticulina münsteri* (axial view). Sample VD 184.3 (Upper Toarcian, Erbaense Zone). $\times 45$.

Fig. 2. Same specimen (oblique view).

Fig. 3. Abraded *Lenticulina* sp. Sample VD 184.3 (Upper Toarcian, Erbaense Zone). $\times 73$.

Fig. 4. “Cigar-shaped” *Lenticulina* sp.. Same sample. $\times 60$.

Fig. 5. *Spiriferiina* (?) (brachial valve). S. VD 185.3 (Middle Toarcian, Erbaense Zone). $\times 30$.

Fig. 6. *Camptocythere toarciana* (left side). S. VD 184.5 (Middle/Upper Toarcian, Erbaense Zone). $\times 50$.

Fig. 7. *Procytherura multicostata* (left side). Same sample. $\times 190$.

Fig. 8. Low-trochospiral gastropod. S. VD 186.5 (Middle Toarcian, Brifrons Zone). $\times 100$.

Fig. 9. Bivalve (*Posidonidae*). S. VD 185.3 (Middle Toarcian, Erbaense Zone). $\times 30$.

Fig. 10. *Stenestroemia* (?) sp. (left side). S. VD 373 (Lower Toarcian, Tenuicostatum Zone). $\times 240$.

Fig. 11. *Monoceratina* sp (left side). Same sample. $\times 154$.




Plate 6: Abraded *Lenticulina* and other organisms.

