

### 3. An imbedding theorem

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **14 (1968)**

Heft 1: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **29.04.2024**

#### **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

#### **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

### 3. AN IMBEDDING THEOREM

*Lemma 3.1.* If  $X$  is a compact complex manifold and  $S$  a coherent analytic sheaf over  $X$ , then  $\Gamma(X, S)$  is a finite dimensional vector space (cf. remark concerning Theorem 2.2).

We will now prove an imbedding theorem (cf. [2], p. 343).

*Theorem 3.2.* If the complex manifold  $X$  is compact, connected, and carries a positive (negative) line bundle, then  $X$  can be imbedded biholomorphically in a complex projective space  $\mathbf{P}^N$ .

*Proof:* Suppose  $F$  is a line bundle on a compact complex manifold  $X$  with the property that for every  $a \in X$  there exists a section  $\sigma \in \Gamma(X, \underline{F})$  with  $\sigma(a) \neq 0$ . Then  $F$  defines a holomorphic mapping of  $X$  into a projective space  $\mathbf{P}^k$  in the following way:

Since  $X$  is compact,  $\Gamma(X, \underline{F})$  is finite-dimensional according to Lemma 3.1.

Let  $\sigma_0, \dots, \sigma_k$  be a basis of  $\Gamma(X, \underline{F})$ . Then the  $\sigma_j$  have no common zeros.

Since  $F$  is locally isomorphic to the product of an open subset of  $X$  and  $\mathbf{C}$ , the  $\sigma_j$  are locally given by holomorphic functions without common zeros.

We map  $X$  into  $\mathbf{P}^k$  by  $x \rightarrow (\sigma_0(x), \dots, \sigma_k(x))$ . The point in the projective space is independent of the isomorphism we are using, for if we use another isomorphism we get a point  $(g(x)\sigma_0(x), \dots, g(x)\sigma_k(x))$ , where  $g(x) \neq 0$  (cf. (1.1)).

We are now going to show that if  $F$  is positive, then there exists an integer  $\gamma$  such that the sections of  $\Gamma(X, \underline{F}^\gamma)$  have no common zeros and such that the corresponding mapping is an imbedding.

For  $a \in X$ , let  $I$  be the sheaf of germs of holomorphic functions vanishing at  $a$ . Since  $I$  is coherent, we can apply the vanishing theorem of Kodaira. We conclude that there exists an integer  $k(a)$  such that  $H^1(X, I \otimes \underline{F}^{k \geq k(a)}) = 0$ .

Since  $\mathcal{O}_a/I_a \approx \mathbf{C}$ , we have the following exact sequence

$$0 \rightarrow I \rightarrow \mathcal{O}(X) \rightarrow \mathbf{C}_a \rightarrow 0,$$

where  $\mathbf{C}_a$  is a sheaf with stalk  $\mathbf{C}$  at  $a$  and zero outside. From this it follows that the sequence

$$0 \rightarrow I \otimes \underline{F}^{k(a)} \rightarrow \underline{F}^{k(a)} \rightarrow \mathbf{C}_a \otimes \underline{F}^{k(a)} \rightarrow 0$$

is exact. We have  $\mathbf{C}_a \otimes \underline{F}^{k(a)} \approx \tilde{F}_a^{k(a)}$ , where  $\tilde{F}_a^{k(a)}$  has stalk  $F_a^{k(a)}$  at  $a$  and

zero outside. Using the fact that  $H^1(X, I \otimes \underline{F}^{k(a)}) = 0$ , the exact cohomology sequence associated to the above sequence of sheaves gives us an exact sequence

$$\Gamma(X, \underline{F}^{k(a)}) \rightarrow \Gamma(X, \tilde{F}_a^{k(a)}) \rightarrow 0.$$

This implies that given  $e \in F_a^{k(a)}$  there exists  $\sigma \in \Gamma(X, \underline{F}^{k(a)})$  such that  $\sigma(a) = e$ . Thus, for every  $a \in X$  we can find an integer  $k(a)$  and a neighbourhood  $V_a$  of  $a$  such that  $\Gamma(X, \underline{F}^{k(a)})$  has a section not vanishing on  $V_a$ . Since  $X$  is compact, there are finitely many such neighbourhoods  $V_i$  ( $i=1, \dots, p$ ) with corresponding sections of  $F^{k_i}$  such that  $X = \bigcup_{i=1}^p V_i$ . Letting  $k = k_1 \cdot k_2 \cdot \dots \cdot k_p$ , we get  $p$  elements of  $\Gamma(X, \underline{F}^k)$  without common zeros, for if  $\sigma \in \Gamma(X, \underline{F})$  and  $\sigma(x) \neq 0$ , then  $\sigma' = \underbrace{\sigma \otimes \dots \otimes \sigma}_{l \text{ times}} \in \Gamma(X, \underline{F}^l)$  and  $\sigma'(x) \neq 0$ .

Let  $\underline{E} = \underline{F}^k$ . Now, for  $a \in X$ , let  $G = q_a^2$ , where  $q_a$  is the ideal of germs of holomorphic functions vanishing at  $a$ . Using the above argument with  $\underline{E}$  and  $G$  instead of  $\underline{F}$  and  $I$ , we see that there exists an integer  $s(a)$  such that the restriction mapping

$$\Gamma(X, \underline{E}^{s(a)}) \rightarrow \{ \mathcal{O}_a / q_a^2 \} \otimes \underline{E}_a^{s(a)}$$

is surjective. Since the residue classes in  $\mathcal{O}_a / q_a^2$  are sets of germs  $f$  of holomorphic functions at  $a$  with fixed values of  $f(a)$  and  $df(a)$ , this implies that we can find a neighbourhood  $U_a$  of  $a$  and sections  $\sigma_1, \dots, \sigma_t \in \Gamma(X, \underline{E}^{s(a)})$  which are nowhere zero in  $U_a$  such that the mapping given by  $\sigma_1, \dots, \sigma_t$  is regular and injective in  $U_a$ . We observe that for every positive integer  $I$  we can find sections  $\sigma_1^{(1)}, \dots, \sigma_t^{(1)} \in \Gamma(X, \underline{E}^{Is(a)})$  which have the same properties in  $U_a$ . In fact, if  $\sigma$  is a section of  $\underline{E}^{s(a)}$  which has no zeros on a set  $M \subset X$ , we set

$$\sigma' = \sigma \otimes \dots \otimes \sigma, (l-1) \text{ times}.$$

Then  $\sigma' \otimes \sigma_1, \dots, \sigma' \otimes \sigma_t$  are sections of  $\underline{E}^{ls(a)}$ , and define the same mapping (at least on  $M$ ) as  $\sigma_1, \dots, \sigma_t$ .

We can cover  $X$  by finitely many such neighbourhoods  $U_1, \dots, U_r$ . If  $s' = s_1 \cdot \dots \cdot s_r$ , then there are elements of  $\Gamma(X, \underline{E}^{s'})$  which give a regular, injective mapping in each  $U_i$  ( $1 \leq i \leq r$ ).

We are now going to show that we can separate points in  $X$  by sections of a suitable  $\underline{E}^z$ . Let  $U = \bigcup_{i=1}^r (U_i \times U_i)$ . For  $(a, b) \in X \times X - U$ , let  $H$  be the sheaf of germs of holomorphic functions vanishing at  $a$  and  $b$ . It is

easily seen that the sequence

$$0 \rightarrow H \rightarrow \mathcal{O}(X) \rightarrow \mathbf{C}_a \oplus \mathbf{C}_b \rightarrow 0$$

is exact. From this we conclude as above that there exists an integer  $s(a, b)$  such that the sequence

$$\Gamma(X, \underline{E}^{s(a,b)}) \rightarrow E_a^{s(a,b)} \oplus E_b^{s(a,b)} \rightarrow 0$$

is exact. Therefore there exists a neighbourhood  $W$  of  $(a, b)$  in  $X \times X$  such that if  $(a', b') \in W$ , then the sections of  $\Gamma(X, \underline{E}^{s(a,b)})$  separate  $a'$  and  $b'$ ; that is, if  $\sigma_0, \dots, \sigma_k$  is a basis of  $\Gamma(X, \underline{E}^{s(a,b)})$ , then  $(\sigma_0(a'), \dots, \sigma_k(a'))$  and  $(\sigma_0(b'), \dots, \sigma_k(b'))$  are different points in  $\mathbf{P}^k$ . Let  $l$  be a positive integer, let  $(a', b') \in W$ , and let  $\sigma$  be a section of  $\Gamma(X, \underline{E}^{s(a,b)})$  such that  $\sigma(a') \neq 0$  and  $\sigma(b') \neq 0$ . Then  $\sigma^{l-1} \otimes \sigma_0, \dots, \sigma^{l-1} \otimes \sigma_k$  are sections of  $\Gamma(X, \underline{E}^{ls(a,b)})$  such that  $((\sigma^{l-1} \otimes \sigma_0)(a'), \dots, (\sigma^{l-1} \otimes \sigma_k)(a'))$  and  $((\sigma^{l-1} \otimes \sigma_0)(b'), \dots, (\sigma^{l-1} \otimes \sigma_k)(b'))$  are different points in  $\mathbf{P}^k$ .

This means that for every positive integer  $l$  the sections of  $\Gamma(X, \underline{E}^{ls(a,b)})$  separate all point pairs in  $W$ . Thus, covering  $X \times X - U$  by finitely many such neighbourhoods and taking  $s''$  to be the product of the corresponding  $s(a, b)$ , we find that the sections of  $\Gamma(X, \underline{E}^{s''})$  separate all point pairs in  $X \times X - U$ .

Let  $\alpha = s's''$  and let  $\sigma_0, \dots, \sigma_d$  be a basis of  $\Gamma(X, \underline{E}^\alpha)$ . We claim that the mapping  $f$  from  $X$  into  $\mathbf{P}^d$  defined by  $f(x) = (\sigma_0(x), \dots, \sigma_d(x))$  is a biholomorphic imbedding of  $X$  into  $\mathbf{P}^d$ . That this mapping is regular follows from the fact that  $\alpha$  is a multiple of  $s'$ . What remains to be proved is that the mapping is injective.

Suppose  $a, b \in X$ ,  $a \neq b$ . If  $(a, b) \in U$ , then  $a, b \in U_i$  for some  $i$ , and since  $\alpha$  is a multiple of  $s'$ , we have  $f(a) \neq f(b)$ . If  $(a, b) \in X \times X - U$ , then  $f(a) \neq f(b)$  since  $\alpha$  is a multiple of  $s''$ . This proves the theorem.

#### 4. LINE BUNDLE ASSOCIATED TO A DIVISOR

Let  $X$  be a complex manifold and  $D$  an analytic subset of  $X$  of pure codimension 1 at every point. Such a set  $D$  is called a *divisor* of  $X$ . We shall construct a line bundle  $F$  on  $X$ , associated to  $D$ .

To do this, we observe that every point of  $X$  has a neighbourhood  $U$  in which there is a holomorphic function  $s$  such that  $U \cap D = \{x \in U; s(x) = 0\}$ , and  $s$  generates, at every point of  $U$ , the ideal of germs of holomorphic functions vanishing on  $D$ . Thus we get a covering of  $X$  by open sets  $U_j$  and