

§5. k-SIMPLE ACYCLIC MAPS

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **25 (1979)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **28.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

(4.4) *Remark.* The group $\tilde{N} = \pi_1(A\tilde{X}_N)$ is a central extension of N (see the appendix) and, as $A\tilde{X}_N$ is acyclic, satisfies $H_1(\tilde{N}) = H_2(\tilde{N}) = 0$. Therefore $\tilde{N}$ is the universal central extension of N (see [K2]), namely one has the exact sequence $0 \rightarrow H_2(N) \rightarrow \tilde{N} \rightarrow N \rightarrow 1$. Therefore, if $f: X \rightarrow X'$ is a map such that $\pi_1(f)$ sends the perfect normal subgroup N of $\pi_1(X)$ isomorphically onto a normal subgroup N' of $\pi_1(X')$, then the induced map $Af: A\tilde{X}_N \rightarrow A\tilde{X}'_N$ induces an isomorphism on the fundamental groups.

§ 5. k -SIMPLE ACYCLIC MAPS

In this section we study acyclic maps having simplicity properties. The first proposition generalizes some results of Dror [D1, Lemma 3.4].

(5.1) PROPOSITION. *Let $f: X \rightarrow Y$ be a map of path connected spaces with $\pi_1(f)$ an isomorphism, and let N be a perfect normal subgroup of $\pi_1(X) = \pi$. If f induces an isomorphism $H_*(X, \mathbb{Z}[\pi/N]) \xrightarrow{\sim} H_*(Y, \mathbb{Z}[\pi/N])$ and an isomorphism $\pi_i(X) \xrightarrow{\sim} \pi_i(Y)$ for $i \leq k-1$, then*

- (1) $\pi_k(f): \pi_k(X) \rightarrow \pi_k(Y)$ is an epimorphism when N acts trivially on $\pi_k(Y)$, and
- (2) $\pi_k(f): \pi_k(X) \rightarrow \pi_k(Y)$ is an isomorphism when N acts trivially on $\pi_k(X)$ and $\pi_k(Y)$.

Proof. Let $F \rightarrow \tilde{X}_N$ be the homotopy fibre of the covering map $\tilde{f}: \tilde{X}_N \rightarrow \tilde{Y}_N$. By hypothesis it follows easily that $\tilde{f}$ induces an isomorphism on integral homology and on $\pi_i(X) \rightarrow \pi_i(Y)$ for $i \leq k-1$. From the Serre spectral sequence we have $H_0(\tilde{Y}_N, H_{k-1}(F)) = H_0(N, H_{k-1}(F)) = 0$. Since $H_{k-1}(F) = \pi_{k-1}(F)$ is a quotient of $\pi_k(Y)$ on which the perfect group N acts trivially, it follows that $\pi_{k-1}(F) = 0$, which proves (1).

Under the hypothesis of (2) we have $\pi_i(F) = 0$ for $i < k$ and $H_0(\tilde{Y}_N, H_k(F)) = H_0(N, \pi_k(F)) = 0$. Since N acts trivially on $\pi_k(X)$ the induced morphism $\pi_k(F) \rightarrow \pi_k(X)$ must be trivial, which proves the proposition.

The following lemma, proved in [D2, Lemma 2.6], follows easily from the homology exact sequence.

$$H_1(G, M'') \rightarrow H_0(G, M') \rightarrow H_0(G, M) \rightarrow H_0(G, M'') \rightarrow 0$$

(5.2) LEMMA. Let $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ be a short exact sequence of $\mathbb{Z}[G]$ -modules where G is a perfect group. Then M' and M'' are trivial G -modules if and only if M is a trivial G -module.

(5.3) DEFINITION. A space X is k -simple provided $\pi_1(X)$ acts trivially on $\pi_k(X)$. A map $f: X \rightarrow Y$ is k -simple provided $\ker \pi_1(f) \subset \pi_1(X)$ acts trivially on $\pi_k(X)$.

(5.4) PROPOSITION. Let $f: X \rightarrow Y$ be a map with homotopy fibre A where $\pi_1(A)$ is perfect. Then f is k -simple if and only if A is k -simple.

Proof. In the homotopy exact sequence of any fibration

$$\pi_{k+1}(Y) \rightarrow \pi_k(A) \rightarrow \pi_k(X) \rightarrow \pi_k(Y),$$

see the appendix, $\pi_1(A)$ acts trivially on $\text{im}(\pi_{k+1}(Y) \rightarrow \pi_k(A)) = M'$. If f is k -simple, then $\text{im}(\pi_1(A)) = \ker(\pi_1(f))$ acts trivially on $\pi_k(X)$. Hence $\pi_1(A)$ acts trivially on $M' \subset \pi_k(A)$ and on the quotient $\pi_k(A)/M'$. By (5.2), it acts trivially on $\pi_k(A)$.

Conversely, $\ker(\pi_1(f))$ acts trivially on $\ker(\pi_k(f)) \subset \pi_k(X)$ and trivially on $\pi_k(Y) \supset \text{im}(\pi_k(f))$. By (5.2), $\ker(\pi_1(f))$ acts trivially on $\pi_1(X)$. This proves the proposition.

(5.5) Notations. For a path connected space X and a perfect normal subgroup N of $\pi_1(X)$, we consider the following conditions:

(P_k). The group N acts trivially on $\pi_i(X)$ for $i \leq k$.

(H_k). The group N acts trivially on $H_i(\tilde{X})$ for $i \leq k$.

(5.6) PROPOSITION. For all natural numbers k we have that P_k implies H_k and H_k implies P_{k-1} . In particular, H_∞ and P_∞ are equivalent.

Proof. Consider the following commutative diagram where the rows and columns are fibrations.

$$\begin{array}{ccccc} T & \longrightarrow & A\tilde{X}_N & \longrightarrow & A(BN) \\ \downarrow & & \downarrow & & \downarrow \\ \tilde{X} & \longrightarrow & \tilde{X}_N & \longrightarrow & BN \\ \downarrow & & \downarrow & & \downarrow \\ F & \longrightarrow & \tilde{X}_N^+ & \longrightarrow & BN^+ \end{array}$$

By (5.4) condition P_k implies that $\pi_1(A\tilde{X}_N)$ acts trivially on $\pi_i(A\tilde{X}_N)$ for $i \leq k$. Since $A\tilde{X}_N$ and $A(BN)$ are both acyclic and $\pi_1(A\tilde{X}_N) \xrightarrow{\sim} \pi_1(A(BN))$ is an isomorphism (by (4.4)), we deduce using (5.1) that $\pi_i(A\tilde{X}_N) \rightarrow \pi_i(A(BN))$ is an isomorphism for $i \leq k$. Thus $\pi_i(T) = 0$ for $i \leq k-1$ and $\pi_k(T)$ is a trivial module $\pi_1(A(BN))$ since it is a quotient of $\pi_{k+1}(A(BN))$. On the other hand, we have $H_0(\pi_1(A(BN)), \pi_k(T)) = 0$ since $\pi_k(T) = H_k(T)$ and thus $H_*(A\tilde{X}_N) \rightarrow H_*(A(BN))$ is an isomorphism. Therefore, $\pi_k(T) = 0$ and $H_i(\tilde{X}) \rightarrow H_i(F)$ is an isomorphism for $i \leq k$. Hence P_k implies H_k since N acts trivially on $H_*(F)$.

Next, assume H_k holds. Then $H_i(\tilde{X}) \rightarrow H_i(F)$ is an isomorphism for $i \leq k$ by the comparison theorem for spectral sequences of fibrations with trivial actions. Since $\pi_1(F)$ is abelian, $\pi_1(F) = 0$ and $\pi_i(T) = 0$ for $i \leq k-1$. Hence $\pi_1(A\tilde{X}_N)$ acts trivially on $\pi_i(A\tilde{X}_N)$ for $i \leq k-1$. Using (5.4), we deduce P_{k-1} and the proposition.

(5.7) THEOREM. *Let $f: X \rightarrow Y$ be an acyclic map between CW-spaces which is k -simple for all $k \geq 2$ with $N = \ker \pi_1(f)$. Then the following is a fiber sequence*

$$\tilde{X} \rightarrow Y \xrightarrow{\alpha'} [B\pi_1(X)]_N^+$$

where α' is induced by $\alpha: X \rightarrow B\pi_1(X)$ as in (3.1) and $\pi_1(\alpha)$ is the identity.

Proof. As in the previous proposition, we have a diagram of fibrations

$$\begin{array}{ccccc} T & \longrightarrow & AX_N & \longrightarrow & A(BN) \\ \downarrow & & \downarrow & & \downarrow \\ \tilde{X} & \longrightarrow & \tilde{X}_N & \longrightarrow & BN \\ \downarrow & & \downarrow & & \downarrow \\ F & \longrightarrow & \tilde{Y} & \longrightarrow & BN^+ \\ & \searrow & \downarrow & & \downarrow \\ & & Y & \longrightarrow & [B\pi_1(X)]_N^+ \end{array}$$

We prove $\tilde{X} \rightarrow F$ is a homotopy equivalence with the same argument used in (5.6) to show P_k implies H_k . Since F is also the fibre of $X_N^+ \rightarrow [B\pi_1(X)]_N^+$ we have proved the theorem.

(5.8) *Remark.* Using (5.1), we see that for an acyclic map $f: X \rightarrow Y$ which is k -simple for all $k \geq 2$, the homotopy groups $\pi_*(Y)$ can be computed in terms of $\pi_*(X)$ and $\pi_*(B\pi_1(X)_N^+) \cong \pi_*(BN)^+$ for $i \geq 2$. Some computations of $\pi_*(BN^+)$ for a certain perfect group N can be found for instance in [H, Chapter 7].

§ 6. ACYCLIC MAPS INTO A GIVEN SPACE

In this section we study acyclic maps $f: X \rightarrow Y$ into a fixed space Y . Two such map $f: X \rightarrow Y$ and $f': X' \rightarrow Y$ are called equivalent provided there is a homotopy equivalence $h: X \rightarrow X'$ with $f \simeq f'h$. Let $AC(Y)$ denote the class of equivalence classes of acyclic $f: X \rightarrow Y$ over Y where X and Y are CW -spaces.

(6.1) DEFINITION. An extension data over a space Y is a triple (Φ, i, ϕ) where

- (a) Φ is an extension $1 \rightarrow N \rightarrow G \rightarrow \pi_1(Y) \rightarrow 1$ with N perfect,
- (b) $i: BG \rightarrow BG_N^+$ is an acyclic map with $\ker(\pi_1(i)) = N$ (whose equivalence class is well defined by (3.5)), and
- (c) $\phi: Y \rightarrow BG_N^+$ is a 2-connected map.

Two triples of extension data (Φ, i, ϕ) and (Φ', i', ϕ') are called equivalent provided there exists an isomorphism $g: G \rightarrow G'$ making the following diagrams commutative (up to homotopy for the second one).

$$\begin{array}{ccc}
 G & \xrightarrow{g} & G' \\
 \searrow & & \swarrow \\
 & \pi_1(Y) &
 \end{array}
 \qquad
 \begin{array}{ccc}
 BG & \xrightarrow{Bg} & BG' \\
 i \downarrow & & \downarrow i' \\
 BG_N^+ & \xrightarrow{Bg^+} & B(G')_N^+ \\
 \swarrow \phi & & \searrow \phi' \\
 & Y &
 \end{array}$$