

Appendix II The Monodromy Groups of the Minimal Hyperbolic Germs

Objektyp: **Appendix**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **25 (1979)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **28.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

The equivalence of Characterizations C2 and C8 is proved in [Wagreich 2, p. 66; Saito 2, Theorem 1.9]. In fact, the zero loci of the germs $\tilde{E}_6$, $\tilde{E}_7$ and $\tilde{E}_8$ have minimal resolution as above with $E^2 = -3$, -2 and -1 respectively.

Characterization C9. The germ $f^{-1}(0)$ is isomorphic to $g^{-1}(0)$, where g is a weighted homogeneous polynomial with weights w_i satisfying $w_0^{-1} + \dots + w_n^{-1} = n/2$.

The equivalence of Characterizations C2 and C9 is proved in [Saito 2, Satz 2.11]. In fact, the germs in Table 2b have the following weights:

GERM	WEIGHTS
P_8	(3, 3, 3)
X_9	(4, 4)
J_{10}	(3, 6)

APPENDIX II

THE MONODROMY GROUPS OF THE MINIMAL HYPERBOLIC GERMS

PROPOSITION. *The monodromy groups of the germs P_9 , X_{10} , and J_{11} have exponential growth.*

In this appendix, we present an (unpublished) proof of this proposition due to E. Looijenga. In fact, we will show that these groups have $PSL(2, \mathbf{Z})$ as subquotient (quotient of a subgroup). We let $O(V)$ denote the orthogonal group of a $\mathbf{Z}$ - or $\mathbf{R}$ -module V equipped with a bilinear form.

Suppose G is a polyhedral graph whose edges are weighed by non-zero integers. By convention, the weight 1 is omitted. Let L_G denote the free $\mathbf{Z}$ -module generated by the vertices $v_1, \dots, v_n$ of G . Define a symmetric bilinear form $(,)$ on L_G by setting $(v_i, v_i) = -2$, and $(v_i, v_j) = 0$ if there is no edge from v_i to v_j , otherwise equal to the weight on this edge. Conversely, given a symmetric integral bilinear form $(,)$ on a free module L with basis $\alpha_1, \dots, \alpha_n$ with the property that $(\alpha_i, \alpha_i) = -2$ for all i , one associates a graph to it in the obvious way.

For $\alpha \in L_G$, let s_α (reflection in α) be the isometry of L_G defined by

$$s_\alpha(\beta) = \beta - 2 \frac{(\alpha, \beta)}{(\alpha, \alpha)} \alpha$$

for $\beta \in L_G$. The reflection group $\mathcal{R}(G)$ of the graph G is defined to be the subgroup of $O(L_G)$ generated by $s_{\alpha_1}, \dots, s_{\alpha_n}$.

Example 1. Consider a reduced irreducible root system in a vector space V . Let $\alpha_1, \dots, \alpha_n$ be a collection of simple roots, let L be the free $\mathbb{Z}$ -module spanned by the α_i , and let $(,)$ be the negative of an invariant bilinear form [Serre, Chapter 5]. If $(\alpha_i, \alpha_i) = -2$ for all i , then the corresponding graph must be of type A_k, D_k, E_6, E_7 or E_8 . The reflection group of these graphs equals the Weyl group, the group generated by reflections in all the roots [Serre, p. V-16]. Furthermore, the reflection group together with the generators $s_{\alpha_1}, \dots, s_{\alpha_n}$ forms a Coxeter system [Bourbaki, p. 92]. (A Coxeter system is a group G , a collection of elements $g_1, \dots, g_n$ and a symmetric integral $n \times n$ matrix $\{m_{ij}\}$ with $m_{ii} = 1$ and $2 \leq m_{ij} \leq \infty$ for $i \neq j$, with the property that G is isomorphic to the free group with generators $g_1, \dots, g_n$ and relations $(g_i, g_j)^{m_{ij}} = 1$, for all i, j .)

Example 2. The monodromy group of a germ f is the reflection group of a quadratic form diagram for f (Sections 13 and 14). When this diagram is a tree (which is only possible for the simple germs), its reflection group together with the generators $T_1, \dots, T_\mu$ forms a Coxeter system. In general, this reflection group is not a Coxeter system [A'Campo 2, II, p. 403].


Lemma [Gabrielov 3]. If the graph G' is a subgraph of the graph G , then $\mathcal{R}(G')$ is a subquotient of $\mathcal{R}(G)$.

Proof. Let $\alpha'_1, \dots, \alpha'_m$ be a basis of $L_{G'}$ corresponding to the vertices of G' , let $\alpha_1, \dots, \alpha_m$ be the corresponding elements in L_G , and extend this to a basis $\alpha_1, \dots, \alpha_m, \alpha_{m+1}, \dots, \alpha_n$ of L_G corresponding to the vertices of G . The map $\alpha'_i \rightarrow \alpha_i$ is an isometric embedding of $L_{G'}$ in L_G . Let $\mathcal{R}'$ be the subgroup of $\mathcal{R}(G)$ generated by $s_{\alpha_1}, \dots, s_{\alpha_m}$; it has a presentation with these generators and certain relations. Any relation among these s_{α_i} holds also for $s_{\alpha_i} \mid L_{G'} = s_{\alpha'_i}$. Thus $\mathcal{R}'$ maps onto $\mathcal{R}(G')$.

Fact. If a subquotient of a group G has exponential growth, then so does G .


Proof of Proposition. 1. A quadratic form diagram for the germs P_9, X_{10} , and J_{11} is given in column 5 of Table 2. These graphs contain a

subgraph of the form $T_{3,3,4}$, $T_{2,4,5}$, and $T_{2,3,7}$ respectively, where $T_{p,q,r}$ is the graph




Hence it suffices to show that the reflection groups of these graphs have exponential growth.

Let Γ be the graph



with vertices corresponding to basis elements $\tilde{\alpha}, \alpha, \beta$ in L_Γ as indicated. We claim that $\mathcal{R}(T_{p,q,r})$ has $\mathcal{R}(\Gamma)$ as subquotient, for $(p, q, r) = (3, 3, 4)$, $(2, 4, 5)$, and $(2, 3, 7)$. Consider (for example) $T_{3,3,4}$, with vertices corresponding to basis elements $\alpha_i \in L_{T_{3,3,4}}$ as indicated:



This contains the graph E_6 . Let

$$\tilde{\alpha} = \alpha_1 + 2\alpha_2 + 3\alpha_3 + 2\alpha_4 + \alpha_5 + 2\alpha_6 \in L_{T_{3,3,4}}$$

be the largest root of E_6 [Bourbaki, p. 165]. Since all the roots of E_6 are the same length, $(\tilde{\alpha}, \tilde{\alpha}) = -2$. The lattice spanned by $\tilde{\alpha}, \alpha$, and β has diagram Γ . The reflections s_α and s_β are in $\mathcal{R}(T_{3,3,4})$. We claim that $s_{\tilde{\alpha}}$ is in $\mathcal{R}(T_{3,3,4})$ as well: The restriction $s_{\tilde{\alpha}}|_{L_{E_6}}$ is in $\mathcal{R}(E_6)$, since E_6 is a root system. Hence $s_{\tilde{\alpha}}|_{L_{E_6}} = (s_{\alpha_{i(1)}} \circ \dots \circ s_{\alpha_{i(m)}})|_{L_{E_6}}$ for some $1 \leq i(1), \dots, i(m) \leq 6$.

Also, $s_{\tilde{\alpha}}$ and $s_{\alpha_{i(1)}} \circ \dots \circ s_{\alpha_{i(m)}}$ are both the identity when restricted to the orthogonal complement of $L_{E_6} \otimes \mathbf{R}$ in $L_{T_{3,3,4}} \otimes \mathbf{R}$. Thus $s_{\tilde{\alpha}} = s_{\alpha_{i(1)}} \circ \dots \circ s_{\alpha_{i(m)}}$, and $\mathcal{R}(T_{3,3,4})$ contains $s_{\tilde{\alpha}}$. A proof similar to that of the lemma then shows that $\mathcal{R}(T_{3,3,4})$ has subquotient $\mathcal{R}(\Gamma)$.

3. Next we show that $\mathcal{R}(\Gamma)$ has subquotient $PSL(2, \mathbf{Z})$. This uses [E. Artin, Chapter V] heavily.

Let V be the 3-dimensional real vector space $L_{\Gamma} \otimes \mathbf{R}$. The bilinear form $(,)$ of Γ extends to V . This form is indefinite since $\tilde{\alpha} + \alpha$ has length 0. Let

$$0'(V) = \{f \in O(V) : \det f = 1 \text{ and spinor norm of } f \text{ equal to } 1 \mathbf{R}^{*2}\}.$$

Since V is indefinite, it is known [E. Artin, p. 200] that

$$(1) \quad 0'(V) \xrightarrow{\cong} PSL(2, \mathbf{R}).$$

Since $PSL(2, \mathbf{R})$ contains $PSL(2, \mathbf{Z})$ as a subgroup, the idea is to find elements of $\mathcal{R}(\Gamma) \subset O(\Gamma)$ which are in $0'(V)$ and map to generators of $PSL(2, \mathbf{Z})$. The standard generators of $PSL(2, \mathbf{Z})$ are

$$S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad T = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$$

with relations $S^2 = (ST)^3 = 1$. By inspection, it is found that the elements $s_{\tilde{\alpha}} s_{\beta}$ and $s_{\beta} s_{\alpha}$ of $O(L)$ satisfy $(s_{\tilde{\alpha}} s_{\beta})^2 = (s_{\beta} s_{\alpha})^3 = 1$, and have determinant equal to 1 and spinor norm equal to $1 \mathbf{R}^{*2}$. Therefore we would like to choose the isomorphism (1) such that $s_{\tilde{\alpha}} s_{\beta}$ maps to S , and $s_{\tilde{\alpha}} s_{\alpha} = (s_{\tilde{\alpha}} s_{\beta})^{-1} (s_{\beta} s_{\alpha})$ maps to $S^{-1} (ST) = T$.

The isomorphism (1) is done in two steps. First, let $D_0(V)$ be the elements of the Clifford algebra of V of norm 1; then [E. Artin, p. 199]

$$(2) \quad D_0(V)/\{\pm 1\} \xrightarrow{\cong} 0'(V).$$

We do not need to know exactly what this isomorphism is, but only that

$$v \circ w \rightarrow s_v s_w$$

for elements v, w in V regarded as a subspace of the Clifford algebra, and $v \circ w$ their product. Hence under the above isomorphisms

$$(3) \quad \frac{1}{2} \tilde{\alpha} \circ \beta \mapsto s_{\tilde{\alpha}} s_{\beta}, \quad \frac{1}{2} \tilde{\alpha} \circ \alpha \mapsto s_{\tilde{\alpha}} s_{\alpha}.$$

Secondly [E. Artin, p. 199],

$$(4) \quad D_0(V) / \{\pm 1\} \cong PSL(2, \mathbf{R}).$$

We examine this more closely. Let

$$A_1 = \sqrt{2}(\alpha + \tilde{\alpha} + \beta/2), \quad A_2 = \tilde{\alpha}/\sqrt{2}, \quad A_3 = \beta/\sqrt{2}.$$

Then A_1, A_2, A_3 is an orthogonal basis of V , and the matrix of $(,)$ with respect to this basis is the diagonal matrix $\langle +1, -1, -1 \rangle$. Let $C^+(V)$ be the subspace of the Clifford algebra of V spanned by the elements of even grading; $C^+(V)$ is generated by $1, i_1, i_2, i_3$, where $i_1 = A_2 \circ A_3$, $i_2 = A_3 \circ A_1$, and $i_3 = A_1 \circ A_2$, and has multiplication table as in [E. Artin, top of p. 200] with $a = -1$. The map

$$C^+(V) \rightarrow M(2, \mathbf{R})$$

(where $M(2, \mathbf{R})$ is the algebra of all 2×2 matrices over $\mathbf{R}$) defined by

$$\begin{aligned} 1 &\mapsto \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, & i_1 &\mapsto \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \\ i_2 &\mapsto \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, & i_3 &\mapsto \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix} \end{aligned}$$

is an isomorphism. (This is slightly different from the isomorphism of [E. Artin, p. 200].), and the restriction of this map to $D_0(V)$ gives the isomorphism (4). Furthermore,

$$(5) \quad \frac{1}{2} \tilde{\alpha} \circ \beta = i_1 \mapsto S, \quad \frac{1}{2} \tilde{\alpha} \circ \alpha = 1 - \frac{1}{2}(i_1 + i_3) \mapsto T$$

under this isomorphism. Combining isomorphisms (2) and (4) gives isomorphism (1), and (3) and (5) show that

$$s_{\tilde{\alpha}} s_{\beta} \mapsto S, \quad s_{\tilde{\alpha}} s_{\alpha} \mapsto T$$

under isomorphism (1). Thus $\mathcal{R}(\Gamma)$ maps onto $PSL(2, \mathbf{Z})$, and hence has $PSL(2, \mathbf{Z})$ as subquotient.

4. Finally, $PSL(2, \mathbf{Z})$ is isomorphic to the free product $(\mathbf{Z}/2\mathbf{Z}) * (\mathbf{Z}/3\mathbf{Z})$ [Serre, ch. 7; Lehner, p. 59], which has exponential growth.

TABLE 1 RATIONAL DOUBLE POINTS

(1) $f(x, y, z)$	(2) (p, q, r)	(3) Dual Graph of Resolution	(4) Name	(5) Subgroup G of S^3	(6) Order of G	(7) Weights
$x^{k+1} + y^2 + z^2$	$(1, 1, k)$		A_k $k \geq 1$	cyclic	$k+1$	$(k+1, 2, 2)$
$x^{k-1} + xy^2 + z^2$	$(2, 2, k-2)$		D_k $k \geq 4$	binary dihedral	$4(k-2)$	$(k-1, 2(k-1)/(k-2), 2)$
$x^3 + y^4 + z^2$	$(2, 3, 3)$		E_6	binary tetrahedral	24	$(3, 4, 2)$
$x^3 + xy^3 + z^2$	$(2, 3, 4)$		E_7	binary octahedral	48	$(3, 9/2, 2)$
$x^3 + y^5 + z^2$	$(2, 3, 5)$		E_8	binary icosahedral	120	$(3, 5, 2)$

Column 1 lists the germ $f(x, y, z)$. Column 3 lists the dual graph of the minimal resolution of $f^{-1}(0)$. The name of the graph is given in column 4. Each graph is of type $T_{p,q,r}$, where (p, q, r) are listed in column 2. Every vertex of the graph represents a nonsingular rational curve of self-intersection -2 . The analytic set $f^{-1}(0)$ is isomorphic to $\mathbb{C}^2/G$, where G is the finite subgroup of S^3 listed in column 5. Each germ f is weighted homogeneous, with weights as listed in column 7.

TABLE 2 GERMS OF LOW MILNOR NUMBER






(1)	(2) Name	(3) μ	(4) $f(z_0, \dots, z_n)$	(5) Quadratic form diagram
a (simple germs)	$A_k, k \geq 1$	k	z_0^{k+1}	
	$D_k, k \geq 4$	k	$z_0^{k+1} + z_0 z_1^2$	
	E_6	6	$z_0^3 + z_1^4$	
	E_7	7	$z_0^3 + z_0 z_1^3$	
	E_8	8	$z_0^3 + z_1^5$	
b (almost-simple germs)	P_8 or $\tilde{E}_6$	8	$z_0^3 + z_1^2 z_2 + a z_0 z_2^2 + b z_2^3, 4a^3 + 27b^2 \neq 0$	
	X_9 or $\tilde{E}_7$	9	$z_0 z_1 (z_0 - z_1) (z_0 - a z_1), a \neq 0, 1$	
	J_{10} or $\tilde{E}_8$	10	$z_0 (z_0 - z_1^2) (z_0 - a z_1^2), a \neq 0, 1$	

TABLE 2 (continuation)


c	P_9 or $T_{3,3,4}$	9	$z_0^2 z_2 + z_1^3 + z_1^2 z_2 + a z_2^4, a \neq 0$	
(minimal hyperbolic germs)	X_{10} or $T_{2,4,5}$	10	$z_0^4 + z_0^2 z_1^2 + a z_1^5, a \neq 0$	
	J_{11} or $T_{2,3,7}$	11	$z_0^3 + \varepsilon z_0^2 z_1^2 + a z_1^7, a \neq 0, \varepsilon = \pm 1$	

Table 2 is divided into parts a, b, and c as indicated in column 1. These parts are referred to in the text as Table 2a, etc. Column 4 contains the equation of the germ; to each equation must be added the quadratic form $z_r^2 + \dots + z_n^2$ in the variables $z_r, \dots, z_n$ not occurring in the equation. Column 2 gives the type of the germ in Arnold's notation, column 3 gives its Milnor number μ , and column 5 gives a quadratic form diagram.

TABLE 3
SIMPLIFICATION TABLE




Table 3 lists some (but not all) of the simplifications that occur among the germs of Table 2.

REFERENCES

- A'CAMPO, N. 1. La fonction zeta d'une monodromie. *Comment. Math. Helvetici* 50 (1975), pp. 233-248.
- 2. Le groupe de monodromie du déploiement des singularités isolées de courbes planes I. *Math. Ann.* 213 (1975), pp. 1-32. II. *Proc. Int. Cong. Math. Vancouver* (1974), Vol. I, pp. 395-404.
- ARNOLD, V. 1. Normal forms of functions near degenerate critical points, the Weyl groups A_k , D_k , E_k , and Lagrange singularities. *Funk. Anal.* 6 (1972), pp. 3-25. (Contains the classification of simple critical points.)
- 2. Classification of unimodal critical points of functions. *Funk. Anal.* 7 (1973), pp. 75-76.
- 3. Remarks on the stationary phase method and Coxeter numbers. *Russ. Math. Surveys* 28 (1973), pp. 19-48. (An excellent survey article, including the results of 1 and 2 above.)
- 4. Critical points of smooth functions. *Proc. Int. Cong. Math. Vancouver* (1974), Vol. I, pp. 19-39.
- 5. Classification of bimodal critical points of functions. *Funk. Anal.* 9 (1975), pp. 49-50.
- ARTIN, E. *Geometric Algebra*. Interscience, New York, 1957.
- ARTIN, M. On isolated rational singularities of surfaces. *Am. J. Math.* 88 (1966), pp. 129-136.
- BOURBAKI, N. *Eléments de mathématique*. Fasc. 34. *Groupes et algèbres de Lie*. Chaps. 4, 5, 6, *Actualités Sci. Indust.*, n° 1337, Hermann, Paris, 1968.
- BRIESKORN, E. 1. Über die Auflösung gewisser Singularitäten von holomorphen Abbildungen. *Math. Ann.* 166 (1966), pp. 76-102.
- 2. Rationale Singularitäten komplexer Flächen. *Inv. Math.* 4 (1968), pp. 336-358.
- 3. Singular elements of semi-simple algebraic groups. *Actes Cong. Int. Math.* (1970), pp. 279-284.
- 4. Die Monodromie der isolierten Singularitäten von Hyperflächen. *Manuscripta math.* 2 (1970), pp. 103-161.
- CARTAN, H. Quotient d'une espace analytique par un groupe d'automorphismes. In: *Algebraic geometry and topology*, pp. 90-102. Princeton Univ. Press, 1957.