

§3. Blowing Down

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **29 (1983)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **28.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

$$\begin{array}{ccc}
 V' & \xrightarrow{g} & V \\
 \gamma \downarrow & & \uparrow h \\
 V'' & \xrightarrow{h} & V
 \end{array}$$

$\mathcal{S}_{\text{Alg}}(V)$ is the set of distinct algebraic structures on V . Hence a natural problem is to compute $\mathcal{S}_{\text{Alg}}(V)$, or at least produce nontrivial elements of this set. For example if we take $M \subset V$ as in Proposition 2.10, then by Theorem 2.12 (V, M) is diffeomorphic to nonsingular algebraic sets (V', M') . Let $|V| = |V'|$ denote the underlying smooth structures and let $V \xrightarrow{g} |V|$, $V' \xrightarrow{g'} |V|$ be the forgetful maps. Then (V, g) and (V', g') are distinct elements of $\mathcal{S}_{\text{Alg}}(|V|)$, otherwise M would be isotopic to a nonsingular algebraic subset of V .

An interesting question is whether algebraic structures on smooth manifolds satisfy the product structure theorem; that is, whether the natural map

$$\mathcal{S}_{\text{Alg}}(M) \times \mathbf{R}^n \rightarrow \mathcal{S}_{\text{Alg}}(M \times \mathbf{R}^n), (V, g) \mapsto (V \times \mathbf{R}^n, g \times \text{id})$$

is surjection. The answer would be negative if one can find a smooth manifold M and $\theta \in H_*(M; \mathbf{Z}/2\mathbf{Z})$ such that M can not be diffeomorphic to a nonsingular algebraic set M' with $\theta \in H_*^A(M'; \mathbf{Z}/2\mathbf{Z})$. To see this, pick any smooth representative $N \xrightarrow{g} M$ of $\theta = g_*[N]$. By graphing g , we can assume $N \subset M \times \mathbf{R}^n$ for some n and g is induced by projection. By Theorem 2.12 we can find a diffeomorphism $\lambda: M \times \mathbf{R}^n \rightarrow V$ to a nonsingular algebraic set V with $\lambda(N)$ is an algebraic subset (one has to modify Theorem 2.12 to apply to this noncompact case). Then there can not exist a birational diffeomorphism $\mu: V \rightarrow M' \times \mathbf{R}^n$ where M' is a nonsingular algebraic set diffeomorphic to M , otherwise $\lambda(N) \xrightarrow{\mu} M' \times \mathbf{R}^n \xrightarrow{\text{projection}} M'$ would represent $\theta \in H_*^A(M'; \mathbf{Z}/2\mathbf{Z})$.

§3. BLOWING DOWN

Real algebraic sets obey some simple but useful topological properties:

PROPOSITION 3.1.

- (a) One point compactification an algebraic set is homeomorphic to an algebraic set.
- (b) Given algebraic sets $L \subset V$, then $V - L$ is homeomorphic to an algebraic set.
- (c) Given algebraic sets $L \subset V$ with V compact then V/L is homeomorphic to an algebraic set.

Proof:

(a) Let $Z \subset \mathbf{R}^n$ be an algebraic set and assume that $Z \neq \mathbf{R}^n$ and $0 \notin Z$ (otherwise translate Z). Let $Z = f^{-1}(0)$ for some polynomial $f(x)$; then define $F(x) = |x|^{2d} f\left(\frac{x}{|x|^2}\right)$, where d is the degree of $f(x)$. Clearly $F(x)$ is a polynomial and $F^{-1}(0)$ is the one point compactification of Z , since $x \mapsto \frac{x}{|x|^2}$ is the inversion through the unit sphere.

(b) Let $V = f^{-1}(0)$, $L = g^{-1}(0)$ for some polynomials $f, g: \mathbf{R}^n \rightarrow \mathbf{R}$. Define $G(x, t) = |f(x)|^2 + |tg(x) - 1|^2$, then $G^{-1}(0) \approx V/L$.

(c) By applying (a) we get the one point compactification of $G^{-1}(0)$ to be an algebraic set; if V is compact this set is homeomorphic to V/L . \square

This proposition implies that a set is homeomorphic to an algebraic set if and only if the one point compactification is homeomorphic to an algebraic set. Hence any noncompact algebraic set has a collar at infinity, since every algebraic set is locally cone-like [M]. Also we get that the reduced suspension $\Sigma^n X = X \times S^n / X \vee S^n$ of any algebraic set X is homeomorphic to an algebraic set.

There is a fancier version of the blowing down operation (c) (Proposition 3.3). First we need to discuss projectively closed algebraic sets. Let $p: \mathbf{R}^n \rightarrow \mathbf{R}$ be a polynomial. Another interpretation of this concept is the following: Let $\lambda: \mathbf{R}^n \rightarrow \mathbf{R}^d$. We call $p(x)$ an *overt polynomial* if $p_d^{-1}(0)$ is either the empty set or $\{0\}$. We call an algebraic set $V = p^{-1}(0)$ a *projectively closed algebraic set* if $p(x)$ is an overt polynomial. Another interpretation of this concept is the following: Let $\lambda: \mathbf{R}^n \rightarrow \mathbf{RP}^n$ be the inclusion $\lambda(x_1, \dots, x_n) = [1; x_1; \dots; x_n]$ then $V = p^{-1}(0)$ is projectively closed if and only if λ is a projective algebraic subset of \mathbf{RP}^n in other words $\lambda(V)$ is Zariski closed in \mathbf{RP}^n (see also [AK₂]). Real algebraic sets along with maps can easily be made projectively closed by the following.

PROPOSITION 3.2. *Let $f: Z \rightarrow W$ be an entire rational function between algebraic sets with Z nonsingular and compact. Then there is a projectively closed algebraic set $V \subset W \times \mathbf{R}^n$ a birational diffeomorphism g which makes the following commute*

$$\begin{array}{ccc} V & \hookrightarrow & W \times \mathbf{R}^n \\ g \uparrow \approx & & \downarrow \pi \\ Z & \xrightarrow{f} & W \end{array}$$

where π is the projection, n is some integer.

Proof: By taking the graph of f we can assume that $Z \subset W \times \mathbf{R}^m \subset \mathbf{R}^r$ for some r , and f is induced by projection. Also identify $\mathbf{R}^r \subset \mathbf{RP}^r$ via λ . Then let \bar{Z} be the Zariski closure of Z in \mathbf{RP}^r . We claim $\dim(\bar{Z} - Z) < \dim(Z)$. This is because if U is an irreducible component of \bar{Z} then $U \cap Z \neq \emptyset$, and therefore $U - Z = U \cap \mathbf{RP}^{r-1}$ is a proper algebraic subset of U where $\mathbf{RP}^{r-1} = \{[0; x_1; \dots; x_r] \in \mathbf{RP}^r\}$. Since U is irreducible $\dim(U - Z) < \dim(U)$, also $\dim(U) = \dim(Z)$. Therefore $\dim(\bar{Z} - Z) < \dim(Z)$. So $\bar{Z} - Z = \text{Sing}(\bar{Z})$. By resolution of singularities [H] (Theorem 1.1) there is a nonsingular algebraic set $V \subset \mathbf{RP}^r \times \prod_i \mathbf{RP}^{a_i}$ such that the projection induces birational diffeomorphism between V and Z . In particular $V \subset \mathbf{R}^r \times \prod_i \mathbf{RP}^{a_i}$.




$$\mathbf{RP}^r \times \prod_i \mathbf{RP}^{a_i} \hookrightarrow \mathbf{R}^{(r+1)^2 + \sum (a_i+1)^2}$$

is a projectively closed algebraic set. Hence V is projectively closed (check details). \square

Now assume that $L \subset W \subset \mathbf{R}^m$ be real algebraic sets, and $V \subset W \times \mathbf{R}^n$ be a projectively closed algebraic set. Let $q: \mathbf{R}^m \rightarrow \mathbf{R}$ be a polynomial with $q^{-1}(0) = L$. Define

$$D_q: W \times \mathbf{R}^n \rightarrow W \times \mathbf{R}^n$$

by $D_q(x, y) = (x, yq(x))$. D_q is a diffeomorphism on $(W - L) \times \mathbf{R}^n$ and $D_q(L \times \mathbf{R}^n) = L \times 0$. Therefore $D_q(V)$ is the quotient space of V by the equivalence relation $(x, y) \sim (x, 0)$ if $x \in L$. We call the operation $V \rightarrow D_q(V) \cup L$ (L is identified by $L \times 0$) *blowing down* V over L .



PROPOSITION 3.3. Given L, W, V as above, then $D_q(V) \cup L$ is an algebraic subset of $W \times \mathbf{R}^n$.

Proof: Let $p: \mathbf{R}^m \times \mathbf{R}^n \rightarrow \mathbf{R}$ be an overt polynomial of degree e with $V = p^{-1}(0)$ and let q be as above. Define a polynomial $r: \mathbf{R}^m \times \mathbf{R}^n \rightarrow \mathbf{R}$ by

$$r(x, y) = q(x)^e p\left(x, \frac{y}{q(x)}\right)$$

We claim $r^{-1}(0) = D_q(V) \cup L$. It is easy to see that

$$r^{-1}(0) \cap (W - L) \times \mathbf{R}^n = D_q(V) \cap (W - L) \times \mathbf{R}^n,$$

so it suffices to show that $r^{-1}(0) \cap (L \times \mathbf{R}^n) = L \times 0$. We decompose $p(x, y) = p_e(x, y) + \alpha(x, y)$ where $p_e(x, y)$ is homogeneous of degree e and $\alpha(x, y)$ is a polynomial of degree less than e . Hence if $(x, y) \in r^{-1}(0) \cap (L \times \mathbf{R}^n)$ then $r(x, y) = 0$ and $q(x) = 0$, which implies $r(x, y) = p_e(0, y) = 0$. Then $y = 0$ since p is overt, so $(x, y) \in L \times 0$. Conversely if $(x, y) \in L \times 0$ then $y = 0$ and $q(x) = 0$. Hence $r(x, y) = p_e(0, 0) = 0$, i.e. $(x, y) \in r^{-1}(0) \cap (L \times \mathbf{R}^n)$. \square

There is a more useful version of Proposition 3.3 which says that after modifying D_q we can get $D_q(V) \cup L$ to be a projectively closed algebraic set (Proposition 3.1 of [AK₆]). This allows us to iterate this blowing down process.

§4. ISOLATED SINGULARITIES

The topology of real algebraic sets with isolated singularities is completely understood by the following Theorem.

THEOREM 4.1 ([AK₂]). X is homeomorphic to an algebraic set with isolated singularities if and only if X is obtained by taking a smooth compact manifold W with boundary $\partial W = \bigcup_{i=1}^r M_i$, where each M_i bounds, then crushing some M_i 's to points and deleting the remaining M_i 's.

