

2. CR STRUCTURES AND THE HEISENBERG GROUP

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **31 (1985)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **02.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Let H be the kernel of Δ in $C_c^\infty(E_2)$. A Hodge decomposition for $C_c^\infty(E_2)$ is

$$C_c^\infty(E_2) = D_1(C_c^\infty(E_1)) \oplus D_2^*(C_c^\infty(E_3)) \oplus H.$$

Hodge studied the de Rham complex on a compact Riemannian manifold. The Riemannian metric induced the metrics on the bundles $\Lambda^p T^*(M)$ as well as the volume element.

In the next section we discuss abstract CR manifolds and look at the Heisenberg group in detail. We write down the $\bar{\partial}_b$ and $\square_b$ operators explicitly and give Folland and Stein's inverse to $\square_b$. In section 3 we introduce the stratified Lie groups and the associated homogeneous structures. We present the continuity theorems of Folland and Rothschild and Stein for convolution operators. In section 4 we prove the decomposition theorem in the general setting of stratified groups.

These results are an extension of the author's dissertation [4]. We wish to express our deep gratitude to M. Kuranishi. We would also like to thank D. Tartakoff for his help and encouragement.

2. CR STRUCTURES AND THE HEISENBERG GROUP

Let M be a C^∞ manifold of dimension $2n + 1$. The complexified tangent bundle of M , $CT(M)$, is the bundle whose fiber is $\mathbb{C} \otimes_{\mathbb{R}} T_m(M)$ where $T_m(M)$ is the tangent space at $m \in M$. When there is no confusion we will drop reference to M in the notation for $T(M)$, $CT(M)$, etc. So, $T(M) = T$ and $CT(M) = CT$, for example.

A CR structure on M is a subbundle $T_{1,0} \subset CT$ such that (i) $T_{1,0} \cap \overline{T_{1,0}} = \{0\}$, (ii) $\text{codim}(T_{1,0} \otimes \overline{T_{1,0}}) = 1$, (iii) if X and Y are smooth sections of $T_{1,0}$ then $[X, Y] = XY - YX$ is a section $T_{1,0}$. We set $T_{0,1} = \overline{T_{1,0}}$. If M has a CR structure it is called a CR manifold.

An example of a CR manifold is a real hypersurface M in a complex manifold M' , $M \subset M'$. Define $T_{1,0}(M) = CT(M) \cap T_{1,0}(M')$ where $T_{1,0}(M')$ is the holomorphic tangent bundle of M' .

If M is a CR manifold set $T^{1,0}$ (resp., $T^{0,1}$) to be the dual space to $T_{1,0}$ (resp., $T_{0,1}$). Let $\Lambda^{p,q}$ be the space of C^∞ sections of $\Lambda^p T^{1,0} \otimes \Lambda^q T^{0,1}$. Define the operator $\bar{\partial}_b: \Lambda^{p,q} \rightarrow \Lambda^{p,q+1}$ as follows: Let $\phi \in \Lambda^{p,q}$ and let $X_1, \dots, X_p$ (resp., $Y_1, \dots, Y_{q+1}$) be sections of $T_{1,0}$ (resp., $T_{0,1}$). Then

$$\begin{aligned}
& \langle \bar{\partial}_b \phi; (X_1 \wedge \dots \wedge X_p) \otimes (Y_1 \wedge \dots \wedge Y_{q+1}) \rangle \\
&= (q+1)^{-1} \sum_{j=1}^{q+1} (-1)^{j+1} Y_j \langle \phi; (X_1 \wedge \dots \wedge X_p) \otimes (Y_1 \wedge \dots \wedge \hat{Y}_j \wedge \dots \wedge Y_{q+1}) \rangle \\
&+ (q+1)^{-1} \sum_{i < j} (-1)^{i+j} \langle \phi; (X_1 \wedge \dots \wedge X_p) \otimes ([Y_i, Y_j] \wedge Y_1 \wedge \dots \wedge \hat{Y}_i \wedge \dots \wedge \hat{Y}_j \wedge \dots \wedge Y_{q+1}) \rangle.
\end{aligned}$$

The $\hat{}$ symbol over a section means as usual that it is deleted from the expression. One can show that

- i) $\bar{\partial}_b^2 = 0$,
- ii) $\bar{\partial}_b(\phi \wedge \psi) = (\bar{\partial}_b \phi) \wedge \psi + (-1)^p \phi \wedge \bar{\partial}_b \psi$ for $\phi \in \Lambda^{0,p}$,
- iii) $\langle \bar{\partial}_b f, Y \rangle = Yf$ for $f \in \Lambda^{0,0}$ and Y a section of $T_{0,1}$.

See [6] for details.

The Heisenberg group, H , is a Lie group with a natural CR structure. The manifold is $\mathbf{C}^n \times \mathbf{R}$. Let $(z, t), (z', t') \in \mathbf{C}^n \times \mathbf{R} = H$. The group law is defined by

$$(z, t) \cdot (z', t') = (z + z', t + t' + 2 \operatorname{Im}(z \cdot z'))$$

where $z \cdot z' = \sum_{j=1}^n z_j \bar{z}'_j$. The identity element is $(0, 0)$ and $(z, t)^{-1} = (-z, -t)$.

Sometimes we will set $u = (z, t)$.

For $j = 1, \dots, n$ if we set $z_j = x_j + iy_j$, the mapping

$$(z, t) \rightarrow (x_1, \dots, x_n, y_1, \dots, y_n, t)$$

defines a C^∞ coordinate system on H . The left invariant vector fields (i.e., the elements of the Lie algebra) are $\mathbf{R}$ -linear combinations of

$$(2) \quad X_j = \frac{\partial}{\partial x_j} + 2y_j \frac{\partial}{\partial t}, \quad Y_j = \frac{\partial}{\partial y_j} - 2x_j \frac{\partial}{\partial t}, \quad T = \frac{\partial}{\partial t}.$$

They satisfy the following commutation relations:

$$(3) \quad [X_j, T] = [Y_j, T] = [X_j, X_k] = [Y_j, Y_k] = 0,$$

$$(4) \quad [X_j, Y_k] = -4\delta_{jk}T.$$

Let CT be the complexified tangent bundle of H . Define

$$(5) \quad Z_j = \frac{1}{2}(X_j - iY_j) = \frac{\partial}{\partial z_j} + i\bar{z}_j \frac{\partial}{\partial t}.$$

Then Z_j , $\bar{Z}_j$, and T form a basis (over $\mathbf{R}$) of the space of left invariant complex tangent vector fields. In particular, they form a global frame of CT . From (3), (4) and (5) we easily see that

$$(6) \quad [Z_j, Z_k] = [\bar{Z}_j, \bar{Z}_k] = [Z_j, T] = [\bar{Z}_j, T] = 0 \\ [Z_j, \bar{Z}_k] = -2i\delta_{jk}T.$$

Let $T_{1,0}$ (resp., $T_{0,1}$) be the subbundle of CT spanned by the Z_j 's (resp., $\bar{Z}_j$'s). Then

$$\bar{T}_{1,0} = T_{0,1} \\ T_{1,0} \cap T_{0,1} = \{0\} \\ \text{codim}(T_{1,0} \otimes T_{0,1}) = 1.$$

Also, if V_1, V_2 are sections of $T^{1,0}$ we can write $V_i = \sum_{j=1}^n f_{ij}Z_j, i = 1, 2$ where the f_{ij} are C^∞ functions on H . Then by (6),

$$[V_1, V_2] = \sum_{k=1}^n \left(\sum_{j=1}^n (f_{1j}Z_j f_{2k} - f_{2j}Z_j f_{1k}) \right) Z_k.$$

So, the splitting of CT defines a CR structure on H .

Impose the left invariant Hermitian metric on CT which makes the Z 's, $\bar{Z}$'s and T an orthonormal frame. Let ω^j and τ be dual to Z_j and T , respectively. Then $\omega^j, \bar{\omega}^j$ and τ form an orthonormal frame for CT^* . The volume element on H is

$$(7) \quad du = 2^n dx_1 \wedge \dots \wedge dx_n \wedge dy_1 \wedge \dots \wedge dy_n \wedge dt.$$

Since H is nilpotent and, hence, unimodular, the volume element is both left and right invariant. One can also verify this directly.

Let $J = (j_1, \dots, j_q)$ be a multi-index with $1 \leq j_i \leq n, i = 1, \dots, q$. Define $|J| = q$ and $\bar{\omega}^J = \bar{\omega}^{j_1} \wedge \dots \wedge \bar{\omega}^{j_q}$. If $\phi \in \Lambda^{0,q}(H)$ we may write $\phi = \sum_{|J|=q} \phi_J \bar{\omega}^J$ where ϕ_J is a C^∞ function from H to $\mathbf{C}$. Let

$$\bar{\omega}^j \lrcorner \bar{\omega}^J = (-1)^k \bar{\omega}^{j_1} \wedge \dots \wedge \widehat{\bar{\omega}^{j_k}} \wedge \dots \wedge \bar{\omega}^{j_q} \quad \text{if } j = j_k \quad \text{and} \quad \bar{\omega}^j \lrcorner \bar{\omega}^J = 0$$

otherwise. Folland and Stein prove that for $\phi \in \Lambda^{0,q}$

$$i) \quad \bar{\partial}_b \phi = \sum_{|J|=q} \sum_{j=1}^n \bar{Z}_j \phi_J \bar{\omega}^j \wedge \bar{\omega}^J,$$

$$(8) \quad \text{ii) } \bar{\partial}_b^* \phi = - \sum_{|J|=q} \sum_{j=1}^n Z_j \phi_J \bar{\omega}^j \lrcorner \bar{\omega}^J,$$

$$\text{iii) } \square_b \phi = \sum_{|J|=q} \left(-\frac{1}{2} \sum_{j=1}^n (Z_j \bar{Z}_j + \bar{Z}_j Z_j) + i(n-2q)T \right) \phi_J \bar{\omega}^J.$$

Define the function

$$\Phi_\alpha(z, t) = (|z|^2 - it)^{-\frac{n+\alpha}{2}} (|z|^2 + it)^{-\frac{n-\alpha}{2}}.$$

Let $\phi \in \Lambda_c^{0,q}$, $q \neq 0, n$. For an appropriate constant, c_q , define

$$(9) \quad K_q \phi(v) = c_q \sum_{|J|=q} \left(\int_H \phi_J(u) \Phi_{n-2q}(u^{-1}v) du \right) \bar{\omega}^J.$$

Folland and Stein prove that for the appropriate c_q

THEOREM 1. Let $\phi \in \Lambda_c^{0,q}$, $q \neq 0, n$. Then $\square_b K_q \phi = K_q \square_b \phi = \phi$.

In [4] we prove a stronger version of the following Hodge decomposition theorem.

THEOREM 2. Let $\phi \in \Lambda_c^{0,q}$, $q \neq 0, n$. Then

- i) $H\phi = 0$ where H is the orthogonal projection onto the kernel of $\square_b$.
- ii) $\phi = \bar{\partial}_b \bar{\partial}_b^* K_q \phi + \bar{\partial}_b^* \bar{\partial}_b K_q \phi$.

We also prove

THEOREM 3. If $\phi \in \Lambda_c^{0,q}$, $q \neq 0, n$ and if $\bar{\partial}_b \phi = 0$ then $\psi = \bar{\partial}_b^* K_q \phi$ satisfies $\bar{\partial}_b \psi = \phi$.

These two theorems are special cases of theorems 6 and 7 proven in section 4.

3. DIFFERENTIAL COMPLEXES ON STRATIFIED GROUPS

We study a class of nilpotent Lie groups which we describe in terms of their Lie algebras. A graded Lie algebra, $\mathfrak{n}$, is a finite dimensional nilpotent algebra which has a direct sum decomposition, $\mathfrak{n} = \bigoplus_{i=1}^r \mathfrak{n}_i$ where the $\mathfrak{n}_i$ satisfy

- i) $[\mathfrak{n}_i, \mathfrak{n}_j] \subseteq \mathfrak{n}_{i+j}$ if $i+j \leq r$,
- ii) $[\mathfrak{n}_i, \mathfrak{n}_j] = 0$ if $i+j > r$.