

4. The Hodge decomposition

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **31 (1985)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **02.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

THEOREM 4. (i) Let K be a convolution operator of type r for $r > 0$. Then K extends from $C_c^\infty(\mathbf{F}^m)$ to a bounded operator from $L^p(\mathbf{F}^m)$ to $L^q(\mathbf{F}^{m'})$ where $1 < p < Q/r$ and $q^{-1} = p^{-1} - r/Q$. (ii) Let K be a convolution operator of type 0. Then K extends from $C_c^\infty(\mathbf{F}^m)$ to a bounded operator from $S_k^p(\mathbf{F}^m)$ to $S_k^p(\mathbf{F}^{m'})$.

Finally, we mention the interaction between the homogeneous convolution operators and the left-invariant differential operators. Let $D: C^\infty(\mathbf{F}^{m'}) \rightarrow C^\infty(\mathbf{F}^{m''})$ be a left-invariant homogeneous differential operator of degree 1 and let K be a homogeneous convolution operator of type r , with $r \geq 1$. Then DK is a homogeneous convolution operator of type $r - 1$. Moreover, if $r > 1$ the kernel of DK is given by $Dk(x)$.

4. THE HODGE DECOMPOSITION

Consider the complex (1) where $E_i = \mathbf{R}^n \times \mathbf{F}^{m_i}$. Assume that each of the D_i is a first order, left-invariant operator, homogeneous of degree 1. So each entry of D_i is of the form $\sum_{j=1}^d a_j X_{1,j}$ where a_j is constant. Construct the Laplacian, Δ , with respect to the euclidian inner products on $\mathbf{F}^{m_i}$, $i = 1, 2, 3$. Assume there exists a homogeneous convolution operator of type 2, K , which inverts Δ . If $f \in C_c^\infty(\mathbf{F}^{m_2})$ then $f(x) = \Delta K f(x) = K \Delta f(x)$.

THEOREM 5. Let $f \in S_2^2(\mathbf{F}^{m_2})$. As distributions, $\Delta f = 0$ if and only if $f = 0$.

Proof. Obviously, if $f = 0$ then $\Delta f = 0$.

Assume $\Delta f = 0$. Let $\{f_j\}$ be a sequence in $C_c^\infty(\mathbf{F}^{m_2})$ such that $f_j \rightarrow f$ in $S_2^2(\mathbf{F}^{m_2})$. Then $f_j \rightarrow f$ in the sense of distributions. Moreover, $\Delta f_j \rightarrow \Delta f = 0$ in $L^2(\mathbf{F}^{m_2})$. Let $g \in C_c^\infty(\mathbf{F}^{m_2})$. Then

$$\langle f, g \rangle = \lim_{j \rightarrow \infty} \langle f_j, g \rangle = \lim_{j \rightarrow \infty} \langle f_j, \Delta K g \rangle = \lim_{j \rightarrow \infty} \langle \Delta f_j, K g \rangle.$$

Because $g \in C_c^\infty(\mathbf{F}^{m_2})$ it is in L^p where $p = 2Q/(Q+4)$. Therefore, by Theorem 4(i), $Kg \in L^q$ where

$$q^{-1} = (Q+4)/2Q - 2/Q = 1/2, \text{ i.e., } Kg \in L^2(\mathbf{F}^{m_2}).$$

For $Q \geq 5$, $1 < p < q < \infty$. So

$$|\langle f, g \rangle| = \lim_{j \rightarrow \infty} |\langle \Delta f_j, K g \rangle| \leq \lim_{j \rightarrow \infty} \|\Delta f_j\|_{L^2(\mathbf{F}^{m_2})} \|Kg\|_{L^2(\mathbf{F}^{m_2})} = 0.$$

So, as a distribution, $f = 0$. This proves the theorem.

We have shown that the only harmonic element in $S_2^2(\mathbf{F}^{m_2})$ is the zero element. Let $f \in S_2^2(\mathbf{F}^{m_2})$ and let $f_j \rightarrow f$ in $S_2^2(\mathbf{F}^{m_2})$ with $f_j \in C_c^\infty(\mathbf{F}^{m_2})$. Then

$$f = \lim_{j \rightarrow \infty} \Delta K f_j = \lim_{j \rightarrow \infty} D_1 D_1^* K f_j + \lim_{j \rightarrow \infty} D_2^* D_2 K f_j = D_1 D_1^* K f + D_2^* D_2 K f.$$

To complete the Hodge decomposition we must prove that

$$D_1 D_1^* K f \perp D_2^* D_2 K f.$$

We need the following notation. Let $D(R) = \{x \in N : |x| < R\}$ and $S(R) = \{x \in N : |x| = R\}$. Endow each set with the left-invariant metric induced by N . The metric gives rise to the corresponding volume elements which, in the case of $D(R)$, is the restriction of dx . Let $d\mu_R$ denote the volume element on $S(R)$. For $f, g \in C_c^\infty(D(R), \mathbf{F}^{m_i})$ define

$$(f, g)_{D(R), i} = \int_{D(R)} (f(x), g(x))_{i, x} dx$$

where $(\cdot, \cdot)_{i, x}$ is the metric on $\mathbf{F}^{m_i}$. Similarly for $f, g \in C^\infty(S(R), \mathbf{F}^{m_i})$ define

$$(f, g)_{S(R), i} = \int_{S(R)} (f(x), g(x))_{i, x} d\mu_R(x).$$

By restriction, any element $f \in C^\infty(N, \mathbf{F}^{m_i})$ gives rise to an element of $C^\infty(S(R), \mathbf{F}^{m_i})$ or $C^\infty(D(R), \mathbf{F}^{m_i})$. In our notation, we will not distinguish f from its restrictions.

We will be integrating by parts on $D(R)$ which will involve a boundary integral on $S(R)$. To that end we define the symbol of our differential operators. Define $\xi(x) = |x| - R$ and let $g \in C^1(N, \mathbf{F}^{m_i})$. Let $x \in S(R)$. Then $\xi(x) = 0$. The symbol of D_i at $x \in S(R)$ acting on $d\xi$ and on g is given by

$$\sigma(D_i, d\xi)g(x) = D_i(\xi g)(x).$$

The integration by parts formula is

$$(13) \quad (D_i g, f)_{D(R), i+1} = (g, D_i^* f)_{D(R), i} + (\sigma(D_i, d\xi)g, f)_{S(R), i}.$$

THEOREM 6. Assume $f \in L^2(\mathbf{F}^{m_2}) \cap L^q(\mathbf{F}^{m_2})$ where $q = Q/Q + 2$. Then $f = D_1 D_1^* K f + D_2^* D_2 K f$ and $D_1 D_1^* K f \perp D_2^* D_2 K f$.

Proof. We have already seen that $f = D_1 D_1^* K f + D_2^* D_2 K f$. To prove the orthogonality we restrict our attention to $D(R)$ for R large.

For brevity let $h = (D_1 D_1^* Kf, D_2^* D_2 Kf)_2$. Also, let

$$h(R) = (D_1 D_1^* Kf, D_2^* D_2 Kf)_{D(R), 2}.$$

Then $\lim_{R \rightarrow \infty} h(R) = h$. Note that $D_1 D_1^* K$ and $D_2^* D_2 K$ are type 0 operators.

Since $f \in L^2(\mathbb{F}^{m_2})$ by theorem 4 (ii) we know that h is defined. Furthermore $h(R)$ is bounded for all R by $\|D_1 D_1^* Kf\|_{L^2(\mathbb{F}^{m_2})} \|D_2^* D_2 Kf\|_{L^2(\mathbb{F}^{m_2})}$.

We can compute $h(R)$ as follows:

$$\begin{aligned} h(R) &= (D_1 D_1^* Kf, D_2^* D_2(Kf))_{D(R), 2} = (D_2 D_1 D_1^* Kf, D_2 Kf)_{D(R), 3} \\ &\quad + (D_1 D_1^* Kf, \sigma(D_2^*, d(|x|)) D_2 Kf)_{S(R), 2} \end{aligned}$$

by (13). We now prove a sequence of lemmas.

LEMMA 1. $h(R)$ is continuous.

Proof. This follows from Lebesgue's dominated convergence theorem.

LEMMA 2. Let $R \geq 1$, $x \in N$ and $|x| = R$. Then

$$|\sigma(D_2^*, d|x|)g(x)| \leq C |g(x)|$$

where C is a constant independent of g .

Proof. Recall that $X_1, \dots, X_d$ is our orthonormal basis for n_1 where $d = \dim(n_1)$. The entries of D_2^* are linear combinations of the X_i , $i = 1, \dots, d$ with coefficients in $\mathbb{F}$. Let D_{ij} be the i, j entry, $1 \leq i \leq m_2$, $1 \leq j \leq m_3$. Then $D_{ij} = \sum_{k=1}^d C_{ij}^k X_k$. Thus, for $x \in S(R)$

$$\begin{aligned} |\sigma(D_2^*, d|x|)g(x)| &\leq C \sum_{i=1}^{m_2} \left| \sum_{j=1}^{m_3} D_{ij}(|x| - R)g_j(x) \right| \\ &\leq C \sum_{i,j,k} |C_{ij}^k X_k(|x| - R)g_j(x)| \\ &\leq C \sum_{jk} |X_k|x|g_j(x)| \quad (\text{since } |x| = R) \\ &\leq C(\max_k |X_k|x|) |g_j(x)|. \end{aligned}$$

We must show $X_k|x|$ is bounded. But $|x|$ is C^∞ away from the origin and homogeneous of degree 1. So $X_k|x|$ is homogeneous of degree 0. Thus, it is determined by its values on $\{|x| = 1\}$. It is C^∞ on this set and, therefore, bounded. This proves the lemma.

Since $h(R) \rightarrow h$ as $R \rightarrow \infty$ we have

LEMMA 3. For $\varepsilon > 0$, $\lim_{r \rightarrow \infty} \frac{1}{2\varepsilon} \int_{r-\varepsilon}^{r+\varepsilon} h(R) dR = h$.

We continue with the proof of our theorem. By the preceding lemma it suffices for us to prove that $\lim_{r \rightarrow \infty} \frac{1}{2\varepsilon} \int_{r-\varepsilon}^{r+\varepsilon} h(R) dR = 0$. But,

$$(14) \quad \begin{aligned} & \frac{1}{2\varepsilon} \int_{r-\varepsilon}^{r+\varepsilon} h(R) dR \\ &= \frac{1}{2\varepsilon} \int_{r-\varepsilon \leq |x| \leq r+\varepsilon} (D_1 D_1^* Kf, \sigma(D_2^*, d|x|) D_2 Kf)_x \|d|x|\| dx \end{aligned}$$

because $d\mu_R dR = \|d|x|\| dx$. We claim that $\|d|x|\|$ is bounded. Let ω^{ij} be dual to X_{ij} where X_{ij} is our orthonormal basis. Then $d|x| = \sum_{i,j} (X_{ij}|x|) \omega^{ij}$. Since $|x|$ is homogeneous of degree 1 and X_{ij} is homogeneous of degree i we have $X_{ij}|x|$ is homogeneous of degree $1-i$. Hence, for $|x| \geq 1$ each $X_{ij}|x|$ is bounded. So $\|d|x|\|$ is bounded.

By assumption, $f \in L^q(\mathbb{F}^{m_2})$, $q = Q/Q + 2$ and we know that $D_2 K$ is type 1. By Theorem 4(i) we know that $D_2 Kf \in L^2(\mathbb{F}^{m_2})$. Thus, by Lemma 2 $|\chi_r \sigma(D_2^*, d|x|) D_2 Kf| \leq C |\chi_r D_2 Kf|$ where χ_r is the characteristic function of $\{r - \varepsilon \leq |x| \leq r + \varepsilon\}$. We conclude that $\chi_r \sigma(D_2^*, d|x|) D_2 Kf \in L^2(\mathbb{F}^{m_2})$. By the Schwarz inequality and the fact that $\|d|x|\|$ is bounded, we get, from (14)

$$\frac{1}{2\varepsilon} \int_{r-\varepsilon}^{r+\varepsilon} h(R) dR \leq \frac{c}{2\varepsilon} \|\chi_r D_1 D_1^* Kf\|_{L^2(\mathbb{F}^{m_2})} \|\chi_r D_2 Kf\|_{L^2(\mathbb{F}^{m_3})}.$$

As $r \rightarrow \infty$, both $\|\chi_r D_1 D_1^* Kf\|_{L^2(\mathbb{F}^{m_2})}$ and $\|\chi_r D_2 Kf\|_{L^2(\mathbb{F}^{m_3})}$ tend to 0. So

$$h = \lim_{r \rightarrow \infty} \frac{1}{2\varepsilon} \int_{r-\varepsilon}^{r+\varepsilon} h(R) dR = 0. \text{ This proves the theorem.}$$

This theorem together with Theorem 5 proves the Hodge decomposition. A similar argument gives the solution to the problem of finding g such that $D_1 g = f$ for a given f .

THEOREM 7. Let $f \in L^2(\mathbb{F}^{m_2}) \cap L^q(\mathbb{F}^{m_2})$ with $q = Q/Q + 2$. Suppose $D_2 f = 0$. Then there exists $g \in L^2(\mathbb{F}^{m_1})$ such that $D_1 g = f$.

Proof. We have $f = D_1 D_1^* Kf + D_2^* D_2 Kf$. It suffices to prove $(f, D_2^* D_2 Kf) = 0$ because this implies $D_2^* D_2 Kf = 0$ since

$$D_1 D_1^* Kf \perp D_2^* D_2 Kf.$$

We may set $g = D_1^* Kf$. Using the same notation as in the preceding theorem we have

$$\begin{aligned} (f, D_2^* D_2 Kf)_2 &= \lim_{R \rightarrow \infty} (f, D_2^* D_2 Kf)_{D(R), 2} \\ &= \lim_{R \rightarrow \infty} ((D_2 f, D_2 Kf)_{D(R), 3} + (f, \sigma(D_2^*, d|x|) D_2 Kf)_{S(R), 2}) \\ &= \lim_{R \rightarrow \infty} (f, \sigma(D_2^*, d|x|) D_2 Kf)_{S(R), 2} \quad (\text{since } D_2 f = 0). \end{aligned}$$

The same argument as in Theorem 6 proves that the limit is zero. This proves the theorem.

REFERENCES

- [1] ATIYAH, M. F. and R. BOTT. A Lefschetz fixed point formula for elliptic complexes, I, II. *Ann. of Math.* 86 (1967), 347-407; 88 (1968), 451-491.
- [2] DADOK, J. and R. HARVEY. The fundamental solution for the Kohn-Laplacian $\bar{\partial}_b$ on the sphere in $\mathbb{C}^n$. *Math. Ann.* 244 (1979), 89-104.
- [3] DE RHAM, G. *Variétés Différentiables*. Paris: Hermann & Cie, 1955.
- [4] DUDDY, J. A. *The boundary Laplacian on the Heisenberg group and the sphere*. Ph. D. dissertation, Columbia University, 1978.
- [5] DOLBEAULT, P. Sur la cohomologie des variétés analytiques complexes. *C. R. Acad. Sci. Paris* 236 (1953), 175-177.
- [6] FOLLAND, G. B. and J. J. KOHN. *The Neumann Problem for the Cauchy-Riemann Complex*. Princeton, N.J.: Princeton University Press, 1972.
- [7] FOLLAND, G. B. and E. M. STEIN. Parametrics and estimates for the $\bar{\partial}_b$ complex on strongly pseudoconvex boundaries. *Bull. Amer. Math. Soc.* 80 (1974), 253-258.
- [8] FOLLAND, G. B. Estimates for the $\bar{\partial}_b$ complex and analysis on the Heisenberg group. *Comm. Pure Appl. Math.* 27 (1974), 429-522.
- [9] ——— Subelliptic estimates and function spaces on nilpotent Lie groups. *Ark. Math.* 13 (1975), 161-207.
- [10] GREENFIELD, S. J. Cauchy-Riemann equations on several variables. *Ann. Scuola Norm. Pisa* 22 (1968), 275-314.
- [11] HARVEY, R. and J. POLKING. Fundamental solutions in complex analysis I and II. *Duke Math. J.* 46 (1979), 253-340.
- [12] HODGE, W. V. D. *The Theory and Applications of Harmonic Integrals*. New York: Cambridge University Press, 1941 (2d ed., 1952).
- [13] KODAIRA, K. Harmonic fields in Riemannian manifolds (generalized potential theory). *Ann. of Math.* 50 (1949), 587-665.
- [14] KOHN, J. J. and H. ROSSI. On the extension of holomorphic functions from the boundary of a complex manifold. *Ann. of Math.* 81 (1965), 451-472.
- [15] LEWY, H. An example of a smooth linear partial differential equation without solution. *Ann. of Math.* 66 (1957), 155-158.