

THE VARIETY OF MODULI OF FOLIATIONS ON A COMPLEX SPACE

Autor(en): **Reiffen, Hans-Jörg**

Objektyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **33 (1987)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **01.06.2024**

Persistenter Link: <https://doi.org/10.5169/seals-87892>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

THE VARIETY OF MODULI OF FOLIATIONS ON A COMPLEX SPACE

by Hans-Jörg REIFFEN

§ 0. Let X be a complex space. By X_s we denote the singular locus of X . Let $\mathcal{F}$ be a coherent analytic sheaf on X . By $S(\mathcal{F})$ we denote the singular locus of $\mathcal{F}$, i.e. the set of points, where $\mathcal{F}$ is not free. $S(\mathcal{F})$ is an analytic subset of X . If X is reduced then $S(\mathcal{F})$ is thin. For a coherent subsheaf $\mathcal{F}'$ of $\mathcal{F}$ we set $S(\mathcal{F} : \mathcal{F}') := S(\mathcal{F}) \cup S(\mathcal{F}/\mathcal{F}')$.

In the following let X be a reduced complex space. If X' is an irreducible component of X , then we denote by $r(\mathcal{F}, X')$ the rank of $\mathcal{F}$ on $X' \setminus (X_s \cup S(\mathcal{F}))$. Ω resp. Θ is the sheaf of holomorphic 1-forms resp. of vector fields on X . We have $S(\Omega) = X_s$.

0.1 DEFINITION. Let Ω' be a coherent subsheaf of Ω and X' an irreducible component of X . Ω' is a X' -foliation, iff there is a thin analytic subset A of X' , $X' \cap X_s \subset A$, such that $\Omega' \mid X' \setminus A$ defines a regular foliation. Ω' is a foliation, iff Ω' is a X' -foliation for every irreducible component X' of X .

In a joint work with G. Bohnhorst (comp. [B/R]) I have developed a general theory of foliations. The coherent foliations as they are introduced in [B/R] correspond to those foliations used in this paper, which are full subsheaves of Ω . We call $\text{codim}_{X'} \Omega' := r(\Omega'; X')$ the codimension of the X' -foliation.

The following homomorphism of sheaves (of groups) is important

$$\delta^p : \Omega^{p+1} \rightarrow \wedge^{p+2} \Omega, \quad \delta^p(\omega^0; \omega^1, \dots, \omega^p) := d\omega^0 \wedge \omega^1 \wedge \dots \wedge \omega^p.$$

0.2 REMARK. Let Ω' be a coherent subsheaf of Ω . Then the following are equivalent:

- (1) Ω' is a X' -foliation of codimension p ;
- (2) $p = r(\Omega'; X')$, $\delta^p \mid (\Omega' \mid X' \setminus X_s)^{p+1} = 0$;
- (3) $p = r(\Omega'; X')$, there is a point $x \in X' \setminus X_s$, such that $\delta^p((\Omega'_x)^{p+1}) = 0$.

For the following compilation of definitions and results the paper by Douady [Dou] is a good reference.

In the following let Y, Z be further complex spaces, that are not necessarily reduced. If $\mathcal{E}$ is a coherent analytic sheaf on Y and $f: Z \rightarrow Y$ a holomorphic mapping then we denote by $f^*\mathcal{E}$ the analytic inverse image of $\mathcal{E}$ on Z .

Let $\mathcal{E}$ be a coherent analytic sheaf on $Y \times X$. For $y \in Y$ we denote by $\mathcal{E}(y)$ the analytic restriction of $\mathcal{E}$ on $X = y \times X$; $\mathcal{E}(y)$ is the analytic inverse image of $\mathcal{E}$ to the injection $X = y \times X \hookrightarrow Y \times X$. If $f: Z \rightarrow Y$ is a holomorphic mapping then we set $\mathcal{E}_Z := (f \times \text{id}_X)^*\mathcal{E}$. We have $\mathcal{E}_Z(z) = \mathcal{E}(f(z))$ for $z \in Z$.

Let the coherent analytic sheaf $\mathcal{E}$ on $Y \times X$ be Y -flat, then $r(\mathcal{E}(y), X')$ is locally constant on Y for every irreducible component X' of X . Therefore

$$R_{X'}^p(Y, \mathcal{E}) := \{y \in Y : r(\mathcal{E}(y), X') = p\}$$

is an open and closed subset of Y for every p . Let $\check{\mathcal{R}}$ be a coherent subsheaf of $\mathcal{E}$, so that $\mathcal{R} := \mathcal{E}/\check{\mathcal{R}}$ is also Y -flat. Then $\check{\mathcal{R}}$ is Y -flat too and we have $\mathcal{R}(y) = \mathcal{E}(y)/\check{\mathcal{R}}(y)$ in a natural way. Let $f: Z \rightarrow Y$ be a holomorphic mapping. Then $\mathcal{E}_Z$ and $\mathcal{R}_Z$ are Z -flat and we have $\mathcal{R}_Z = \mathcal{E}_Z/\check{\mathcal{R}}_Z$ in a natural way.

Let $\mathcal{F}$ be a coherent analytic sheaf on X . Then the analytic inverse image $\mathcal{F}^Y := \pi^*\mathcal{F}$ to the projection $\pi: Y \times X \rightarrow X$ is Y -flat and we have $\mathcal{F}^Y(y) = \mathcal{F}$ in a natural way. In the following let $\check{\mathcal{R}}$ be a coherent subsheaf of Ω^Y , such that $\mathcal{R} := \Omega^Y/\check{\mathcal{R}}$ is Y -flat. Let X' be an irreducible component of X . We set:

$$F_{X'}^p(Y, \check{\mathcal{R}}) := \{y \in Y : \check{\mathcal{R}}(y) \text{ is a } X'\text{-foliation of codimension } p\},$$

$$F_{X'}(Y, \check{\mathcal{R}}) := \{y \in Y : \check{\mathcal{R}}(y) \text{ is a } X'\text{-foliation}\},$$

$$F(Y, \check{\mathcal{R}}) := \{y \in Y : \check{\mathcal{R}}(y) \text{ is a foliation}\}.$$

We will show:

0.3 THEOREM. $F_{X'}^p(Y, \check{\mathcal{R}})$ is an analytic subset of Y .

By 0.3 we get:

0.4 COROLLARY.

- (1) $F_{X'}(Y, \check{\mathcal{R}})$ is an analytic subset of Y ;
- (2) $F(Y, \check{\mathcal{R}})$ is an analytic subset of Y .

Because $\mathcal{R}$ is Y -flat we get the following:

0.5 REMARK. $S(\Omega^Y: \check{\mathcal{R}}) \cap (y \times X) = S(\Omega: \check{\mathcal{R}}(y))$.

In 0.5 we identify $X = y \times X$.

Let $f: Z \rightarrow Y$ be a holomorphic mapping. Then $\Omega_Z^Y = \Omega^Z$ and we obviously get:

0.6 REMARK. $F_{X'}^p(Z, \check{\mathcal{R}}_Z) = f^{-1}(F_{X'}^p(Y, \check{\mathcal{R}}))$.

$F_{X'}^p(Y, \check{\mathcal{R}})$ is an analytic subset of $R_{X'}^p(Y, \check{\mathcal{R}})$. Using the homomorphism δ^p we construct a canonical complex structure on $F_{X'}^p(Y, \check{\mathcal{R}})$ and show:

0.7 THEOREM. The holomorphic mapping $f: Z \rightarrow Y$ induces a holomorphic mapping $F_{X'}^p(Z, \check{\mathcal{R}}_Z) \rightarrow F_{X'}^p(Y, \check{\mathcal{R}})$ related to the canonical complex structures.

We can apply $F(Y, \check{\mathcal{R}})$ with the structure given by the spaces $F_{X'}^p(Y, \check{\mathcal{R}})$. We get the following (comp. [Dou]):

0.8 COROLLARY. Let X be compact and let H be the Douady space of $\Omega, \mathcal{R}$ the universal Douady sheaf of Ω . Then the complex subspace $F(H, \check{\mathcal{R}})$ of H has the following property:

If Z is a complex space and $\mathcal{S} = \Omega^Z/\check{\mathcal{S}}$ a coherent and Z -flat sheaf, then the unique holomorphic mapping $f: Z \rightarrow H$ with $\mathcal{S} = \check{\mathcal{R}}_Z$ factorizes over $F(Z, \check{\mathcal{S}}), F(H, \check{\mathcal{R}})$.

In this sense $F(H, \check{\mathcal{R}})$ is a moduli space of the foliations on X .

0.9 PROPOSITION. Let X be a connected compact manifold. Then the following sets are complements of analytic subsets in $F(Y, \check{\mathcal{R}})$:

$$F_r(Y, \check{\mathcal{R}}) := \{y \in Y: \check{\mathcal{R}}(y) \text{ is a regular foliation}\},$$

$$F_f(Y, \check{\mathcal{R}}) := \{y \in F(Y, \check{\mathcal{R}}): \check{\mathcal{R}}(y) \text{ is locally free}\},$$

$${}^k F(Y, \check{\mathcal{R}}) := \{y \in F(Y, \check{\mathcal{R}}): \text{codim } S(\Omega: \check{\mathcal{R}}(y)) \geq k\}.$$

$F_f(Y, \check{\mathcal{R}}) \cap {}^2 F(Y, \check{\mathcal{R}})$ is the set of points $y \in Y$ such that $\check{\mathcal{R}}(y)$ is a free foliation in the sense of [B/R]. By the theorem of Frobenius-Malgrange all foliations $\check{\mathcal{R}}(y), y \in F_f(Y, \check{\mathcal{R}}) \cap {}^3 F(Y, \check{\mathcal{R}})$ are locally integrable.

In an earlier version of this paper ([Re]) I gave a proof of 0.3 by Banach analytic technics. With these technics G. Pourcin has gotten results similar to those in this paper ([Pou]). Similar results were also proved by X. Gomez-Mont ([GM]). He considers foliations defined by vector fields; his technics are closer to those of this paper. The advantage of my approach is the simplicity. For 0.3, the canonical complex structure on $F_{X'}^p(Y, \check{\mathcal{R}})$, $F(Y, \check{\mathcal{R}})$ and 0.7 we need no compactness.

§ 1. We prove 0.3: Because 0.3 is a local theorem related to Y and because of 0.2 we may assume the following:

Y is an analytic subspace of a polycylinder $P = \{s \in \mathbb{C} : |s| < \sigma\}^m$ in $\mathbb{C}^m$ defined by an ideal sheaf $\mathcal{I}$. We denote the coordinates of $\mathbb{C}^m$ by $y = (y_1, \dots, y_m)$. $Y = R_X^p(Y, \tilde{\mathcal{R}})$.

X is a polycylinder $\{t \in \mathbb{C} : |t| < \tau\}^n$ in $\mathbb{C}^n$. We denote the coordinates of $\mathbb{C}^n$ by $x = (x_1, \dots, x_n)$. $S(\Omega^Y : \tilde{\mathcal{R}}) = \emptyset$.

We can interpret $(\bigwedge^r \Omega)^P$ as a submodule of $\bigwedge^r \Omega_{P \times X} : (\bigwedge^r \Omega)^P$ consists of all r -forms η of the form $\eta = \sum_{1 \leq v_1 < \dots < v_r \leq n} \eta_{v_1 \dots v_r}(y, x) dx_{v_1} \wedge \dots \wedge dx_{v_r}$.

We notice this by $\eta = \eta(y, x; dx)$. We get $(\bigwedge^r \Omega)^Y = (\bigwedge^r \Omega)^P / \mathcal{I} \cdot (\bigwedge^r \Omega)^P$.

We may assume, that there are forms $\eta^1, \dots, \eta^q \in \Gamma(P \times X, \Omega^P)$ generating $\tilde{\mathcal{R}}$ everywhere. We shorten $F := F_X^p(Y, \tilde{\mathcal{R}})$. For $y^0 \in Y$ we get:

$$y^0 \in F \Leftrightarrow \delta^p(\eta^{j_0}(y^0, x; dx); \eta^{j_1}(y^0, x; dx), \dots, \eta^{j_p}(y^0, x; dx)) = 0$$

$$\text{for all } 1 \leq j_0, j_1, \dots, j_p \leq q.$$

We consider an arbitrary system $\omega^0, \omega^1, \dots, \omega^p \in \Gamma(P \times X, \Omega^P)$. By fixing y we define

$$\mathfrak{P}^p(\omega^0; \omega^1, \dots, \omega^p)$$

$$:= \delta^p(\omega^0(y, x; dx); \omega^1(y, x; dx), \dots, \omega^p(y, x; dx)) \in \Gamma(P \times X, (\bigwedge^{p+2} \Omega)^P).$$

We have

$$\mathfrak{P}^p(\omega^0; \omega^1, \dots, \omega^p) = \sum_{1 \leq v_1 < \dots < v_{p+2} \leq n} A_{v_1 \dots v_{p+2}} dx_{v_1} \wedge \dots \wedge dx_{v_{p+2}}$$

with holomorphic coefficients

$$A_{v_1 \dots v_{p+2}} = A_{v_1 \dots v_{p+2}}(\omega^0; \dots, \omega^p) \in \Gamma(P \times X, \mathcal{O}_{P \times X}).$$

We form the serial expansions

$$A_{v_1 \dots v_{p+2}} = \sum_{l \in \mathbb{N}^n} A_{v_1 \dots v_{p+2}; l} x^l$$

with holomorphic coefficients $A_{v_1 \dots v_{p+2}; l} = A_{v_1 \dots v_{p+2}; l}(\omega^0, \dots, \omega^p) \in \Gamma(P, \mathcal{O}_P)$. With these notations we get:

The set F is defined on Y by the following (infinite) system of holomorphic functions:

$$A_{v_1 \dots v_{p+2}; l}(\eta^{j_0}; \eta^{j_1}, \dots, \eta^{j_p});$$

$$1 \leq j_0, j_1, \dots, j_p \leq q, 1 \leq v_1 < \dots < v_{p+2} \leq n, l \in \mathbb{N}^n.$$

F is an analytic subset of Y .

§ 2. We refine the considerations of § 1. We can interpret $\mathfrak{Y}^p$ as a homomorphism of sheaves (of groups) $\mathfrak{Y}^p: (\Omega^P)^{p+1} \rightarrow (\bigwedge^{p+2} \Omega)^P$. Obviously $\mathfrak{Y}^p$ induces a homomorphism $\mathfrak{Y}_Y^p: (\Omega^Y)^{p+1} \rightarrow (\bigwedge^{p+2} Y)$, which does not depend on the representation of Y as a subspace of a number space. Let $\mathcal{J}$ be the ideal sheaf on P , defined by $\mathcal{J}$ and the functions $A_{v_1 \dots v_{p+2}; l}(\eta^{j_0}; \eta^{j_1}, \dots, \eta^{j_p})$ (comp. [Fr]). We apply F with the structure sheaf $\mathcal{H} := \mathcal{O}_P / \mathcal{J}$ and consider the natural injection $\iota: F \rightarrow Y$.

2.1 PROPOSITION. $\mathcal{H}$ is the maximal complex structure on F , such that $\mathfrak{Y}_{(F, \mathcal{H})}^p | (\tilde{\mathcal{R}}_{(F, \mathcal{H})})^{p+1} = 0$, i.e.

- (1) $\mathfrak{Y}_{(F, \mathcal{H})}^p | (\tilde{\mathcal{R}}_{(F, \mathcal{H})})^{p+1} = 0$;
- (2) if $(F, \tilde{\mathcal{H}})$ is an analytic subspace of Y , such that $\mathfrak{Y}_{(F, \tilde{\mathcal{H}})}^p | (\tilde{\mathcal{R}}_{(F, \tilde{\mathcal{H}})})^{p+1} = 0$, then $(F, \tilde{\mathcal{H}})$ is an analytic subspace of $(F, \mathcal{H})$.

Proof. (1) Let $\omega^0, \omega^1, \dots, \omega^p \in \Gamma(P \times X, \Omega)$ induce elements of $\Gamma(F \times X, \tilde{\mathcal{R}}_H)$. We may assume, that there are representations

$$\omega^k = \sum_{j=1}^q a_{kj} \eta^j \bmod \Gamma(P \times X, \mathcal{J} \cdot \Omega^p).$$

Because of

$$\eta^{j_0} \wedge \eta^{j_1} \wedge \dots \wedge \eta^{j_p} \in \Gamma(P \times X, \mathcal{J} \cdot (\bigwedge^{p+1} \Omega)^P)$$

we get $\bmod \Gamma(P \times X, \mathcal{J} \cdot (\bigwedge^{p+2} \Omega)^P)$:

$$\mathfrak{Y}^p(\omega^0; \omega^1, \dots, \omega^p) = \sum_{1 \leq j_0, \dots, j_p \leq q} a_{0j_0} \dots a_{pj_p} \mathfrak{Y}^p(\eta^{j_0}; \eta^{j_1}, \dots, \eta^{j_p}).$$

Therefore we get

$$A_{v_1 \dots v_{p+2}; l}(\omega^0; \omega^1, \dots, \omega^p) \in \Gamma(P, \mathcal{J}), \mathfrak{Y}^p(\omega^0; \omega, \dots, \omega^p) \in \Gamma(P \times X, \mathcal{J} \cdot (\bigwedge^{p+2} \Omega)^P).$$

(2) Let $\tilde{\mathcal{J}}$ be the ideal sheaf of $\mathcal{O}_P$ defining $\tilde{\mathcal{H}}$. We show $\mathcal{J} \subset \tilde{\mathcal{J}}$. We have

$$\mathfrak{Y}^p(\eta^{j_0}; \eta^{j_1}, \dots, \eta^{j_p}) \in \Gamma(P \times X, \tilde{\mathcal{J}} \cdot (\bigwedge^{p+2} \Omega)^P)$$

and therefore

$$A_{v_1 \dots v_{p+2}; l}(\eta^{j_0}; \eta^{j_1}, \dots, \eta^{j_p}) \in \Gamma(P, \tilde{\mathcal{J}}).$$

Now we consider an arbitrary reduced complex space X and an arbitrary complex space Y . We again set $F := F_X^p(Y, \tilde{\mathcal{R}})$.

Let $y^0 \in F$. We consider $X'(y^0) := X \setminus S(\Omega: \tilde{\mathcal{R}}(y^0))$. If $x^0 \in X'(y^0)$ then we can realize the situation of § 1 related to open neighbourhoods V_0 of y^0 and U_0 of x^0 . Let $\mathcal{H}(x^0)$ denote the structure sheaf of $V_0 \cap F$ according to 2.1. We get $\mathcal{H}(x^0)_{y^0} = \mathcal{H}(x)_{y^0}$ for every $x \in U_0$. Because $X'(y^0)$ is connected, we get $\mathcal{H}(x^0)_{y^0} = \mathcal{H}(x)_{y^0}$ for every $x \in X'(y^0)$. Let $\mathcal{H}$ be the structure sheaf on F defined by $\mathcal{H}_y := \mathcal{H}(x)_y, x \in X'(y)$.

If $\tilde{\mathcal{H}}$ is any structure sheaf on F such that $(F, \tilde{\mathcal{H}})$ is an analytic subspace of Y we get by 0.5

$$S(\Omega^{(F, \tilde{\mathcal{H}})}: \tilde{\mathcal{R}}_{(F, \tilde{\mathcal{H}})}) = S(\Omega^Y: \tilde{\mathcal{R}}) \cap (F \times X).$$

We set

$$(F \times X)^0 := (F \times X) \setminus S(\Omega^Y: \tilde{\mathcal{R}}).$$

By 2.1 we get:

2.2 PROPOSITION. $\mathcal{H}$ is the maximal complex structure on F , such that $\mathfrak{g}_{(F, \mathcal{H})}^p | (\tilde{\mathcal{R}}_{(F, \mathcal{H})} | (F \times X)^0)^{p+1} = 0$.

We prove 0.7: We may assume that Z, X and Y, X fulfill the assumptions of § 1 simultaneously. We denote the objects related to Z by the index Z . Further we may assume, that f is given by a holomorphic mapping $g: P_Z \rightarrow P$ with $g^*\mathcal{J} \subset \mathcal{J}_Z$. We consider $\omega^j(z, x; dx) := \eta^j(g(z), x; dx)$. Then ω^j generates an element of $\Gamma(Z \times X, \tilde{\mathcal{R}}_Z)$ and we get

$$\begin{aligned} & A_{v_1 \dots v_{p+2}; l}(\eta^{j_0}; \eta^{j_1}, \dots, \eta^{j_p}) \circ (f \times \text{id}_X) \\ &= A_{v_1 \dots v_{p+2}; l}(\omega^{j_0}; \omega^{j_1}, \dots, \omega^{j_p}) \in \Gamma(P_Z, \mathcal{J}_Z). \end{aligned}$$

We prove 0.9: Let $\pi: Y \times X \rightarrow Y$ be the projection then $\pi(S(\Omega^Y: \tilde{\mathcal{R}}))$ and $\pi(S(\tilde{\mathcal{R}}))$ are analytic subsets of Y . $\{(y, x) \in Y \times X: \dim_{(y, x)}(S(\Omega^Y: \tilde{\mathcal{R}}) \cap (y \times X)) > \dim X - k\}$ is analytic and therefore $\{y \in Y: \text{codim } S(\Omega: \tilde{\mathcal{R}}(y)) < k\}$ too.

REFERENCES

- [B·R] BOHNHORST, G. und H.-J. REIFFEN. Holomorphe Blätterungen mit Singularitäten. *Math. Gottingensis* 5 (1985).
- [Dou] DOUADY, A. Le problème des modules pour les sous-espaces analytiques compacts d'un espace analytique donné. *Ann. Inst. Fourier* 16 (1) (1966), 1-95.
- [Fr] FRISCH, J. Points de platitude d'un morphisme d'espaces analytiques; *Inv. math.* 4 (1967), 118-138.
- [GM] GOMEZ-MONT, X. The Transverse Dynamics of a Holomorphic Flow. *Pub. Prel. dal Inst. de Mat., Univ. Nac. Aut. México, Núm. 109* (1986).
- [Pou] POURCIN, G. Deformations of coherent foliations on a compact normal space. Preprint (1986). Deformations of Singular Holomorphic Foliations on Reduced Compact $\mathbb{C}$ -analytic Spaces. Preprint (1986).
- [Re] REIFFEN, H.-J. The Variety of Moduli of Foliations on a Compact Complex Space. *Osn. Schriften z. Math., Reihe P, Heft 89* (1986).

(Reçu le 26 novembre 1986)

Hans-Jörg Reiffen

Universität Osnabrück
Fachbereich Mathematik/Informatik
Albrechtstrasse 28
D-45 Osnabrück