

3. Elliptic spaces

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **39 (1993)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **27.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

It remains to deduce the inequality (2) from (2.3). If the inequality (2) holds for some power series $h(z)$ it will also hold for $h(z^k)$, at the cost of replacing K by $K^{\frac{1}{2k}}$. By (2.3) we are thus reduced to showing that the power series

$$\sum_{i=0}^{\infty} q_i z^i = \prod_{i=0}^{\infty} (1 + z^i)$$

satisfies (2). But this is an immediate consequence of a theorem of Hardy and Ramanujan [10]. $\square$

COROLLARY OF PROOF. *If G satisfies the hypotheses of Theorem 2.1 (2) then for some $k \in \mathbb{N}$,*

$$G(z) \underset{c}{\geq} \prod_{i=1}^{\infty} [1 + (z^k)^i]. \quad \square$$

3. ELLIPTIC SPACES

In this section we establish the ellipticity of the spaces listed in the introduction.

3.1. Finite simply connected H -spaces, X .

Because X is an H -space, $H_*(\Omega X; \mathbb{F}_p)$ is commutative, all p . Since it has finite depth [3; Theorem A] it is elliptic [7; Prop. 3.2]. Hence X is elliptic.

3.2. Simply connected homogeneous spaces, $G // H$.

We may suppose that G is simply connected, and hence elliptic by §3. The fibration $G \rightarrow G/H \rightarrow BH$ loops to the fibration $\Omega G \rightarrow \Omega(G/H) \rightarrow H$ in which $\pi_1(H)$ acts trivially in $H_*(\Omega G; \mathbb{F}_p)$ [1; Lemma 5.1]. Thus we can use the Serre spectral sequence to deduce polynomial growth for $H_*(\Omega(G/H); \mathbb{F}_p)$ from the same property for $H_*(\Omega G; \mathbb{F}_p)$.

3.3. Fibrations $F \rightarrow X \rightarrow B$ with F, B elliptic.

Here all spaces are simply connected and we can apply the Serre spectral sequence to deduce that $H_*(X; \mathbb{Z})$ is concentrated in finitely many degrees, and finitely generated in each. Hence X has the weak homotopy type of a finite CW complex. Loop the fibration $F \rightarrow X \rightarrow B$ and use the fact that $H_*(\Omega F; \mathbb{F}_p)$ and $H_*(\Omega B; \mathbb{F}_p)$ grow polynomially to deduce the same property for $H_*(\Omega X; \mathbb{F}_p)$.

3.4. *Simply connected Poincaré complexes X with $H^*(X; \mathbb{F}_p)$ at most doubly generated.*

Suppose $p \neq 2$ and $H = H^*(X; \mathbb{F}_p)$ contains an element of odd degree. Then it has an odd generator α . Using Poincaré duality it is easy to see that there are only three possibilities for the algebra H :

$$H = \Lambda \alpha \quad \text{or} \quad \Lambda \alpha \otimes \Lambda \beta \quad \text{or} \quad \Lambda \alpha \otimes \mathbb{F}_p[\beta] / \beta^k.$$

In each case a simple, classical computation [11] produces $\text{Tor}^H(\mathbb{F}_p, \mathbb{F}_p)$ and shows that it grows polynomially. Since the Eilenberg-Moore spectral sequence converges from $\text{Tor}^H(\mathbb{F}_p, \mathbb{F}_p)$ to $H^*(\Omega X; \mathbb{F}_p)$, $H^*(\Omega X; \mathbb{F}_p)$ also has this property.

In all other cases ($p = 2$ or H concentrated in even degrees) H is a commutative local ring in the classic sense. Because H satisfies Poincaré duality it is a Gorenstein ring. Now a theorem of Wiebe [12; Korollar p. 268] asserts (because H has at most two generators) that H is a polynomial algebra divided by a regular sequence. It is thus easy (and classical [11]) to compute $\text{Tor}^H(\mathbb{F}_p, \mathbb{F}_p)$, and deduce that it grows polynomially. Hence so does $H_*(\Omega X; \mathbb{F}_p)$.

3.5. *Simply connected Dupin hypersurfaces E in S^{n+1} .*

In [9; Table 2.1] are listed the possibilities for $H_*(E; \mathbb{Z})$. We divide these into three cases, using the notation of [9].

Case (a): E has the same integral homology as S^k or as $S^k \times S^l$.

In this case Poincaré duality shows that E has the same integral cohomology ring as S^k or as $S^k \times S^l$, and we can apply 3.4.

Case (b): E has the rational homotopy type of $A_3(2)$, $A_3(4)$, $A_3(8)$, $A_4(2)$ or $A_6(2)$.

In these cases the calculations of [9; §6] show explicitly that the ring $H^*(E; \mathbb{Z})$ is torsion free and generated by two elements. Thus each $H^*(E; \mathbb{F}_p)$ is doubly generated, and we can apply Wiebe's result as in 3.4.

Case (c): E has the integral homology of $S^k \times S^l \times S^{k+l}$, with $k < l$.

We need, in this case, to recall from [9; §2] that there are linear sphere bundles

$$S^k \rightarrow E \xrightarrow{\pi_0} B \quad \text{and} \quad S^l \rightarrow E \xrightarrow{\pi_1} B_1$$

with B_0, B_1 simply connected focal submanifolds of S^{n+1} . Moreover if D_0, D_1 denote the corresponding disk bundles with boundary E then $S^{n+1} = D_0 \cup_E D_1$.

Fix $p \geq 0$ and consider the Serre spectral sequence for the fibration $S^k \rightarrow E \rightarrow B_0$ with coefficients in $\mathbb{F}_p$. If this fails to collapse then $H^k(\pi_0): H^k(B_0; \mathbb{F}_p) \rightarrow H^k(E; \mathbb{F}_p)$ is surjective. Since $l > k$ it is always true that $H^k(\pi_1)$ is surjective. Choose classes $\alpha \in H^k(B_0; \mathbb{F}_p)$, $\beta \in H^k(B_1; \mathbb{F}_p)$ mapping to the same non-zero class in $H^k(E; \mathbb{F}_p)$. The Mayer-Vietoris sequence for the decomposition $S^{n+1} = D_0 \cup_E D_1$ then gives a class $\gamma \in H^k(S^{n+1}; \mathbb{F}_p)$ restricting to α and β , which is absurd.

Thus the spectral sequence for $S^k \rightarrow E \rightarrow B_0$ collapses and so $H_*(B_0; \mathbb{F}_p) \cong H_*(S^l \times S^{l+k}; \mathbb{F}_p)$. Using Poincaré duality for B_0 we see that $H^*(B_0; \mathbb{F}_p)$ and $H^*(S^l \times S^{l+k}; \mathbb{F}_p)$ are isomorphic as graded algebras. Thus B_0 is elliptic by 3.4 and E is elliptic by 3.3.

3.6. Simply connected closed manifolds M with a smooth action by a compact Lie group G , having a simply connected codimension one orbit.

Here we may assume G is connected. Let the orbit be G/K , and convert the inclusion of G/K into a fibration $F \rightarrow G/K \rightarrow M$. From [9; Table 1.5] we see that for any p , $\dim H_i(F; \mathbb{F}_p) \leq 2$, all i . Thus applying the Serre spectral sequence to the fibration $\Omega(G/K) \rightarrow \Omega M \rightarrow F$ and using 3.1 for G/K we see that $H_*(\Omega M; \mathbb{F}_p)$ grows polynomially.

3.7. Simply connected manifolds $M \# N$ with each of the rings $H^(M; \mathbb{Z})$, $H^*(N; \mathbb{Z})$ generated by a single class.*

By Van Kampen's theorem both M and N are simply connected, and so their fundamental cohomology classes are not torsion. Since each ring is monogenic, $H^*(M; \mathbb{Z})$ and $H^*(N; \mathbb{Z})$ are torsion free. Thus $H^*(M; \mathbb{F}_p)$ and $H^*(N; \mathbb{F}_p)$ are also monogenic, and so $H^*(M \# N; \mathbb{F}_p)$ is doubly generated. Now apply 3.4.

REFERENCES

- [1] BROWDER, W. On differential Hopf algebras. *Trans. Amer. Math. Soc.* 107 (1963), 153-176.
- [2] FELIX, Y., S. HALPERIN and J.-C. THOMAS. The homotopy Lie algebra for finite complexes. *Publ. de l'Institut des Hautes Etudes Scientifiques* 56 (1983), 387-410.
- [3] FELIX, Y., S. HALPERIN, J.-M. LEMAIRE and J.-C. THOMAS. Mod p loop space homology. *Invent. Math.* 95 (1989), 247-262.
- [4] FELIX, Y., S. HALPERIN and J.-C. THOMAS. Hopf algebras of polynomial growth. *J. of Algebra* 125 (1989), 408-417.
- [5] FELIX, Y., S. HALPERIN and J.-C. THOMAS. Loop space homology of spaces of LS category one and two. *Math. Ann.* 287 (1990), 377-387.