

4. \bar{W} IS CONNECTED

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **39 (1993)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **28.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

From Lemma 3.5, it follows then that

$$B \subseteq \bigcup_{\varepsilon_1, \dots, \varepsilon_{35} \in \{0, 1\}} \left[\left(\sum_{j=1}^n \varepsilon_j z^j \right) + \left(\frac{1}{2} + O(|\delta|) \right) z^n B \right]$$

so for sufficiently small δ , we may apply Lemma 3.1 to deduce $z \in \bar{W}$. \square

We now combine all the results of this section.

THEOREM 3.1. *There is an open neighborhood of $\{z: |z| = 1, z \neq 1\}$ contained in \bar{W} .*

Proof. Apply Propositions 3.2 and 3.3. \square

COROLLARY 3.1. *If $z \in (-1, -1 + \delta)$ for sufficiently small δ then z is a multiple zero of some 0, 1 power series.*

Proof. By Theorem 3.1, if δ is small enough we can pick 0, 1 power series f_n and zeros z_n of f_n such that $z_n \notin \mathbf{R}$ and $z_n \rightarrow z$ as $n \rightarrow \infty$. By taking a subsequence we may assume that the coefficient of z^k in f_n is eventually constant for large n , for each k . By a Rouché's Theorem argument, the pairs of zeros $\{z_n, \bar{z}_n\}$ of f_n must converge to (at least) a double zero at z of $\lim_{n \rightarrow \infty} f_n$. \square

4. \bar{W} IS CONNECTED

Since W is countable, we cannot hope to prove W is connected. We prove instead that \bar{W} is connected. First we need some topological lemmas.

Give $\{0, 1\}$ the discrete topology and $\{0, 1\}^\omega$ the product topology, as usual. If $v = (v_1, v_2, \dots, v_n)$ is a finite vector of 0's and 1's, let S_v be the set of sequences in $\{0, 1\}^\omega$ which start with v . The following lemma is the key ingredient in the connectivity proof.

LEMMA 4.1. *Let Y be a topological space. Suppose $f: \{0, 1\}^\omega \rightarrow Y$ is a continuous map such that*

$$(4.1) \quad f(S_{v0}) \cap f(S_{v1}) \neq \emptyset$$

for all $v \in \{0, 1\}^n$, and all $n \geq 0$. (Here $v0$ denotes the vector v with 0 appended, etc.) Then the image of f is path connected.

Proof. Let $w(0) = f(x'_0)$ and $w(1) = f(x_1)$ be elements of the image we wish to connect by a path. Find $x_{1/2}, x'_{1/2} \in \{0, 1\}^\omega$ such that $x'_0, x_{1/2}$ have the same first coordinate, and $x'_{1/2}, x_1$ have the same first coordinate and $f(x_{1/2}) = f(x'_{1/2})$. (If x'_0, x_1 have the same first coordinate, take $x_{1/2} = x'_{1/2} = x'_0$; otherwise apply the hypothesis (4.1) with v as the empty vector.) Let $w(1/2)$ be this common value.

Next find $x_{1/4}, x'_{1/4} \in \{0, 1\}^\omega$, using the same argument, such that $x'_0, x_{1/4}$ agree in the first two coordinates, $x'_{1/4}, x_{1/2}$ agree in the first two coordinates, and $f(x_{1/4}) = f(x'_{1/4})$. Let $w(1/4)$ be this common value. Do the analogous thing at $3/4$.

By induction, we may continue to define $x_{d/2^n}, x'_{d/2^n}, w(d/2^n)$ at all dyadic rationals $d/2^n$ in $[0, 1]$, such that $x'_{d/2^n}$ and $x_{(d+1)/2^n}$ agree in the first n coordinates and

$$w(d/2^n) = f(x_{d/2^n}) = f(x'_{d/2^n}).$$

By induction, we see that all the x'_q with $q \in [d/2^n, (d+1)/2^n)$ agree in the first n coordinates. Hence for

$$r = \sum_{i=1}^{\infty} \varepsilon_i 2^{-i} \in [0, 1], \quad \varepsilon_i \in \{0, 1\}$$

not a dyadic rational, we may define

$$x_r = x'_r = \lim_{n \rightarrow \infty} x'_{\sigma(n)} \quad \text{where} \quad \sigma(n) = \sum_{i=1}^n \varepsilon_i 2^{-i},$$

and $w(r) = f(x_r)$. Then w maps $[d/2^n, (d+1)/2^n]$ into $f(S_v)$ where $v \in \{0, 1\}^n$ is the first n coordinates of $x'_r, r \in [d/2^n, (d+1)/2^n)$ and of $x_{(d+1)/2^n}$.

We now show that w is continuous at $r \in [0, 1]$. Let U be an open set of Y containing $w(r)$. Then $f^{-1}(U)$ contains S_v and $S_{v'}$ for some finite substrings v, v' of x_r, x'_r respectively, by continuity of f . By the last sentence of the previous paragraph it follows that

$$w^{-1}(U) \supseteq w^{-1}(f(S_v) \cup f(S_{v'}))$$

will contain a neighborhood of r .

Thus $w: [0, 1] \rightarrow \text{image}(f)$ is a continuous path, and $\text{image}(f)$ is path connected. \square

Let M be a topological space. Give M^n the product topology and let the symmetric group S_n act on M^n by permuting the coordinates. The space

M^n/S_n , which parameterizes n -element multisets, can be given the quotient topology.

LEMMA 4.2. *If $A \subseteq M^n/S_n$ is connected, and the multiset $\{P, P, \dots, P\}$ is in A for some $P \in M$, then the subset $B \subseteq M$ of all coordinates of points in A is connected.*

Proof. Suppose not. Then there are open sets $U, V \subseteq M$ such that $U \cap B$ and $V \cap B$ are disjoint nonempty sets with union B . Without loss of generality, $P \in U$. Let

$$\begin{aligned} U' &= U \times U \times \cdots \times U, \\ V' &= (V \times M \times M \times \cdots \times M) \\ &\quad \cup (M \times V \times M \times \cdots \times M) \\ &\quad \vdots \\ &\quad \cup (M \times M \times M \times \cdots \times V). \end{aligned}$$

Then U', V' are open sets in M^n which are stable under S_n , so they project to open sets U'', V'' in M^n/S_n . Also $A \subseteq U'' \cup V''$ since a point in A must have all coordinates in U , or else at least one coordinate in $B \setminus U \subseteq V$. Furthermore $P \in U'' \cap A$, and $V'' \cap A$ is nonempty also, since at least one point of A has a coordinate in V , since $V \cap B \neq \emptyset$. Finally $U'' \cap V'' \cap A = \emptyset$, since it is not possible for a point of A to have all coordinates in U , yet have some coordinate in V . This contradicts the connectedness of A . \square

THEOREM 4.1. \bar{W} is connected.

Proof. First we show that for $\delta \in (0, 1)$,

$$\bar{W}_\delta = (\bar{W} \cap \{z: |z| \leq 1\}) \cup \{z: 1 - \delta \leq |z| \leq 1\}$$

is connected. The idea is to apply Lemma 4.1 to the function f which assigns to $(\varepsilon_1, \varepsilon_2, \dots)$ the set of zeros of

$$1 + \varepsilon_1 z + \varepsilon_2 z^2 + \cdots$$

inside $\{z: |z| < 1 - \delta\}$. To make a continuous map of this requires some manipulation.

By Jensen's theorem, as was shown in Section 2, there is an upper bound n on the number of zeros that a power series with 0, 1 coefficients can have inside $\{z: |z| < 1 - \delta\}$. Let M be $\{z: |z| \leq 1\}$ with the annulus

$\{z: 1 - \delta \leq |z| \leq 1\}$ shrunk to a point P . (Therefore M is topologically a sphere.) To each power series $1 + \sum_{i=1}^{\infty} \varepsilon_i z^i$, $\varepsilon_i \in \{0, 1\}$, we assign the set of zeros inside $\{z: |z| < 1 - \delta\}$, (counted with multiplicities) and throw in extra copies of the point P as necessary to bring the total number of points to n . Since the order of these n elements of M is unspecified, we obtain a point of M^n/S_n . Let $f((\varepsilon_1, \varepsilon_2, \dots))$ be this point.

We claim that this map

$$f: \{0, 1\}^{\omega} \rightarrow M^n/S_n$$

is continuous. This follows easily from Rouché's theorem; if two power series agree in the first m coordinates for m sufficiently large then their zeros inside $\{z: |z| < 1 - \delta\}$ will be within ε . Some may escape or enter the disk, but this is not a problem, since in the topology on M , P is close to all points z with $|z|$ sufficiently near $1 - \delta$.

We next check condition (4.1) of Lemma 4.1. This is easily done using the following trick: given

$$v = (v_1, v_2, \dots, v_n) \in \{0, 1\}^n,$$

let $w = (v_1, v_2, \dots, v_n, 1, v_1, v_2, \dots, v_n)$. Then $v \in S_{v_0}$, $w \in S_{v_1}$, and $f(v) = f(w)$ (we extend v, w to infinite vectors by appending 0's), since

$$1 + v_1 z + v_2 z^2 + \dots + v_n z^n$$

and

$$\begin{aligned} &1 + v_1 z + v_2 z^2 + \dots + v_n z^n + z^{n+1} + v_1 z^{n+2} + \dots + v_n z^{2n+1} \\ &= (1 + v_1 z + v_2 z^2 + \dots + v_n z^n) (1 + z^{n+1}) \end{aligned}$$

have the same zeros inside $\{z: |z| < 1 - \delta\}$. Therefore we may apply Lemma 4.1 and deduce that the image of f is path connected.

Since $f((0, 0, \dots)) = (P, P, P, \dots, P)$, we may apply Lemma 4.2 with $A = \text{image}(f)$ to deduce that \bar{W}_δ with the annulus $\{z: 1 - \delta \leq |z| \leq 1\}$ shrunk to a point P is a connected subset of M . This is equivalent to the connectivity of \bar{W}_δ .

Since $\bar{W} \cap \{z: |z| \leq 1\}$ is the decreasing intersection of the compact connected sets $\bar{W}_{1/m}$, it too is connected. So is its image under $z \mapsto 1/z$. Finally, \bar{W} is the union of these two sets, which meet on the unit circle, so \bar{W} is connected as well. \square