

2.2 Homotopy types with a given cohomology ring

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **41 (1995)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **29.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Proof. W_o is characteristic for F if and only if $q_{\bar{F}} = \bar{F}^t(W_o)$.

In terms of a $\mathbf{Z}$ -basis $\{e_1, \dots, e_b\}$ for H the condition $q_{\bar{F}} \in \text{Im}(\bar{F}^t)$ translates into a simple rank condition over $\mathbf{Z}_{/2}$: the $\mathbf{Z}_{/2}$ -rank of the $b \times \binom{b+1}{2}$ -matrix A representing $\bar{F}^t$ must be equal to the $\mathbf{Z}_{/2}$ -rank of the matrix A extended by the column $(\bar{e}_i \cdot \bar{e}_j \cdot (\bar{e}_i + \bar{e}_j))_{1 \leq i \leq j \leq b}$

EXAMPLE 3. Let $H = \mathbf{Z}e_1 \oplus \mathbf{Z}e_2$ be free of rank 2, $F \in S^3 H^\vee$ given by $e_1^3 = a$, $e_1^2 e_2 = b$, $e_1 e_2^2 = c$, $e_2^3 = d$ with $a, b, c, d \in \mathbf{Z}$. The rank condition becomes

$$rk_2 \begin{bmatrix} \bar{a} & \bar{b} \\ \bar{c} & \bar{d} \\ \bar{b} & \bar{c} \end{bmatrix} = rk_2 \begin{bmatrix} \bar{a} & \bar{b} & \bar{0} \\ \bar{c} & \bar{d} & \bar{0} \\ \bar{b} & \bar{c} & \overline{b+c} \end{bmatrix}$$

2.2 HOMOTOPY TYPES WITH A GIVEN COHOMOLOGY RING

Our next task is to describe the set of oriented homotopy types of 1-connected, closed, oriented, 6-dimensional manifolds with a fixed torsion-free cohomology ring.

From Žubr's classification theorem we know that in algebraic terms this means the following: fix a non-negative integer r_o , a finitely generated free abelian group H_o , and a symmetric trilinear form $F_o \in S^3 H_o^\vee$ which admits characteristic elements.

Let $\mathcal{M}(r_o, H_o, F_o)$ be the set of 1-connected, closed, oriented, 6-dimensional manifolds X with $b_3(X) = 2r_o$, such that there exists an isomorphism $\alpha: H_o \rightarrow H^2(X, \mathbf{Z})$ with $\alpha^* F_X = F_o$. Denote by $\text{Aut}(F_o)$ the subgroup of $\mathbf{Z}$ -automorphisms of H_o which leave $F_o \in S^3 H_o^\vee$ invariant; $\text{Aut}(F_o)$ acts on pairs $(w, [l]) \in \bar{H}_o \times H_o^\vee / {}_{48}H_o^\vee / {}_{U_{F_o}}$ in a natural way:

$$\gamma \cdot (w, [l]) := (\gamma(w), (\gamma^{-1})^* [l]) .$$

Let ${}_{\text{Aut}(F_o)} \backslash \bar{H}_o \times H_o^\vee / {}_{48}H_o^\vee / {}_{U_{F_o}}$ be the set of $\text{Aut}(F_o)$ -orbits.

A manifold X in $\mathcal{M}(r_o, H_o, F_o)$ and an isomorphism $\alpha: H_o \rightarrow H^2(X, \mathbf{Z})$ with $\alpha^* F_X = F_o$ yields a well-defined $\text{Aut}(F_o)$ -orbit:

$$(\alpha^{-1}(w_2(X)), \alpha^* [p_1(X) + 24T]) \text{ (modulo } \text{Aut}(F_o) \text{)},$$

where $T \in H^4(X, \mathbf{Z})$ is an arbitrary integral lifting of $\tau(X) \in H^4(X, \mathbf{Z}_{/2})$.

The set of oriented homotopy types $\mathcal{M}(r_o, H_o, F_o) / \simeq$ of manifolds in $\mathcal{M}(r_o, H_o, F_o)$ can now be described in the following way:

PROPOSITION 3. *The assignment $X \mapsto (\alpha^{-1}(w_2(X)), \alpha^*[p_1(X) + 24T])$ (modulo $\text{Aut}(F_o)$) defines an injection.*

$$I: \mathcal{M}(r_o, H_o, F_o)/\simeq \rightarrow_{\text{Aut}(F_o)} \bar{H}_o \times H_o^\vee /_{48H_o^\vee} /_{U_{F_o}}.$$

Proof. Suppose X and X' are manifolds in $\mathcal{M}(r_o, H_o, F_o)$, $\alpha: H_o \rightarrow H^2(X, \mathbf{Z})$ and $\alpha': H_o \rightarrow H^2(X', \mathbf{Z})$ isomorphisms with $\alpha^*F_X = F_o$ and $(\alpha')^*F_{X'} = F_o$. X and X' have the same image under I iff there exists an automorphism $\gamma \in \text{Aut}(F_o)$ with $\gamma\alpha^{-1}(w_2(X)) = (\alpha')^{-1}w_2(X')$ and $(\gamma^{-1})^*\alpha^*[p_1(X) + 24T] = (\alpha')^*[p_1(X') + 24T']$. Consider $\beta := \alpha \circ \gamma \circ \alpha^{-1}: H^2(X, \mathbf{Z}) \rightarrow H^2(X', \mathbf{Z})$; β is obviously an isomorphism with $\beta^*F_{X'} = F_X$, $\beta w_2(X) = w_2(X')$, and $\beta^*[p_1(X') + 24T'] = [p_1(X) + 24T]$; but this means that the systems of invariants associated with X and X' are weakly equivalent, and therefore X and X' oriented homotopy equivalent.

A complete description of the set $\mathcal{M}(r_o, H_o, F_o)/\simeq$ i.e. of the image of I is only possible if the automorphism group $\text{Aut}(F_o)$ is known; this can be a serious problem, but we will see that the ‘general’ automorphism group is finite (and usually small), so that the next proposition gives a reasonable estimate for the number of elements in $\mathcal{M}(r_o, H_o, F_o)/\simeq$.

PROPOSITION 4. *Fix $r_o \in \mathbf{N}$, a finitely generated free abelian group H_o , and a symmetric trilinear form $F_o \in S^3 H_o^\vee$ which admits characteristic elements. Set $b := rk_{\mathbf{Z}} H_o$, $s := rk_{\mathbf{Z}/2}(\bar{F}_o^t)$, and let $t := rk_{\mathbf{Z}/2}(\cdot_{\bar{F}_o})$ be the $\mathbf{Z}/2$ -rank of the $\mathbf{Z}/2$ -linear square map $\cdot_{\bar{F}_o}: \bar{H}_o \rightarrow \bar{H}_o^\vee$ sending $\bar{u} \in \bar{H}_o$ to $\bar{u}^2 \in \bar{H}_o^\vee$. Then $\mathcal{M}(r_o, H_o, F_o)/\simeq$ contains at most 2^{2b-s-t} elements.*

Proof. Fix any admissible system of invariants $(r_o, H_o, w_o, \tau_o, F_o, p_o)$ for a manifold in $\mathcal{M}(r_o, H_o, F_o)$. Given (r_o, H_o, F_o) , we know from the last lemma that the possible elements w_o form a coset of $\text{Ker}(\bar{F}_o^t)$ in $\bar{H}_o$, so that there exist precisely 2^{b-s} such elements. It remains to count the classes $[l] \in H_o^\vee /_{48H_o^\vee} /_{U_{F_o}}$, such that the $\text{Aut}(F_o)$ -orbit of $(w_o, [p_o + 24T_o + l])$ lies in the image of I .

To understand the latter condition we fix integral liftings $W_o, \in H_o, T_o \in H_o^\vee$ of w_o and τ_o satisfying the admissibility conditions

- i) $W_o^3 \equiv (p_o + 24T_o)(W_o) \pmod{48}$
- ii) $p_o(x) \equiv 4x^3 + 6x^2 W_o + 3x W_o^2 \pmod{24} \quad \forall x \in H_o$.

Clearly the $\text{Aut}(F_o)$ -orbit of $(w_o, [p_o + 24T_o + l])$ lies in the image of I if and only if

$$\text{i}') \quad W_o^3 \equiv (p_o + 24T_o + l)(W_o) \pmod{48},$$

$$\text{ii}') \quad (p_o + l)(x) \equiv 4x^3 + 6x^2 W_o + 3x W_o^2 \pmod{24} \quad \forall x \in H_o,$$

which is equivalent to $l(W_o) \equiv 0 \pmod{48}$, and $l \equiv 0 \pmod{24H_o^\vee}$ because of i) and ii).

Now, by definition of the subgroup $U_{F_o} \subset H_o^\vee / {}_{48}H_o^\vee$ we have the following commutative diagram with exact rows and columns:

$$\begin{array}{ccccccc}
 & & \text{Ker}(\cdot \bar{F}_o) & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 0 \rightarrow \text{Ker}(24 \cdot \bar{F}_o) & \hookrightarrow & H_o / {}_2H_o & \xrightarrow{24 \cdot \bar{F}_o} & U_{F_o} & \rightarrow & 0 \\
 & & \downarrow \cdot \bar{F}_o & & \downarrow & & \\
 & 0 \rightarrow & H_o^\vee / {}_2H_o^\vee & \xrightarrow{24} & H_o^\vee / {}_{48}H_o^\vee & \rightarrow & H_o^\vee / {}_{24}H_o^\vee \rightarrow 0 \\
 & & \downarrow & & \downarrow & & \parallel \\
 0 \rightarrow \text{Coker}(\cdot \bar{F}_o) & \rightarrow & H_o^\vee / {}_{48}H_o^\vee / U_{F_o} & \rightarrow & H_o^\vee / {}_{24}H_o^\vee & \rightarrow & 0 \\
 & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & &
 \end{array}$$

The number of elements $[l] \in H_o^\vee / {}_{48}H_o^\vee / U_{F_o}$ to be counted coincides therefore with the cardinality of the kernel of the map $ev(w_o): \text{Coker}(\cdot \bar{F}_o) \rightarrow \mathbf{Z}_{/2}$ induced by evaluation in w_o . This number is at most $2^{b-t}(2^{b-t-1}$ if $w_o \neq 0$ and $t \neq b$).

COROLLARY 2. *If the $\mathbf{Z}_{/2}$ -rank $s = rk_{\mathbf{Z}_{/2}}(\cdot \bar{F}_o)$ is maximal, then $\mathcal{M}(r_o, H_o, F_o) / \simeq$ contains at most one class.*

Proof. Suppose $\cdot \bar{F}_o: \bar{H}_o \rightarrow \bar{H}_o^\vee$ is surjective; then $\bar{F}_o^t: \bar{H}_o \rightarrow S^2 \bar{H}_o^\vee$ must have a trivial kernel, since $\bar{h}\bar{x}^2 = 0$ for all $\bar{x} \in \bar{H}_o$ implies $\bar{h} = 0$ if every linear form is a square. But this means $s = t = b$, so that $\mathcal{M}(r_o, H_o, F_o) / \simeq$ has at most one element.

EXAMPLE 4. Let $H_o = \mathbf{Z}e_1 \oplus \mathbf{Z}e_2$, $e_1^3 = a$, $e_1^2 e_2 = b$, $e_1 e_2^2 = c$, $e_2^3 = d$. If $\bar{b} \equiv \bar{c} \pmod{2}$, and $\bar{a}\bar{d} - \bar{b}\bar{c} \equiv 1 \pmod{2}$, then $\mathcal{M}(r_o, H_o, F_o) / \simeq$ contains precisely one class for every $r_o \geq 0$.