

4.2 Standard constructions

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **41 (1995)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **29.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

REMARK 12. It is very likely that there exist non-integrable almost complex structures on manifolds X as above, but probably this is hard to prove. It is also not unlikely that the Chern numbers of complex 3-folds are not topological invariants. A possible way to check this would be, to run a computer search for 3-folds given by certain standard constructions.

4.2 STANDARD CONSTRUCTIONS

For later use we investigate the topological invariants of complex 3-folds which can be obtained by certain simple standard constructions like complete intersections, simple cyclic coverings, blow-ups of points and curves, and projective bundles.

PROPOSITION 11 (Libgober/Wood). *Let $X \subset \mathbf{P}^{3+r}$ be a smooth complete intersection of multidegree $\underline{d} = (d_1, \dots, d_r)$. Choose a normalized basis $e \in H^2(X, \mathbf{Z})$, and let $\varepsilon \in H^4(X, \mathbf{Z})$ be defined by $\varepsilon(e) = 1$. Then the invariants of X are:*

$$F_X(xe) = dx^3 \quad \text{where} \quad d = \prod_{i=1}^r d_i, \quad w_2(X) \equiv (4 + r - \sum_{i=1}^r d_i)e,$$

$$p_1(X) = d(4 + r - \sum_{i=1}^r d_i^2)\varepsilon, \quad \text{and}$$

$$b_3(X) = 4 - \frac{d}{6} \left[(4 + r - \sum_{i=1}^r d_i)^3 - 3(4 + r - \sum_{i=1}^r d_i)(4 + r - \sum_{i=1}^r d_i^2) + 2(4 + r - \sum_{i=1}^r d_i^3) \right].$$

Proof. [L/W].

PROPOSITION 12. *Let X be a smooth, 1-connected, complex projective 3-fold, and let $\pi: X' \rightarrow X$ be a simple cyclic covering of degree d branched along a non-singular ample divisor $B \in |L^{\otimes d}|$. X' is smooth, projective, 1-connected, and $\pi^*: H^2(X, \mathbf{Z}) \rightarrow H^2(X', \mathbf{Z})$ is an isomorphism. The invariants of X and X' are related by the formulae:*

$$(\pi^*)^* F_{X'} = dF_X, \quad w_2(X') - \pi^* w_2(X) \equiv (d-1)\pi^* c_1(L),$$

$$p_1(X') - \pi^* p_1(X) = (1-d)(1+d)\pi^* c_1(L)^2, \quad \text{and}$$

$$b_3(X') = db_3(X) + (d-1)(b_2(B) - 2b_2(X)).$$

Proof. X' is clearly smooth and projective. By a theorem of M. Cornalba $\pi: X' \rightarrow X$ is a 3-equivalence, i.e. $\pi_*: \pi_i(X') \rightarrow \pi_i(X)$ is bijective for $i \leq 2$, and surjective for $i = 3$ [Co]. X' is therefore 1-connected, and $\pi^*: H^2(X, \mathbf{Z}) \rightarrow H^2(X', \mathbf{Z})$ is an isomorphism. The relation between $F_{X'}$ and F_X is obvious, whereas the formula for $b_3(X')$ follows from $\pi_1(B) = \{1\}$ and standard properties of Euler numbers.

In order to calculate $w_2(X')$ and $p_1(X')$ we compute the Chern classes of X' : $c_1(X') - \pi^*c_1(X) = (1-d)\pi^*c_1(L)$, $c_2(X') - \pi^*c_2(X) = (1-d)\pi^*[c_1(X)c_1(L) - dc_1(L)^2]$.

The latter formulae follow from the description of X' as a divisor in the total space of the line bundle L .

EXAMPLE 9. Let X be a d -fold, simple cyclic covering of $\mathbf{P}^3$ branched along a smooth surface $B \subset \mathbf{P}^3$ of degree dl , $l \geq 1$. Let $e \in H^2(X, \mathbf{Z})$ correspond to the preimage of a plane in $\mathbf{P}^3$. The invariants of X are then given by:

$$F_X(xe) = dx^3, w_2(X) \equiv (4 + (1-d)l)e, p_1(X) = d[4 + (1-d)(1+d)l^2]\varepsilon \\ (\varepsilon(e) = 1), b_3(X) = (d-1)(d^2l^2 - 4dl + 6)dl.$$

PROPOSITION 13. Let $\sigma: \hat{X} \rightarrow X$ be the blow-up of a complex 3-fold X in a point, and let $e \in H^2(\hat{X}, \mathbf{Z})$ be the class of the exceptional divisor. The invariants of $\hat{X}$ and X are related by the following formulae:

$$F_{\hat{X}}(\sigma^*h + xe) = F_X(h) + x^3 \quad \forall h \in H^2(X, \mathbf{Z}), x \in \mathbf{Z}, w_2(\hat{X}) = \sigma^*w_2(X), \\ p_1(\hat{X}) = \sigma^*p_1(X) + 4(e^2 - \sigma^*c_1(X) \cdot e), b_3(\hat{X}) = b_3(X).$$

Proof. Standard arguments, see [G/H]. The Chern classes are related by $c_1(\hat{X}) = \sigma^*c_1(X) - 2e$, $c_2(\hat{X}) = \sigma^*c_2(X)$.

PROPOSITION 14. Let $\sigma: \hat{X} \rightarrow X$ be the blow-up of a complex 3-fold X along a smooth curve C of genus g , and let $e \in H^2(\hat{X}, \mathbf{Z})$ be the class of the exceptional divisor. The invariants of $\hat{X}$ and X are related by:

$$F_{\hat{X}}(\sigma^*h + xe) = F_X(h) - 3h \cdot Cx^2 - \deg N_{C/X}x^3 \quad \forall h \in H^2(X, \mathbf{Z}), \\ x \in \mathbf{Z}, w_2(\hat{X}) \equiv \sigma^*w_2(X) + e, p_1(\hat{X}) = \sigma^*p_1(X) + (e^2 - 2\sigma^*C), \\ b_3(\hat{X}) = b_3(X) + 2g.$$

Proof. [G/H]. The Chern classes are given by $c_1(\hat{X}) = \sigma^*c_1(X) - c$, $c_2(\hat{X}) = \sigma^*(c_2(X) + C) - \sigma^*c_1(X) \cdot e$.

PROPOSITION 15. Let E be a holomorphic vector bundle of rank 2 with Chern classes $c_i(E)$, $i = 1, 2$ over a 1-connected, compact complex surface Y , and let $\pi: \mathbf{P}(E) \rightarrow Y$ be the projective bundle of lines in the fibers of E . The cup-form of $\mathbf{P}(E)$ is given by

$$F_{\mathbf{P}(E)}(h + x\xi) = x[(3h^2) - (3c_1(E) \cdot h)x + (c_1(E)^2 - c_2(E))x^2],$$

where $\xi = c_1(\mathcal{O}_{\mathbf{P}(E)}(1))$, $h \in H^2(Y, \mathbf{Z})$, and $x \in \mathbf{Z}$. The other topological invariants of $\mathbf{P}(E)$ are:

$$\begin{aligned} w_2(\mathbf{P}(E)) &\equiv \pi^*(w_2(Y) + c_1(E)), p_1(E)) \\ &= \pi^*[c_1(Y)^2 - 2c_2(Y) + c_1(E)^2 - 4c_2(E)], b_3(\mathbf{P}(E)) = 0. \end{aligned}$$

Proof. The Leray-Hirsch theorem identifies the cohomology ring $H^*(\mathbf{P}(E), \mathbf{Z})$ with the ring $H^*(Y, \mathbf{Z})[\xi]/\langle \xi^2 + c_1(E) \cdot \xi + c_2(E) \rangle$; this determines the cup-form. In order to calculate the characteristic classes one uses the exact sequence $0 \rightarrow \mathcal{O}_{\mathbf{P}(E)} \rightarrow \pi^*E \otimes \mathcal{O}_{\mathbf{P}(E)}(1) \rightarrow T_{\mathbf{P}(E)} \rightarrow \pi^*T_Y \rightarrow 0$. $b_3(\mathbf{P}(E)) = 0$ follows from $b_1(Y) = 0$ and the Leray-Hirsch theorem.

4.3 EXAMPLES OF 1-CONNECTED NON-KÄHLERIAN 3-FOLDS

Recall that the Hessian of a symmetric trilinear form $F \in S^3 H^\vee$ on a free $\mathbf{Z}$ -module H of finite rank was defined as the composition $H_F: H \xrightarrow{F^t} S^2 H^\vee \xrightarrow{\text{disc}} \mathbf{Z}$. In terms of coordinates $\xi_1, \dots, \xi_b$ on H it is given by the determinant $\det\left(\frac{\partial^2 f}{\partial \xi_i \partial \xi_j}\right)$, where $f \in \mathbf{C}[H_{\mathbf{C}}]_3$ is the homogeneous cubic polynomial associated with F .

PROPOSITION 16. *Let F be a symmetric trilinear form whose Hessian vanishes identically. Then F is not realizable as cup-form of a Kählerian 3-fold.*

Proof. Let X be a complex 3-fold with a Kähler metric g . The Kähler class $[\omega_g] \in H^2(X, \mathbf{R})$ defines a multiplication map $\cdot [\omega_g]: H^2(X, \mathbf{R}) \rightarrow H^4(X, \mathbf{R})$, which is an isomorphism by the Hard Lefschetz Theorem [G/H]. In section 3.1 we have seen that this is not possible if the Hessian of the cup-form vanishes.

COROLLARY 6. *Cubic forms $f \in \mathbf{C}[H_{\mathbf{C}}]_3$ which depend on strictly less than $b = \text{rk}_{\mathbf{Z}} H$ variables are not realizable as cup-forms of Kählerian 3-folds with $b_2 = b$.*

By considering the Hessian of a cup-form over the reals one obtains further conditions.

DEFINITION 4. *Let $F \in S^3 H^\vee$ be a symmetric trilinear form on a free $\mathbf{Z}$ -module of rank b .*

The Hesse cone of F is the subset $\mathcal{H}_F \subset H_{\mathbf{R}}$ defined by $\mathcal{H}_F := \{h \in H_{\mathbf{R}} \mid (-1)^b \det(F^t(h)) < 0\}$.