

6. The τ -ABEL transform

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **45 (1999)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **28.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

5.5. COROLLARY. Let $E(\tau_l)$ denote the projection of $\mathcal{H}_{l,s}$ onto the K -isotypic subspace of type τ_l . Then

$$(5.43) \quad \zeta_{l,s}(g) = \frac{1}{d_l} \operatorname{tr}[E(\tau_l)T_{l,s}(g)].$$

$(T_{l,s}, \mathcal{H}_{l,s})$ is infinitesimally equivalent to a unitary representation if and only if the corresponding irreducible subquotient of $(\mathcal{H}'_{l,s})_K$ is unitarizable. The following theorem is thus a consequence of Theorems 5.1 and 5.3 and of Proposition 5.4.

5.6. THEOREM. $\zeta_{l,s} = \zeta_{l,-s}$ is positive definite if and only if one of the following cases occurs:

1. $s = i\nu$, $\nu \in \mathbf{R}$.
2. If $2l \geq 2n - 1$: $\pm s = s_j := 2(l - n - j) + 1$ for integers $j \geq 0$ so that $s_j > 0$. (discrete series)
3. If $2l < 2n - 1$: $s \in (2l - \rho + 2, -2l + \rho - 2)$. (complementary series)

The situation for s real and nonnegative is represented in Figure 6.1.

6. THE τ_l -ABEL TRANSFORM

Proposition 3.2 proves that the τ_l -Abel transform is a $*$ -homomorphism of $\mathcal{D}(G; \chi_l)$ into the convolution algebra $\mathcal{D}_+(\mathbf{R})$ consisting of the even C^∞ functions on $\mathbf{R}$ with compact support. The main theorem of this section states that the τ_l -Abel transform is also a bijection of $\mathcal{D}(G; \chi_l)$ onto $\mathcal{D}_+(\mathbf{R})$, and gives a formula for its inverse.

Identify A with $\mathbf{R}$ under the map $t \mapsto a_t$. Restriction to A then identifies $\mathcal{D}(G; \chi_l)$ with $\mathcal{D}_+(\mathbf{R})$. Let $\mathcal{D}([1, \infty))$ denote the set of the compactly supported C^∞ functions on $[1, \infty)$ (right differentiability at 1 is considered). Define a map H by

$$(Hf)(\cosh t) := f(a_t) \equiv f(t)$$




FIGURE 6.1

Positive definite $\zeta_{l,s}$ for real $s \geq 0$

for $f \in \mathcal{D}(G; \chi_l)$. Lemma 2 and its corollary in [Rou] imply

6.1. LEMMA. H is a bijection of $\mathcal{D}(G; \chi_l)$ onto $\mathcal{D}([1, \infty))$.

For every $\mu \in \mathbf{C}$ with $\Re \mu > 0$, the Weyl fractional integral transform of $\varphi \in \mathcal{D}([1, \infty))$ is defined by

$$(6.44) \quad \mathcal{W}_\mu \varphi(x) = \frac{1}{\Gamma(\mu)} \int_x^\infty \varphi(u)(u-x)^{\mu-1} du, \quad x \in [1, \infty).$$

Analytic continuation of $\mathcal{W}_\mu$ to $\Re \mu \leq 0$ is obtained via repeated integration by parts of (6.44): for every integer $m \geq 0$

$$\mathcal{W}_\mu \varphi(u) = \frac{(-1)^m}{\Gamma(\mu+m)} \int_u^\infty \frac{d^m \varphi}{dx^m}(x) (x-u)^{\mu+m-1} dx.$$

For every integer $m \geq 0$, the Gegenbauer transform (of dimension 4) of $\varphi \in \mathcal{D}([1, \infty))$ is defined by

$$(6.45) \quad \mathcal{G}_m \varphi(u) = \frac{4\pi}{m+1} \int_u^\infty \varphi(x) C_m^1\left(\frac{u}{x}\right) (x^2 - u^2)^{\frac{1}{2}} x \, dx, \quad u \in [1, \infty),$$

where

$$(6.46) \quad C_m^1(y) = (m+1)F\left(-m, m+2; \frac{3}{2}; \frac{1-y}{2}\right)$$

is the Gegenbauer polynomial of indices $(1, m)$ (cf. e.g. [E⁺], 3.15 (3)).

6.2. LEMMA ([K1], Theorem 3.2; [Dea], Formulas (28) and (29)).

1. For every $\mu \in \mathbf{C}$, $\mathcal{W}_\mu$ is a bijection of $\mathcal{D}([1, \infty))$ onto itself. The inverse mapping of $\mathcal{W}_\mu$ is $\mathcal{W}_{-\mu}$.

2. For every integer $m \geq 0$, $\mathcal{G}_m$ is a bijection of $\mathcal{D}([1, \infty))$ onto itself. The inverse mapping of $\mathcal{G}_m$ is given by

$$(6.47) \quad \mathcal{G}_m^{-1} \psi(x) = -\frac{1}{2\pi^2(m+1)} \frac{1}{x^2} \int_x^\infty \frac{d^3 \psi}{du^3}(u) C_m^1\left(\frac{u}{x}\right) (u^2 - x^2)^{\frac{1}{2}} du$$

for all $\psi \in \mathcal{D}([1, \infty))$ and all $x \in [1, \infty)$.

6.3. THEOREM. The τ_l -Abel transform is a bijection of $\mathcal{D}(G; \chi_l)$ onto $\mathcal{D}_+(\mathbf{R})$. It can be written as the composition

$$\mathcal{A}_l = \frac{(2\pi)^{2(n-1)}}{d_l^2} H^{-1} \circ \mathcal{W}_{2n-2} \circ \mathcal{G}_{2l} \circ H,$$

and its inverse is given by

$$\mathcal{A}_l^{-1} = \frac{d_l^2}{(2\pi)^{2(n-1)}} H^{-1} \circ \mathcal{G}_{2l}^{-1} \circ \mathcal{W}_{2-2n} \circ H.$$

Moreover, the support of the restriction to $A \equiv \mathbf{R}$ of $f \in \mathcal{D}(G; \chi_l)$ is contained in $[-R, R]$ if and only if the support of $\mathcal{A}_l f$ is contained in $[-R, R]$.

Proof. Identify the set of pure quaternions $w = i b + j c + k d \in \mathbf{H}$ with $\mathbf{R}^3$, and $\mathbf{H}^{n-1}$ with $\mathbf{R}^{4(n-1)}$. If $z \in \mathbf{H}^{n-1}$, then $-[z, z] \equiv |z|^2$ is the square of the Euclidean norm of z in $\mathbf{R}^{4(n-1)}$. For $a_t \in A$ and $n = n(w, z) \in N$ we have

$$(a_t n)_{00} = \cosh t + e^t (w + \tfrac{1}{2} |z|^2).$$

Let $f \in \mathcal{D}(G; \chi_l)$. Applying Lemma 3.3 and Formulas (1.5), we obtain

$$\begin{aligned}
\mathcal{A}f(t) &= \frac{1}{d_l^2} e^{\rho t} \int_N f(a_t n) \, dn \\
&= \frac{1}{d_l^3} e^{\rho t} \int_N \chi_l \left(\frac{(a_t n)_{00}}{|(a_t n)_{00}|} \right) Hf(|(a_t n)_{00}|) \, dn \\
&= \frac{1}{d_l^3} e^{\rho t} \int_{\mathbf{R}^{4(n-1)}} \int_{\mathbf{R}^3} \chi_l \left(\frac{\cosh t + e^t(w + \frac{1}{2}|z|^2)}{|\cosh t + e^t(w + \frac{1}{2}|z|^2)|} \right) \\
&\quad \times Hf(|\cosh t + e^t(w + \frac{1}{2}|z|^2)|) \, dz \, dw \\
&= \frac{1}{d_l^3} e^{\rho t} \int_{\mathbf{R}^{4(n-1)}} \int_{\mathbf{R}^3} C_{2l}^1 \left(\frac{\cosh t + \frac{1}{2}e^t|z|^2}{[(\cosh t + \frac{1}{2}e^t|z|^2)^2 + e^{2t}|w|^2]^{\frac{1}{2}}} \right) \\
&\quad \times Hf([(\cosh t + \frac{1}{2}e^t|z|^2)^2 + e^{2t}|w|^2]^{\frac{1}{2}}) \, dz \, dw \\
&\quad \text{(by Formula (4.40))}
\end{aligned}$$

$$\begin{aligned}
&= \frac{4^{n-1}}{d_l^3} \int_{\mathbf{R}^{4(n-1)}} \int_{\mathbf{R}^3} C_{2l}^1 \left(\frac{\cosh t + |X|^2}{[(\cosh t + |X|^2)^2 + |Y|^2]^{\frac{1}{2}}} \right) \\
&\quad \times Hf([(\cosh t + |X|^2)^2 + |Y|^2]^{\frac{1}{2}}) \, dX \, dY \\
&\quad \text{(by substituting } X = \frac{1}{\sqrt{2}}e^{\frac{t}{2}}z, Y = e^t w)
\end{aligned}$$

$$\begin{aligned}
&= \frac{2^\rho}{d_l^3} \frac{\pi^{2n-1}}{\Gamma(2n-2)} \int_0^\infty \int_0^\infty C_{2l}^1 \left(\frac{\cosh t + r^2}{[(\cosh t + r^2)^2 + s^2]^{\frac{1}{2}}} \right) \\
&\quad \times Hf([(\cosh t + r^2)^2 + s^2]^{\frac{1}{2}}) r^{4n-5} s^2 \, ds \, dr
\end{aligned}$$

(by passing to spherical coordinates in $\mathbf{R}^{4(n-1)}$ and in $\mathbf{R}^3$)

$$= \frac{2^{2n}}{d_l^3} \frac{\pi^{2n-1}}{\Gamma(2n-2)} \int_{\cosh t}^{\infty} \left[\int_0^{\infty} C_{2l}^1 \left(\frac{u}{[u^2 + s^2]^{\frac{1}{2}}} \right) Hf([u^2 + s^2]^{\frac{1}{2}}) s^2 ds \right] \\ \times (u - \cosh t)^{2n-3} du$$

(by setting $u = \cosh t + r^2$)

(6.48)

$$= \frac{2^{2n} \pi^{2n-1}}{d_l^3 \Gamma(2n-2)} \int_{\cosh t}^{\infty} \left[\int_u^{\infty} C_{2l}^1 \left(\frac{u}{x} \right) Hf(x) (x^2 - u^2)^{\frac{1}{2}} x dx \right] (u - \cosh t)^{2n-3} du$$

(by setting $x = [u^2 + s^2]^{\frac{1}{2}}$)

$$= \frac{(2\pi)^{2(n-1)}}{d_l^2 \Gamma(2n-1)} \int_{\cosh t}^{\infty} (\mathcal{G}_{2l} Hf)(u) (u - \cosh t)^{2n-3} du$$

$$= \frac{(2\pi)^{2(n-1)}}{d_l^2} \mathcal{W}_{2n-2} \mathcal{G}_{2l} Hf(\cosh t)$$

$$= \frac{(2\pi)^{2(n-1)}}{d_l^2} (H^{-1} \mathcal{W}_{2n-2} \mathcal{G}_{2l} Hf)(t),$$

i.e.

$$\mathcal{A}_l = \frac{(2\pi)^{2(n-1)}}{d_l^2} (H^{-1} \circ \mathcal{W}_{2n-2} \circ \mathcal{G}_{2l} \circ H).$$

The inversion formula immediately follows from Lemma 6.2.

The restriction to $A \equiv \mathbf{R}$ of $f \in \mathcal{D}(G; \chi_l)$ has its support $\text{supp } f$ contained in $[-R, R]$ if and only if $\text{supp } Hf \subset [1, \cosh R]$. Moreover, if $\text{supp } \varphi \subset [1, \cosh R]$, then $\text{supp } \mathcal{W}_\mu \varphi$, $\text{supp } \mathcal{G}_m \varphi$ and $\text{supp } \mathcal{G}_m^{-1} \varphi$ are also contained in $[1, \cosh R]$. The last statement then follows from the formulas for $\mathcal{A}_l$ and $\mathcal{A}_l^{-1}$. $\square$

The τ_l -spherical transform of $f \in \mathcal{D}(G; \chi_l)$ is the function $\widehat{f}_l$ on $\mathbf{C}$ defined by

$$\widehat{f}_l(s) = \int_G f(g) \zeta_{l,s}(g) dg, \quad s \in \mathbf{C}.$$

Let $S_l: f \mapsto \widehat{f}_l$ denote the τ_l -spherical transform, and let $\mathcal{F}$ denote the Fourier-Laplace transform on $\mathbf{R}$. Formulas (3.20) and (3.22) yield

$$(6.49) \quad S_l = \mathcal{F} \circ \mathcal{A}_l.$$

Let $\mathcal{H}_+^R(\mathbf{R})$ denote the set of even functions h on $\mathbf{C}$ which are entire rapidly decreasing functions of exponential type R : for every integer $N \geq 0$ there is a constant $C_N > 0$ so that

$$|h(s)| \leq C_N (1 + |s|)^{-N} e^{R|\Re s|} \quad \text{for all } s \in \mathbf{C}.$$

Set $\mathcal{H}_+(\mathbf{R}) := \bigcup_{R>0} \mathcal{H}_+^R(\mathbf{R})$. Theorem 6.3 and the Paley-Wiener Theorem for the Fourier-Laplace transform of even functions on $\mathbf{R}$ prove the following theorem.

6.4. THEOREM (Paley-Wiener Theorem). *The τ_l -spherical transform is a bijection of $\mathcal{D}(G; \chi_l)$ onto $\mathcal{H}_+(\mathbf{R})$. Moreover, the restriction of $f \in \mathcal{D}(G; \chi_l)$ to $A \equiv \mathbf{R}$ has support in $[-R, R]$ if and only if $\widehat{f}_l \in \mathcal{H}_+^R(\mathbf{R})$.*

We conclude this section by observing that the τ_l -Abel transform is related, as one should expect, to the Abel transform of [K2], §5.

Reversing the order of integration and substituting $x = \cosh \tau$ and $u = \cosh w$, we obtain from (6.48)

$$(6.50) \quad \mathcal{A}_l f(t) = \int_t^\infty A_l(t, \tau) f(\tau) d\tau$$

where

$$\begin{aligned} A_l(t, \tau) := & \frac{(2\pi)^{2n-1}}{d_l^3 \Gamma(2n-2)} \sinh(2\tau) \int_t^\tau C_{2l}^1 \left(\frac{\cosh w}{\cosh \tau} \right) (\cosh^2 \tau - \cosh^2 w)^{\frac{1}{2}} \\ & \times (\cosh w - \cosh t)^{2n-3} \sinh w dw. \end{aligned}$$

Substituting also $y = \frac{\cosh \tau - \cosh w}{\cosh \tau - \cosh t}$ and setting

$$\gamma(t, \tau) = \frac{\cosh \tau - \cosh t}{2 \cosh \tau} \quad \text{and} \quad K_l = \frac{(2\pi)^{2n-1}}{d_l^3 \Gamma(2n-2)},$$

we get from Formula (6.46)

$$\begin{aligned}
 A_l(t, \tau) &= \sqrt{2} K_l \sinh(2\tau) (\cosh \tau - \cosh t)^{2n-\frac{3}{2}} (\cosh \tau)^{\frac{1}{2}} \\
 &\quad \times \int_0^1 C_{2l}^1 (1 - 2\gamma(t, \tau)y) y^{\frac{1}{2}} (1-y)^{2n-3} (1 - \gamma(t, \tau)y)^{\frac{1}{2}} dy \\
 &= \sqrt{2} (2l+1) K_l \sinh(2\tau) (\cosh \tau - \cosh t)^{2n-\frac{3}{2}} (\cosh \tau)^{\frac{1}{2}} \\
 &\quad \times \int_0^1 F\left(\frac{3}{2} + 2l, -2l - \frac{1}{2}; \frac{3}{2}; \gamma(t, \tau)y\right) y^{\frac{1}{2}} (1-y)^{2n-3} (1 - \gamma(t, \tau)y)^{\frac{1}{2}} dy.
 \end{aligned}$$

If we now apply the relation ([E⁺], 2.9(2))

$$(6.51) \quad F(a, b; c; z) = (1-z)^{c-a-b} F(c-a, c-b; c; z)$$

and Bateman's Formula ([E⁺], 2.4(2))

$$\begin{aligned}
 (6.52) \quad F(a, b; c; z) &= \frac{\Gamma(c)}{\Gamma(s)\Gamma(c-s)} \int_0^1 x^{s-1} (1-x)^{c-s-1} F(a, b; s; xz) dx \\
 &\quad \text{for } \Re c > \Re s > 0, z \neq 1, |\arg(1-z)| < \pi,
 \end{aligned}$$

we finally obtain

$$\begin{aligned}
 (6.53) \quad A_l(t, \tau) &= \frac{(2\pi)^{2n-\frac{1}{2}}}{2\Gamma(2n-\frac{1}{2})} \frac{1}{d_l^2} \sinh(2\tau) (\cosh \tau - \cosh t)^{2n-\frac{3}{2}} (\cosh \tau)^{\frac{1}{2}} \\
 &\quad \times F\left(\frac{3}{2} + 2l, -2l - \frac{1}{2}, 2n - \frac{1}{2}, \gamma(t, \tau)\right).
 \end{aligned}$$

The comparison of Formula (6.53) with the kernel $A_{2n-1, 2l+1}(t, \tau)$ in [K2], Formula (5.60), gives

$$\begin{aligned}
 (6.54) \quad A_l(t, \tau) &= \frac{1}{2} \frac{1}{d_l^2} \left(\frac{\pi}{4}\right)^{2n} \frac{1}{\Gamma(2n)} 2^{-4l} (\cosh \tau)^{-2l} A_{2n-1, 2l+1}(t, \tau) \\
 &= 2^{-4l} C_l (\cosh \tau)^{-2l} A_{2n-1, 2l+1}(t, \tau)
 \end{aligned}$$

where we have set

$$(6.55) \quad C_l := \frac{1}{2} \frac{1}{(2l+1)^2} \left(\frac{\pi}{4}\right)^{2n} \frac{1}{\Gamma(2n)}.$$