

5. Lower bounds

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **45 (1999)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **27.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

5. LOWER BOUNDS

Now we will consider generalized Følner sequences for functions f such that

$$Pf \geq \|P\|f.$$

This will enable us to obtain some lower bounds on the norm of random walk operators on graphs.

As in Section 3, let X be a connected, locally finite graph and let P be the simple random walk operator on X .

In this section we will prove the following lower bound on the norm $\|P\|$:

THEOREM 8. *Let X be a graph such that at each vertex there are at most k edges. Then*

$$\|P\| \geq \frac{2\sqrt{k-1}}{k}.$$

The norm of the random walk operator $\|P\|$ is equal to $\frac{2\sqrt{k-1}}{k}$ for the random walk on the tree which has k edges at each vertex. In [9] Kesten proved this lower bound in the case of Cayley graphs.

Proof of Theorem 8. Let us consider a graph X such that at each vertex there are at most k edges. We can suppose that $k \geq 3$ because for $k = 2$ we obtain subgraphs of $\mathbf{Z}$ or finite graphs, and necessarily $\|P\| = 1$. As it is enough to prove the desired bound for any connected component of X , we can suppose that X is connected.

In order to show that $\|P\|$ is large enough, we will construct a sequence of functions $f_n \in l^2(X, N)$ such that

$$\limsup_{n \rightarrow +\infty} \frac{\|Pf_n\|_{l^2(X, N)}}{\|f_n\|_{l^2(X, N)}} \geq \frac{2\sqrt{k-1}}{k}.$$

Let us endow the set of vertices of X with a metric. The distance between two vertices is the smallest number of edges needed to connect them. Let us choose a vertex e in X and for a vertex v let us denote by $|v|$ its distance from e .

Let f be the unique (up to translations and multiplications) radial eigenfunction of P on the homogeneous tree of degree k , corresponding to the eigenvalue $\frac{2\sqrt{k-1}}{k}$, which is the norm of P on this tree, i.e.

$$(15) \quad f(v) = g(|v|) = \left(\frac{k-2}{k} |v| + 1 \right) \left(\frac{1}{\sqrt{k-1}} \right)^{|v|}.$$

Using (15) we can define f on X . We then prove

LEMMA 6. *For any vertex $v \in X$,*

$$Pf(v) \geq \frac{2\sqrt{k-1}}{k} f(v).$$

Proof. If $v = e$ the result is clearly true. Let us consider then a vertex $v \in X$ such that $n = |v| \geq 1$. Let the number of neighbors of v which are at a distance $n-1$ or n from e be equal respectively to p and q . So the number of neighbors of v which are at a distance $n+1$ is equal to $N(v) - p - q$. Hence

$$Pf(v) = \frac{1}{N(v)} (pg(n-1) + qg(n) + (N(v) - p - q)g(n+1)).$$

As $p \geq 1$ and g is a decreasing function,

$$Pf(v) \geq \frac{1}{N(v)} (g(n-1) + (N(v) - 1)g(n+1)).$$

As $N(v) \leq k$ and $g(n-1) \geq g(n+1)$,

$$Pf(v) \geq \frac{1}{k} (g(n-1) + (k-1)g(n+1)) = \frac{2\sqrt{k-1}}{k} g(n). \quad \square$$

Let us denote by S_n and B_n the vertices which are respectively at a distance n and less than or equal to n .

LEMMA 7.

$$\frac{\sum_{v \in S_{n+1}} f^2(v) N(v)}{\sum_{v \in B_n} f^2(v) N(v)} \rightarrow_{n \rightarrow \infty} 0.$$

Proof. As $1 \leq N(v) \leq k$ it is enough to show that

$$\frac{\sum_{v \in S_{n+1}} f^2(v)}{\sum_{v \in B_n} f^2(v)} \rightarrow_{n \rightarrow \infty} 0.$$

Let us denote

$$a_n = \sum_{v \in S_n} f^2(v) = |S_n| g^2(n).$$

As $|S_{n+1}| \leq (k-1)|S_n|$ one has

$$(16) \quad a_{n+1} = |S_{n+1}|g^2(n+1) \leq (k-1)|S_n|g^2(n+1) = \left(1 + \frac{k-2}{(k-2)n+k}\right)^2 a_n.$$

We have to show that

$$(17) \quad \frac{\sum_{v \in S_{n+1}} f^2(v)}{\sum_{v \in B_n} f^2(v)} = \frac{a_{n+1}}{a_1 + \cdots + a_n} \xrightarrow{n \rightarrow \infty} 0.$$

It is a standard exercise to show that (16) implies (17). $\square$

Let f_n be the sequence of functions which are restrictions of f to the vertices that are at a distance not greater than n :

$$f_n = f|_{B_n}.$$

By Lemma 6 and Lemma 7 it follows that

$$\limsup_{n \rightarrow +\infty} \frac{\|Pf_n\|_{l^2(X,N)}}{\|f_n\|_{l^2(X,N)}} \geq \frac{2\sqrt{k-1}}{k},$$

which proves Theorem 8. $\square$

Some examples of upper bounds on the norm of the simple random walk operator on graphs and their comparison with the lower bound from Theorem 8 can be found in [22].

REFERENCES

- [1] AOMOTO, K. and Y. KATO. Green functions and spectra on free products of cyclic groups. *Ann. Inst. Fourier (Grenoble)* 38, 1 (1988), 59–85.
- [2] BOUGEROL, PH. and L. ÉLIE. Existence of positive harmonic functions on groups and on covering manifolds. *Ann. Inst. H. Poincaré Probab. Statist.* 31, No. 1 (1995), 59–80.
- [3] CARTWRIGHT, D. I. and P. M. SOARDI. Random walks on free products, quotients and amalgams. *Nagoya Math. J.* 102 (1986), 163–180.
- [4] FØLNER, E. On groups with full Banach mean value. *Math. Scand.* 3 (1955), 243–254.
- [5] GOODMAN, F. M., P. DE LA HARPE and V. F. R. JONES. Coxeter graphs and towers of algebras. *Math. Sci. Res. Inst. Publ.* 14. Springer-Verlag (1989).
- [6] GROMOV, M. Groups of polynomial growth and expanding maps. *Inst. Hautes Études Sci. Publ. Math.* 53 (1981), 53–78.
- [7] KAIMANOVICH, V. Dirichlet norms, capacities and generalized isoperimetric inequalities for Markov operators. *Potential Anal.*, 1 (1992), 61–82.