

4.1 Totally geodesic submanifolds

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **47 (2001)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **27.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

$$u = (R^*Ru) * T.$$

Though the question can be tackled by harmonic analysis on X (cf. Section 5), a G -invariant linear differential operator D can sometimes be found directly, such that $DS = \delta$. Then (2) follows from the equality $u = u * DS = D(u * S)$. Indeed, for any test function φ ,

$$\begin{aligned} \langle D(u * S), \varphi \rangle &= \langle u * S, {}^t D\varphi \rangle \\ &= \langle u(g \cdot x_o), \langle S, ({}^t D\varphi) \circ \tau(g) \rangle \rangle \quad \text{by (1)} \\ &= \langle u(g \cdot x_o), \langle S, {}^t D(\varphi \circ \tau(g)) \rangle \rangle, \end{aligned}$$

since the transpose operator ${}^t D$ is G -invariant too, as follows from the existence of a G -invariant measure on X . Finally,

$$\begin{aligned} \langle D(u * S), \varphi \rangle &= \langle u(g \cdot x_o), \langle DS, \varphi \circ \tau(g) \rangle \rangle \\ &= \langle u * DS, \varphi \rangle, \end{aligned}$$

as claimed; assuming G unimodular (as in [9], p. 291) is thus unnecessary here.

The method applies whenever we can find a G -invariant differential operator D on X with given fundamental solution S . We shall now investigate this question on the basis of Propositions 4 and 5.

4. RADON TRANSFORMS ON ISOTROPIC SPACES

Throughout this section X will be an *isotropic* connected *noncompact* Riemannian manifold, that is a Euclidean space or a Riemannian globally symmetric space of rank one:

$$X = \mathbf{R}^n \text{ or } H^m(\mathbf{R}), H^{2m}(\mathbf{C}), H^{4m}(\mathbf{H}), H^{16}(\mathbf{O}),$$

where all superscripts denote the real dimension of these real, complex, quaternionic or Cayley hyperbolic spaces (cf. Wolf [18], §8.12). We first try to invert the d -geodesic Radon transform on X , defined by integrating over a family of d -dimensional totally geodesic submanifolds of X . At the end of this section we shall see that the same tools provide an inversion formula for the horocycle Radon transform on $H^{2k+1}(\mathbf{R})$.

4.1 TOTALLY GEODESIC SUBMANIFOLDS

Our first goal is to describe these submanifolds and the corresponding functions S in Proposition 4.

a. Let $X = G/K$ be a *Riemannian symmetric space of the noncompact type (of arbitrary rank)*, where G is a connected semisimple Lie group and K a maximal compact subgroup (see Notations, **c** and **d**).

At the Lie algebra level, a totally geodesic submanifold of X is defined by a *Lie triple system*, i.e. a vector subspace $\mathfrak{s}$ of $\mathfrak{p}$ such that $[\mathfrak{s}, [\mathfrak{s}, \mathfrak{s}]] \subset \mathfrak{s}$. Then $\text{Exp } \mathfrak{s}$ is totally geodesic in X and contains the origin x_o . Besides $\mathfrak{k}' = [\mathfrak{s}, \mathfrak{s}] \subset \mathfrak{k}$ and $\mathfrak{g}' = \mathfrak{k}' \oplus \mathfrak{s}$ are Lie subalgebras of $\mathfrak{g}$. Let G' be the (closed) Lie subgroup with Lie algebra $\mathfrak{g}'$, and K' (with Lie algebra $\mathfrak{k}'$) be the isotropy subgroup of x_o in G' . Then

$$\text{Exp } \mathfrak{s} = G'/K' = G' \cdot x_o,$$

a closed symmetric subspace of X ([8], p.224–226, or [15], p.234 sq.).

Now let Y be the set of all d -dimensional totally geodesic submanifolds $y = g \cdot y_o$ of X , with $g \in G$ and $y_o = \text{Exp } \mathfrak{s} = G' \cdot x_o$. *Lemma 1* applies: if H is the subgroup of all $h \in G$ such that $h \cdot y_o = y_o$, then $y_o = H \cdot x_o$, $Y = G/H$ and the incidence relation is $x \in y$.

It will be useful to note that the Lie algebra $\mathfrak{h}$ of H satisfies

$$(3) \quad \mathfrak{h} = (\mathfrak{h} \cap \mathfrak{k}) \oplus (\mathfrak{h} \cap \mathfrak{p}), \quad \mathfrak{h} \cap \mathfrak{k} \supset [\mathfrak{s}, \mathfrak{s}], \quad \mathfrak{h} \cap \mathfrak{p} = \mathfrak{s}.$$

Indeed the definition of H shows its invariance under the Cartan involution of G , whence the direct sum decomposition of $\mathfrak{h}$. Besides $\mathfrak{h}$ contains $\mathfrak{g}' = [\mathfrak{s}, \mathfrak{s}] \oplus \mathfrak{s}$ by Lemma 1 and, for $V \in \mathfrak{h} \cap \mathfrak{p}$, the point $\exp V \cdot x_o = \text{Exp } V$ belongs to $H \cdot x_o = \text{Exp } \mathfrak{s}$, thus $V \in \mathfrak{s}$ by the injectivity of Exp on $\mathfrak{p}$.

By Lemma 1 the Radon transform of $u \in C_c(X)$ is given by

$$Ru(y) = \int_y u(x) dm_y(x) = \int_{\text{Exp } \mathfrak{s}} u(g \cdot x) dm_{y_o}(x),$$

where dm_{y_o} is the Riemannian measure induced by X on its submanifold $y_o = \text{Exp } \mathfrak{s}$.

b. RANK ONE CASE. We now restrict to the rank one case (hyperbolic spaces). Let $H \in \mathfrak{s}$ be a fixed non zero vector. The line $\mathfrak{a} = \mathbf{R}H$ is a maximal abelian subspace of $\mathfrak{p}$ and $\mathfrak{s}$, and $\text{Exp } \mathfrak{s}$ is again a symmetric space of rank one. The classical decomposition

$$\mathfrak{p} = \mathfrak{a} \oplus \mathfrak{p}_\alpha \oplus \mathfrak{p}_{2\alpha}$$

into eigenspaces of $(\text{ad } H)^2$, with respective eigenvalues 0 , $(\alpha(H))^2$, $(2\alpha(H))^2$ (where α and 2α are the positive roots of the pair $(\mathfrak{g}, \mathfrak{a})$), implies a similar decomposition of the invariant subspace $\mathfrak{s}$:

$$\mathfrak{s} = \mathfrak{a} \oplus \mathfrak{s}_\alpha \oplus \mathfrak{s}_{2\alpha},$$

with $\mathfrak{s}_\alpha = \mathfrak{s} \cap \mathfrak{p}_\alpha$ and $\mathfrak{s}_{2\alpha} = \mathfrak{s} \cap \mathfrak{p}_{2\alpha}$. We set

$$\begin{aligned} p &= \dim \mathfrak{p}_\alpha, & q &= \dim \mathfrak{p}_{2\alpha}, & n &= \dim X = p + q + 1, \\ p' &= \dim \mathfrak{s}_\alpha, & q' &= \dim \mathfrak{s}_{2\alpha}, & d &= \dim \mathfrak{s} = p' + q' + 1, \end{aligned}$$

with $q = q' = 0$ when 2α is not a root (case of real hyperbolic spaces).

Let us normalize the vector H by the condition $\alpha(H) = 1$. Multiplying if necessary the Riemannian metric of X by a constant factor, we may assume that the corresponding Euclidean norm on $\mathfrak{p}$ satisfies $\|H\| = 1$. Since Exp is a diffeomorphism of $\mathfrak{p}$ onto X , the integral of a function $u \in C_c(X)$ can be computed as

$$\int_X u(x) dx = \int_{\mathfrak{p}} u(\text{Exp } Z) J(Z) dZ,$$

where $J(Z) = \det_{\mathfrak{p}}(\sinh \text{ad } Z / \text{ad } Z)$ is the Jacobian of Exp , a K -invariant function on $\mathfrak{p}$. If u is K -invariant on X , we simply write $u(r)$ for $u(\text{Exp } Z) = u(\text{Exp } rH)$ with $r = \|Z\|$ whence, computing with spherical coordinates on $\mathfrak{p}$,

$$\int_X u(x) dx = \int_0^\infty u(r) A(r) dr,$$

where $A(r) = \omega_n r^{n-1} \det_{\mathfrak{p}}(\sinh \text{ad } rH / \text{ad } rH)$ is the area of the sphere with center x_o and radius r in X , and $\omega_n = 2\pi^{n/2} / \Gamma(n/2)$ is the area of the unit sphere in $\mathbf{R}^n$. Taking account of the eigenvalues of $(\text{ad } H)^2$ we obtain, with a parameter ε explained in the next remark,

$$(4) \quad A(r) = \omega_n \left(\frac{\sinh \varepsilon r}{\varepsilon} \right)^p \left(\frac{\sinh 2\varepsilon r}{2\varepsilon} \right)^q = \omega_n \left(\frac{\sinh \varepsilon r}{\varepsilon} \right)^{n-1} (\cosh \varepsilon r)^q.$$

A similar expression gives $A_o(r)$ for the submanifold y_o (with d, p', q' instead of n, p, q). The distribution S in Proposition 4 is thus defined by the radial function

$$(5) \quad S(r) = A_o(r)/A(r) = (\omega_d/\omega_n) \left(\frac{\sinh \varepsilon r}{\varepsilon} \right)^{d-n} (\cosh \varepsilon r)^{q'-q}.$$

REMARK. Here $\varepsilon = 1$ for spaces of the noncompact type, but (4) and (5) remain valid in the Euclidean case too, setting $\varepsilon = 0$ and $(\sinh \varepsilon r)/\varepsilon = r$: when $X = \mathbf{R}^n$ the geodesic submanifolds are the affine d -planes, $1 \leq d \leq n-1$, and

$$S(r) = (\omega_d/\omega_n) r^{d-n}.$$

The compact cases (projective spaces) might be dealt with similarly. One should then normalize H by $\alpha(H) = i$ and replace ε by i . Integrals with respect to r should run from 0 to the diameter ℓ of X , i.e. the first number $\ell > 0$ such that $A(\ell) = 0$.

4.2 AN INVERSION FORMULA

The G -invariant differential operators on an isotropic space X are the polynomials of its Laplace-Beltrami operator L ([9], p. 288). In order to invert the d -geodesic Radon transform on X , Section 3.2 suggests looking for a polynomial P such that the above distribution S is a fundamental solution of $P(L)$.

Motivated by (4) and (5), we introduce the family of radial functions $f_{a,b}$ on X defined by

$$f_{a,b}(r) = \left(\frac{\sinh \varepsilon r}{\varepsilon} \right)^a (\cosh \varepsilon r)^b = \left(\frac{\sinh \varepsilon r}{\varepsilon} \right)^{a-b} \left(\frac{\sinh 2\varepsilon r}{2\varepsilon} \right)^b,$$

where a and b are real constants and r is the distance from the origin x_o ; in particular $f_{a,b}(r) = r^a$ for $\varepsilon = 0$. Thus

$$A(r) = \omega_n f_{n-1,q}, \quad S(r) = (\omega_d/\omega_n) f_{d-n,q'-q}$$

with q, q', n and d as defined above.

PROPOSITION 6. Assume $\varepsilon = 0$ (Euclidean case), or $\varepsilon = 1$ and $b = 0$, or else $\varepsilon = 1$ and $b = 1 - q$ (hyperbolic cases). Then, for any integer $k \geq 1$, the function $f_{2k-n,b}$ defines a K -invariant distribution $F_{2k-n,b}$ on X such that

$$P_k(L)F_{2k-n,b} = \omega_n 2^{k-1} (k-1)! (2-n)(4-n) \cdots (2k-n) \delta,$$

where δ is the Dirac distribution at the origin x_o and P_k is the polynomial

$$P_k(x) = \prod_{j=1}^k (x + \varepsilon^2(n - 2j - b)(2j + b + q - 1)).$$

REMARK. The case $b = 0, n = 2k + 2$ was given by Schimming and Schlichtkrull [17], Theorem 6.1, as an example in their beautiful study of Hadamard's method and Helmholtz operators on harmonic manifolds.