

2.2 Symplectic characteristic classes and Chern classes

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **47 (2001)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **28.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

$\phi(\tilde{\rho}(z))$ of any generator $z \in \mathbf{Z}/p\mathbf{Z}$ is conjugate to a $Y \in \mathrm{Sp}(2n, \mathbf{Z})$. Then $d_j(\rho) = \tilde{\rho}^*(c_j) = c_j(\rho)$. We define the total Chern class of a representation $\tilde{\rho}$ to be

$$c(\tilde{\rho}) := 1 + c_1(\tilde{\rho}) + c_2(\tilde{\rho}) + \cdots + c_n(\tilde{\rho}).$$

It has the well-known properties $c(\rho \oplus \sigma) = c(\rho)c(\sigma)$, $c(m\rho) = c(\rho)^m$, where ρ, σ are representations and m is a positive integer.

2.2 SYMPLECTIC CHARACTERISTIC CLASSES AND CHERN CLASSES

THEOREM 2.1. *Let p be an odd prime. Then for any $n = 1, \dots, (p-1)/2$ there exists a representation $\tilde{\rho}: \mathbf{Z}/p\mathbf{Z} \rightarrow \mathrm{U}((p-1)/2)$ such that the n -th Chern class $c_n(\tilde{\rho})$ is nonzero and the representation $\phi \circ \tilde{\rho}: \mathbf{Z}/p\mathbf{Z} \rightarrow \mathrm{Sp}(p-1, \mathbf{R})$ factors, up to conjugation, through a representation $\rho: \mathbf{Z}/p\mathbf{Z} \rightarrow \mathrm{Sp}(p-1, \mathbf{Z})$.*

The representation $\tilde{\rho}$ factors through $\mathrm{Sp}(p-1, \mathbf{Z})$ if the image $\tilde{\rho}(z)$ of a generator $z \in \mathbf{Z}/p\mathbf{Z}$ satisfies the condition stated in Theorem 1.2. Then, because $c_n(\tilde{\rho}) \neq 0$, we have $d_n(\rho) \neq 0$ where $\rho: \mathbf{Z}/p\mathbf{Z} \rightarrow \mathrm{Sp}(p-1, \mathbf{Z})$ is the representation corresponding to $\tilde{\rho}$.

Proof of Theorem 2.1. Let $\mathcal{U}$ be the set of subsets $\mathcal{I} \subset (\mathbf{Z}/p\mathbf{Z})^*$ of cardinality $|\mathcal{I}| = (p-1)/2$, and $j \in \mathcal{I}$ implies $p-j \notin \mathcal{I}$. The cardinality of $\mathcal{U}$ is $2^{(p-1)/2}$. We always assume the elements $j \in \mathcal{I}$ to be represented by integers j with $1 \leq j < p$. Note that we will use the same notation for the elements of $\mathcal{I}$ and their representatives. For $j = 1, \dots, p-1$ let $\tilde{\rho}_j: \mathbf{Z}/p\mathbf{Z} \rightarrow \mathrm{U}(1)$ be the one-dimensional representation with $\tilde{\rho}_j(z) := e^{j2\pi i/p}$ for a fixed generator $z \in \mathbf{Z}/p\mathbf{Z}$. For a given $\mathcal{I}$ we define $\tilde{\rho}_{\mathcal{I}}$ to be the direct sum of the representations $\tilde{\rho}_j$, $j \in \mathcal{I}$. Let $x := c_1(\tilde{\rho}_1)$, then the total Chern class of $\tilde{\rho}_{\mathcal{I}}$ is

$$c(\tilde{\rho}_{\mathcal{I}}) = c\left(\bigoplus_{j \in \mathcal{I}} \tilde{\rho}_j\right) = \prod_{j \in \mathcal{I}} (1 + jx).$$

The representations $\tilde{\rho}_{\mathcal{I}}$ are those which factor through $\mathrm{Sp}(p-1, \mathbf{Z})$. For a given $\mathcal{I} \in \mathcal{U}$ we define $-\mathcal{I} := \{p-j \mid j \in \mathcal{I}\}$. Then $-\mathcal{I} \in \mathcal{U}$ and $\mathcal{I} \cup -\mathcal{I} = (\mathbf{Z}/p\mathbf{Z})^*$. Moreover, we get $c(\tilde{\rho}_{\mathcal{I}})c(\tilde{\rho}_{-\mathcal{I}}) = 1 - x^{p-1}$. The n -th Chern class $c_n(\tilde{\rho}_{\mathcal{I}})$ is nonzero if and only if the coefficient a_n of x^n in the total Chern class $c(\tilde{\rho}_{\mathcal{I}})$ is nonzero. Let $\mathcal{I} := \{j_1, \dots, j_{(p-1)/2}\} \in \mathcal{U}$; then we define

$$\mathcal{I}_l := \{j_1, \dots, j_{l-1}, -j_l, j_{l+1}, \dots, j_{(p-1)/2}\} \in \mathcal{U}.$$

We assume that $1 \leq n \leq (p-1)/2$ exists such that for each set $\mathcal{I} \in \mathcal{U}$ the coefficient a_n of x^n in $c(\tilde{\rho}_{\mathcal{I}})$ is zero. It is impossible that $n = (p-1)/2$

because $a_{(p-1)/2}$ is the product of the $j \in \mathcal{I}$ and therefore nonzero. Now let $n \neq 0$, $n \neq (p-1)/2$; then we define for any $l = 1, \dots, (p-1)/2$

$$b_n^l := \sum_{\substack{J \subseteq \mathcal{I} \setminus \{j_l\} \\ |J|=n}} \prod_{j \in J} j, \quad b_0^l := 1.$$

Then the coefficient of x^n in $c(\tilde{\rho}_{\mathcal{I}})$ is $a_n = b_n^l + j_l b_{n-1}^l$. Because of our assumption, the coefficients of x^n in $c(\tilde{\rho}_{\mathcal{I}})$ and in $c(\tilde{\rho}_{\mathcal{I}_l})$ are $b_n^l + j_l b_{n-1}^l = 0$ and $b_n^l - j_l b_{n-1}^l = 0$ respectively. This implies that $b_n^l = 0$, $b_{n-1}^l = 0$ and

$$\begin{aligned} a_{n+1} &= \sum_{\substack{J \subseteq \mathcal{I} \\ |J|=n+1}} \prod_{j \in J} j \\ &= \frac{1}{n+1} \sum_{j_l \in \mathcal{I}} \left(j_l \sum_{\substack{J \subseteq \mathcal{I} \setminus \{j_l\} \\ |J|=n}} \prod_{j \in J} j \right) = \frac{1}{n+1} \sum_{j_l \in \mathcal{I}} j_l b_n^l = 0. \end{aligned}$$

The factor $1/(n+1)$ appears because in the second line we have $n+1$ times each term appearing in the sum of the first line. Therefore $a_{n+1} = 0$ for each set $\mathcal{I} \in \mathcal{U}$, and by induction we get $a_{(p-1)/2} = 0$ for each set $\mathcal{I} \in \mathcal{U}$, which is impossible. $\square$

Let $\mathrm{Sp}(\mathbf{Z}) := \bigcup_{n \geq 1} \mathrm{Sp}(2n, \mathbf{Z})$.

THEOREM 2.2. *For every $j \geq 1$, $d_j(\mathbf{Z}) \in H^{2j}(\mathrm{Sp}(\mathbf{Z}), \mathbf{Z})$ has infinite order.*

Proof. This theorem is a corollary of Theorem 2.1. A consequence of the stability result stated in section 2.1 is that for $p-1 > 8j+8$ the inclusion

$$\mathrm{Sp}(p-1, \mathbf{Z}) \longrightarrow \mathrm{Sp}(\mathbf{Z})$$

induces an isomorphism

$$H^{2j}(\mathrm{Sp}(\mathbf{Z}), \mathbf{Z}) \xrightarrow{\cong} H^{2j}(\mathrm{Sp}(p-1, \mathbf{Z}), \mathbf{Z}).$$

In Theorem 2.1 we have shown that for any odd prime p and any integer $j = 1, \dots, (p-1)/2$ a representation $\tilde{\rho}_{\mathcal{I}}: \mathbf{Z}/p\mathbf{Z} \rightarrow \mathrm{U}((p-1)/2)$ exists that factors through $\mathrm{Sp}(p-1, \mathbf{Z})$ and for which the j -th Chern class $c_j(\tilde{\rho}_{\mathcal{I}})$ is nonzero. Then the j -th symplectic class $d_j(\rho_{\mathcal{I}})$ is also nonzero. Here the

representation $\rho_I: \mathbf{Z}/p\mathbf{Z} \rightarrow \mathrm{Sp}(p-1, \mathbf{Z})$ is the one corresponding to $\tilde{\rho}_I$. We have an induced homomorphism

$$\begin{aligned} \rho_I^*: H^{2j}(\mathrm{Sp}(p-1, \mathbf{Z}), \mathbf{Z}) &\longrightarrow H^{2j}(\mathbf{Z}/p\mathbf{Z}, \mathbf{Z}) \\ d_j(\mathbf{Z}) &\longmapsto d_j(\rho_I). \end{aligned}$$

Herewith for any p the class $d_j(\mathbf{Z}) \in H^{2j}(\mathrm{Sp}(p-1, \mathbf{Z}), \mathbf{Z})$ is nonzero and has either infinite order or finite order divisible by p , since it restricts non-trivially to $H^{2j}(\mathbf{Z}/p\mathbf{Z}, \mathbf{Z})$. This shows that $d_j(\mathbf{Z}) \in H^{2j}(\mathrm{Sp}(\mathbf{Z}), \mathbf{Z})$ has infinite order. $\square$

This is a new proof of a result of A. Borel [3]. He proved that $H^*(\mathrm{Sp}(\mathbf{Z}), \mathbf{Q}) = \mathbf{Q}[d_1, d_3, \dots]$. Moreover, each d_{2i} can be expressed as a polynomial in the d_{2j+1} 's. This implies that all the $d_i(\mathbf{Z})$'s have infinite order.

REFERENCES

- [1] ALEXANDER, J. P., P. E. CONNER and G. C. HAMRICK. *Odd order group actions and Witt classifications of innerproducts*. Lecture Notes in Mathematics 625. Springer, 1977.
- [2] ALEXANDER, J. P., P. E. CONNER, G. C. HAMRICK and J. W. VICK. Witt classes of integral representations of an abelian p -group. *Bull. Amer. Math. Soc.* 80 (1974), 1179–1182.
- [3] BOREL, A. Stable real cohomology of arithmetic groups. *Ann. Sci. École Norm. Sup.* (4) 7 (1974), 235–272.
- [4] BROWN, K. S. Euler characteristics of discrete groups and G -spaces. *Invent. Math.* 27 (1974), 229–264.
- [5] BÜRGISSER, B. Elements of finite order in symplectic groups. *Arch. Math. (Basel)* 39 (1982), 501–509.
- [6] BUSCH, C. Symplectic characteristic classes and the Farrell cohomology of $\mathrm{Sp}(p-1, \mathbf{Z})$. Dissertation ETH no. 13506, ETH Zurich (2000).
- [7] CHARNEY, R. A generalization of a theorem of Vogtmann. *J. Pure Appl. Algebra* 44 (1987), 107–125.
- [8] GARBANATI, D. A. Unit signatures, and even class numbers, and relative class numbers. *J. reine angew. Math.* 274–275 (1975), 376–384.
- [9] NEUKIRCH, J. *Algebraische Zahlentheorie*. Springer, 1992.
- [10] ———. *Class Field Theory*. Grundlehren der mathematischen Wissenschaften 280. Springer, 1986.
- [11] SJERVE, D. and Q. YANG. Conjugacy classes of p -torsion in $\mathrm{Sp}_{p-1}(\mathbf{Z})$. *J. Algebra* 195 (1997), 580–603.