

8. The Whitehead product and the positive cone

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **47 (2001)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **27.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

8. THE WHITEHEAD PRODUCT AND THE POSITIVE CONE

We will establish an interesting connection between the positive cone of a product $S^{2n} \times S^{2m}$ and the Whitehead product structure on the homotopy groups of the spaces $BU(k)$. As an application we will get some precise information on the positive cone of $S^{2n} \times S^{2m}$.

Let us first recall the basic properties of the Whitehead product (the reader may refer to [White]). The product $S^p \times S^q$ has a cell structure obtained by attaching a $(p+q)$ -cell to $S^p \vee S^q$. More precisely, there exists a suitable pointed map $f_0: S^{p+q-1} \longrightarrow S^p \vee S^q$ such that $S^p \times S^q$ is homeomorphic to the mapping cone of f_0 :

$$S^p \times S^q \cong C_{f_0} = (S^p \vee S^q) \cup_{f_0} e^{p+q}$$

Given a pointed map $g = \alpha \vee \beta: S^p \vee S^q \longrightarrow X$, where X is a CW-complex, there exists (up to homotopy) an extension $\bar{g}: S^p \times S^q \longrightarrow X$ of g if and only if the composition $g \circ f_0$ is homotopically trivial. Now, considering α and β as elements of the homotopy groups $\pi_p(X)$ and $\pi_q(X)$ respectively, the composition $(\alpha \vee \beta) \circ f_0$ determines an element in the homotopy group $\pi_{p+q-1}(X)$. This defines a map

$$\pi_p(X) \times \pi_q(X) \longrightarrow \pi_{p+q-1}(X), \quad (\alpha, \beta) \longmapsto [\alpha, \beta] := (\alpha \vee \beta) \circ f_0,$$

which by definition is the Whitehead product. One can show that it is $\mathbb{Z}$ -bilinear (provided that $p, q \geq 2$), i.e.

$$[\alpha_1 + \alpha_2, \beta] = [\alpha_1, \beta] + [\alpha_2, \beta] \quad \text{and} \quad [\alpha, \beta_1 + \beta_2] = [\alpha, \beta_1] + [\alpha, \beta_2].$$

Moreover, the Whitehead product is natural with respect to pointed maps, i.e. if $f: X \longrightarrow Y$ is a pointed map between CW-complexes, then

$$[f_*(\alpha), f_*(\beta)] = f_*([\alpha, \beta]).$$

We now want to study the case where $X = BU(l)$. Let x_1 and x_2 be two generators of $\tilde{K}(S^{2n})$ and $\tilde{K}(S^{2m})$ respectively, and assume $1 \leq n \leq m$. By Theorem 4.1, we know that $\text{g-dim}(x_1) = n$ and that $\text{g-dim}(x_2) = m$. Letting $q \geq m$, we consider x_1 and x_2 as maps from S^{2n} (respectively S^{2m}) to BU that lift to $BU(q)$. The element $x_1 + x_2$ of $\tilde{K}(S^{2n} \vee S^{2m}) = \tilde{K}(S^{2n}) \oplus \tilde{K}(S^{2m})$ can be represented by the map $x_1 \vee x_2: S^{2n} \vee S^{2m} \longrightarrow BU$, and it also lifts to a map $z: S^{2n} \vee S^{2m} \longrightarrow BU(q)$.

CLAIM. For $k \in \{m, m+1, \dots, m+n-1\}$, there is no extension of $z = x_1 \vee x_2: S^{2n} \vee S^{2m} \longrightarrow BU(k)$ to a map $S^{2n} \times S^{2m} \longrightarrow BU(k)$.

Let $y: S^{2n} \times S^{2m} \longrightarrow BU(s)$ be an extension of z for some $s \geq m$. Let x be the composition of y with the map $i_s: BU(s) \longrightarrow BU$. This means that $\text{g-dim}(x) \leq s$ and that $\iota^*(x) = x_1 + x_2 \in \tilde{K}(S^{2n} \vee S^{2m})$, where ι is the inclusion of $S^{2n} \vee S^{2m}$ in the product $S^{2n} \times S^{2m}$. Recall that $(\iota^*)^{-1}(x_1 + x_2) = x_1 + x_2 + \mathbf{Z} \cdot x_1 x_2 \subset \tilde{K}(S^{2n} \times S^{2m})$. So, there exists an integer l such that $x = x_1 + x_2 + lx_1 x_2$, and consequently

$$\gamma^{n+m}(x) = (-1)^{n+m-1} (l(n+m-1)! - (n-1)!(m-1)!) \cdot x_1 x_2 \neq 0.$$

We see that $s \geq \text{g-dim}(x) \geq \gamma\text{-dim}(x) \geq n+m$. This proves the claim.

As a direct consequence, by considering x_1 and x_2 as elements (in fact generators) of $\pi_{2n}(BU(k))$ and $\pi_{2m}(BU(k))$ respectively, we get the following result on the Whitehead product:

$$[x_1, x_2] \neq 0 \text{ in } \pi_{2n+2m-1}(BU(k)), \text{ for } m \leq k < n+m.$$

We would now like to get some information on the order of $[x_1, x_2]$ in the homotopy group $\pi_{2n+2m-1}(BU(k))$. By $\mathbf{Z}$ -bilinearity of the Whitehead product, we have $ab[x_1, x_2] = [ax_1, bx_2]$ for any integers a and b . Replacing x_1 by ax_1 and x_2 by bx_2 in the preceding computation (in particular $x = ax_1 + bx_2 + lx_1 x_2$ for some l), one easily verifies that

$$(*) \quad \left. \begin{array}{l} ab[x_1, x_2] = 0 \\ \text{in } \pi_{2n+2m-1}(BU(k)) \\ \text{for } m \leq k < n+m \end{array} \right\} \implies l(n+m-1)! - ab(n-1)!(m-1)! = 0$$

and this implies that ab is a multiple of $(n+m-1)!/((n-1)!(m-1)!)$. Notice that $[x_1, x_2] \in \pi_{2n+2m-1}(BU(k))$ has to be a torsion element. Indeed, by Lemma 4.2, the group $\pi_{2n+2m-1}(BU(m))$ is finite, and the result follows from naturality of the Whitehead product. (In fact, one can show that any group $\pi_{2i+1}(BU(j))$ is finite; this is proved like Lemma 4.2, by appealing to a result of Borel and Hirzebruch: see Remark i) in Section 9.) We have thus obtained the following theorem.

THEOREM 8.1. *Let $1 \leq n \leq m$ and $m \leq k < n+m$. Let x_1 and x_2 be generators of the homotopy groups $\pi_{2n}(BU(k)) \cong \mathbf{Z}$ and $\pi_{2m}(BU(k)) \cong \mathbf{Z}$ respectively. Then the Whitehead product $[x_1, x_2] \in \pi_{2n+2m-1}(BU(k))$ is non-zero. Moreover, its order is a multiple of $\frac{(n+m-1)!}{(n-1)!(m-1)!}$.*

By the implication (*), if $ab[x_1, x_2] = 0$ in $\pi_{2n+2m-1}(BU(k))$ for some k with $m \leq k < n + m$, then for $l = ab(n-1)!(m-1)!/(n+m-1)!$, the geometric dimension of $x := ax_1 + bx_2 + lx_1x_2$ is $\leq k$ (and for any other value of l , $\text{g-dim}(x)$ is $m + n$, provided that $ab \neq 0$). Surprisingly, this condition only depends on l and on the product ab . Consequently, from Theorem 2.3 together with Theorem 7.1, we obtain the following result.

THEOREM 8.2. *The geometric dimension on $\tilde{K}(S^{2n} \times S^{2m})$, with $n \leq m$, is given as follows: for $x = ax_1 + bx_2 + lx_1x_2 \in \tilde{K}(S^{2n} \times S^{2m})$,*

$$\text{g-dim}(x) = \begin{cases} 0 & \text{if } a = b = l = 0 \\ n & \text{if } a \neq 0, b = l = 0 \\ m & \text{if } a = 0, b \neq 0, l = 0 \\ s(ab) & \text{if } b \neq 0, l = ab(n-1)!(m-1)!/(n+m-1)! \\ n+m & \text{if } l \neq ab(n-1)!(m-1)!/(n+m-1)! \end{cases}$$

where $s(ab) \in \{m, m+1, \dots, n+m-1\}$ only depends on the product ab (for fixed n and m).

As a direct consequence of Theorems 8.1 and 8.2, we have

COROLLARY 8.3. *The order of the Whitehead product $[x_1, x_2]$ in $\pi_{2n+2m-1}(BU(n+m-1))$ is exactly $(n+m-1)!/((n-1)!(m-1)!)$.*

REMARK 8.4.

i) This result has been established only using information on the γ -cone of $S^{2n} \times S^{2m}$ (and Serre's theorem on the rational homotopy of spheres). If one is able to compute its positive cone, one then can easily compute the exact order of $[x_1, x_2]$ in the various homotopy groups $\pi_{2n+2m-1}(BU(k))$, for $m \leq k < n + m$: it is given by

$$\min \left\{ l \geq 1 \mid \text{g-dim} \left(l \frac{(n+m-1)!}{(n-1)!(m-1)!} x_1 + x_2 + lx_1x_2 \right) \leq k \right\}.$$

ii) In 1960, Bott [Bott3] has proved Corollary 8.3 by different methods.