

1.2 Elementary properties of $\mathbb{Q}_m$

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **49 (2003)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **29.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

EXAMPLE 1.2. The group $W = \mathbf{Z}/2$ acts on $\mathfrak{h} = \mathbf{C}$ by $s(v) = -v$. In this case m is a non negative integer and $\Sigma = \{s\}$. So this definition says that q is in Q_m iff $q(x) - q(-x)$ is divisible by x^{2m+1} . It is very easy to write a basis of Q_m . It is given by the polynomials $\{x^{2i} \mid i \geq 0\} \cup \{x^{2i+1} \mid i \geq m\}$.

1.2 ELEMENTARY PROPERTIES OF Q_m

Some elementary properties of Q_m are collected in the following proposition.

PROPOSITION 1.3 (see [FV] and references therein).

- 1) $\mathbf{C}[\mathfrak{h}]^W \subset Q_m \subseteq \mathbf{C}[\mathfrak{h}]$, $Q_0 = \mathbf{C}[\mathfrak{h}]$, $Q_m \subset Q_{m'}$ if $m \geq m'$, $\bigcap_m Q_m = \mathbf{C}[\mathfrak{h}]^W$.
- 2) Q_m is a graded subalgebra of $\mathbf{C}[\mathfrak{h}]$.
- 3) The fraction field of Q_m is equal to $\mathbf{C}(\mathfrak{h})$.
- 4) Q_m is a finite $\mathbf{C}[\mathfrak{h}]^W$ -module and a finitely generated algebra. $\mathbf{C}[\mathfrak{h}]$ is a finite Q_m -module.

Proof. 1) is immediate and has already been mentioned in 1.1.

2) Clearly Q_m is closed under addition. Let $p, q \in Q_m$. Let $s \in \Sigma$. Then

$$p(x)q(x) - p(sx)q(sx) = (p(x) - p(sx))q(x) + p(sx)(q(x) - q(sx)).$$

Since both $p(x) - p(sx)$ and $q(x) - q(sx)$ are divisible by $\alpha_s^{2m_s+1}$, we deduce that $p(x)q(x) - p(sx)q(sx)$ is also divisible by $\alpha_s^{2m_s+1}$, proving the claim.

3) Consider the polynomial

$$\delta_{2m+1}(x) = \prod_{s \in \Sigma} \alpha_s(x)^{2m_s+1}.$$

This polynomial is uniquely defined up to scaling. One has $\delta_{2m+1}(sx) = -\delta_{2m+1}(x)$ for each $s \in \Sigma$, hence $\delta_{2m+1} \in Q_m$. Take $f(x) \in \mathbf{C}[\mathfrak{h}]$. We claim that $f(x)\delta_{2m+1}(x) \in Q_m$. As a matter of fact,

$$f(x)\delta_{2m+1}(x) - f(sx)\delta_{2m+1}(sx) = (f(x) + f(sx))\delta_{2m+1}(x),$$

and by its definition $\delta_{2m+1}(x)$ is divisible by $\alpha_s(x)^{2m_s+1}$ for all $s \in \Sigma$. This implies 3).

4) By Hilbert's theorem on the finiteness of invariants, we get that $\mathbf{C}[\mathfrak{h}]^W$ is a finitely generated algebra over $\mathbf{C}$ and $\mathbf{C}[\mathfrak{h}]$ is a finite $\mathbf{C}[\mathfrak{h}]^W$ -module and hence a finite Q_m -module, proving the second part of 4).

Now $Q_m \subset \mathbf{C}[\mathfrak{h}]$ is a submodule of the finite module $\mathbf{C}[\mathfrak{h}]$ over the Noetherian ring $\mathbf{C}[\mathfrak{h}]^W$. Hence it is finite. This immediately implies that Q_m is a finitely generated algebra over $\mathbf{C}$. $\square$

REMARK. In fact, since W is a finite Coxeter group, a celebrated result of Chevalley says that the algebra $\mathbf{C}[\mathfrak{h}]^W$ is not only a finitely generated $\mathbf{C}$ -algebra but actually a free (=polynomial) algebra. Namely, it is of the form $\mathbf{C}[q_1, \dots, q_n]$, where the q_i are homogeneous polynomials of some degrees d_i . Furthermore, if we denote by H the subspace of $\mathbf{C}[\mathfrak{h}]$ of harmonic polynomials, i.e. of polynomials killed by W -invariant differential operators with constant coefficients without constant term, then the multiplication map

$$\mathbf{C}[\mathfrak{h}]^W \otimes H \rightarrow \mathbf{C}[\mathfrak{h}]$$

is an isomorphism of $\mathbf{C}[\mathfrak{h}]^W$ - and of W -modules. In particular, $\mathbf{C}[\mathfrak{h}]$ is a free $\mathbf{C}[\mathfrak{h}]^W$ -module of rank $|W|$.

1.3 THE VARIETY X_m AND ITS BIJECTIVE NORMALIZATION

Using Proposition 1.3, we can define the irreducible affine variety $X_m = \text{Spec}(Q_m)$. The inclusion $Q_m \subset \mathbf{C}[\mathfrak{h}]$ induces a morphism

$$\pi: \mathfrak{h} \rightarrow X_m,$$

which again by Proposition 1.3 is birational and surjective. (Notice that in particular this implies that X_m is singular for all $m \neq 0$.)

In fact, not only is π birational, but a stronger result is true.

PROPOSITION 1.4 (Berest, see [BEG]). *π is a bijection.*

Proof. By the above remarks, we only have to show that π is injective. In order to achieve this, we need to prove that quasi-invariants separate points of $\mathfrak{h}$, i.e. that if $z, y \in \mathfrak{h}$ and $z \neq y$, then there exists $p \in Q_m$ such that $p(z) \neq p(y)$. This is obtained in the following way. Let $W_z \subset W$ be the stabilizer of z and choose $f \in \mathbf{C}[\mathfrak{h}]$ such that $f(z) \neq 0$, $f(y) = 0$. Set

$$p(x) = \prod_{s \in \Sigma, sz \neq z} \alpha_s(x)^{2m_s+1} \prod_{w \in W_z} f(wx).$$

We claim that $p(x) \in Q_m$. Indeed, let $s \in \Sigma$ and assume that $s(z) \neq z$.

We have by definition $p(x) = \alpha_s(x)^{2m_s+1} \tilde{p}(x)$, with $\tilde{p}(x)$ a polynomial. So

$$p(x) - p(sx) = \alpha_s(x)^{2m_s+1} \tilde{p}(x) - \alpha_s(sx)^{2m_s+1} \tilde{p}(sx) = \alpha_s(x)^{2m_s+1} (\tilde{p}(x) + \tilde{p}(sx)).$$

If on the other hand, $sz = z$, i.e. $s \in W_z$, then s preserves the set $W \setminus W_z$, and hence preserves $\prod_{s \in \Sigma \cap (W \setminus W_z)} \alpha_s(x)^{2m_s+1}$ (as it acts by -1 on the products $\prod_{s \in \Sigma} \alpha_s(x)^{2m_s+1}$ and $\prod_{s \in \Sigma \cap W_z} \alpha_s(x)^{2m_s+1}$). Since $\prod_{w \in W_z} f(wx)$ is