

1.3. Operations on analytic spaces.

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **14 (1968)**

Heft 1: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **24.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

for some coherent sheaf \mathcal{I} of ideals of \mathcal{O}_X . An *open analytic subspace* of (X, \mathcal{O}_X) is just a restriction $(U, \mathcal{O}_X \mid U)$, U open in X . An *analytic subspace* of an analytic space (X, \mathcal{O}_X) is a closed analytic subspace (Y, \mathcal{O}_Y) of the open analytic subspace $(\mathbb{C} \bar{Y} \cup Y, \mathcal{O}_{\mathbb{C} \bar{Y} \cup Y})$ of (X, \mathcal{O}_X) , provided $\mathbb{C} \bar{Y} \cup Y$ is indeed open in X , i.e. Y is locally closed in X .

Examples. The “single point” $(0, \mathbb{C})$ is an analytic subspace of the “double point” $(0, \mathbb{C} \{x\}/(x^2))$, but not conversely. The double point is, however, a closed analytic subspace of, e.g., $(\mathbb{C}, \mathcal{O}_{\mathbb{C}})$. A “point” of an analytic space will always mean a single point embedded in (X, \mathcal{O}_X) by means of a map $(0, \mathbb{C}) \rightarrow (X, \mathcal{O}_X)$.

1.3. Operations on analytic spaces.

In this section we shall write X for the analytic space (X, \mathcal{O}_X) .

a) *Product.* By a general definition in the theory of categories, a product of two analytic spaces X, X' is a triple (Z, π, π') where Z is an analytic space and $\pi : Z \rightarrow X, \pi' : Z \rightarrow X'$ are two morphisms with the following property:

Given any analytic space Y and any pair $f : Y \rightarrow X, f' : Y \rightarrow X'$ of morphisms there exists a unique morphism $g : Y \rightarrow Z$ such that $f = \pi \circ g, f' = \pi' \circ g$.

For example, the product of \mathbb{C}^p and \mathbb{C}^q is \mathbb{C}^{p+q} , according to proposition 1.2.4.

We shall see that a product of analytic spaces always exists. The uniqueness of g clearly implies the uniqueness of the product (Z, π, π') up to isomorphism; we denote one such Z by $X \times X'$.

To prove that the product always exists, let us suppose first that X and X' are special models, i.e. X is defined by a triple (U, f, F) where U is open in \mathbb{C}^n, F is a finite-dimensional complex linear space, and $f : U \rightarrow F$ is an analytic map; similarly for X' . We claim that the special model Z defined by $(U \times U', f \times f', F \times F')$ is a product. Indeed, from the description of the morphisms into a special model provided by Proposition 1.2.5. it follows that we have natural maps $\pi : Z \rightarrow X, \pi' : Z \rightarrow X'$ induced by the projections $U \times U' \rightarrow U, U \times U' \rightarrow U'$. Also, if $f : Y \rightarrow X$ and $f' : Y \rightarrow X'$ are given, $g : Y \rightarrow Z$ is determined by

$$\begin{array}{ccccc} & f \nearrow & X & \rightarrow & U & \searrow & & \\ Y & & & & & & U \times U' & \\ & f \searrow & X' & \rightarrow & U' & \nearrow & & \end{array}$$

In the general case we take $X \times X'$ as the ringed space whose topological underlying space is the cartesian product of the underlying space of X and X' , and whose structure sheaf is given locally by the product of local models for X and X' . (From the uniqueness “up to isomorphism” of the product results that these sheaves stick together in a well-determined way).

b) *Kernel of a double arrow.* If $X \begin{matrix} \xrightarrow{u} \\ \rightrightarrows \\ \xrightarrow{v} \end{matrix} Y$ is a double arrow, i.e. a pair of morphisms, a kernel X' of (u, v) is an analytic subspace of X such that the morphisms of an arbitrary analytic space Z into X' are exactly the morphisms h of Z into X such that $u \circ h = v \circ h$. In other words, if $i : X' \rightarrow X$ is the natural map of X' into X , the morphisms $h : Z \rightarrow X'$ satisfy $u \circ i \circ h = v \circ i \circ h$ and if a morphism $g : Z \rightarrow X$ satisfies $u \circ g = v \circ g$, then $g = i \circ h$ for some $h : Z \rightarrow X'$. To prove the existence of the kernel it suffices, again, to do this locally, i.e. for special models. If X is defined by (U, f, F) and Y by (V, g, G) we may (perhaps, after restricting U) extend u and v to maps $\bar{u}, \bar{v} : U \rightarrow E$ where E denotes the complex linear space of which V is an open subset. The kernel is then defined by the triple

$$(U, f \times (\bar{u} - \bar{v}), F \times E).$$

It follows from the Proposition 1.2.5. that this special model satisfies the universal property of kernels.

Example 1. The kernel of $\mathbf{C} \begin{matrix} \xrightarrow{t} \\ \rightrightarrows \\ \xrightarrow{-t} \end{matrix} \mathbf{C}$ is the simple point $\{0\}$, t denoting the identity of \mathbf{C} .

Example 2. The kernel of $\mathbf{C} \begin{matrix} \xrightarrow{t} \\ \rightrightarrows \\ \xrightarrow{t+t^2} \end{matrix} \mathbf{C}$ is $\{0\}$ counted as a double point.

c) *Fiber product.* If $u : X \rightarrow S$ and $v : Y \rightarrow S$ are given morphisms of analytic spaces, the fiber product $X \times_s Y$ of X and Y over S is the kernel of the double arrow

$$X \times Y \begin{matrix} \xrightarrow{u \circ \pi} \\ \rightrightarrows \\ \xrightarrow{v \circ \pi'} \end{matrix} S$$

where $\pi : X \times Y \rightarrow X$ and $\pi' : X \times Y \rightarrow Y$ are the maps defined by the product. Note that when S is a simple point, $X \times_s Y = X \times Y$.

One may also introduce the category of analytic spaces over S . Its objects are morphisms $u : X \rightarrow S$ of an analytic space X onto S and its morphisms are morphisms $f : X \rightarrow Y$ such that the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ u \searrow & & \swarrow v \\ & S & \end{array}$$

is commutative. The product in this category, i.e. the object satisfying the universal property given above for the product $X \times Y$, is then exactly the fiber product $X \times_S Y$. If S is a point, we have the category of analytic spaces.

Example 3. If U and V are open subspaces of an analytic space X , the open subspace $U \cap V$ is isomorphic to $U \times_X V$. We may thus define, in general, the intersection of two analytic subspaces $X' \rightarrow X$ and $X'' \rightarrow X$ of X to be the fiber product $X' \times_X X''$.

Example 4. If $\varphi : Y \rightarrow X$ is a morphism of analytic spaces and $a \in X$ a point, i.e. a map $a : (0, \mathbf{C}) \rightarrow X$ we may consider the space $Y(a) = Y \times_X a$. It is natural to call this the inverse image of a under φ and to denote it by $\varphi^{-1}(a)$; its underlying space is exactly $\varphi_0^{-1}(a)$.

If $\varphi_0(b) = a$, then $\mathcal{O}_{Y(a),b}$ is $\mathcal{O}_{Y,b}$ taken modulo the image under $\varphi^1 : \mathcal{O}_{X,a} \rightarrow \mathcal{O}_{Y,b}$ of the maximal ideal in $\mathcal{O}_{X,a}$.

Example 5. The pull-back of a linear bundle E over X by a map $Y \rightarrow X$ is exactly $Y \times_X E$.

1.4. Relations between reduced and non-reduced spaces.

We shall first characterize those analytic spaces which are reduced.

Proposition 1.4.1. A analytic space (X, \mathcal{O}_X) is reduced if and only if $\mathcal{O}_{X,x}$ has no nilpotent element for x arbitrary in X .

Proof. The necessity of the condition is obvious for \mathcal{O}_X can be considered as a submodule of \mathcal{C}_X if (X, \mathcal{O}_X) is reduced.

Conversely, if $\mathcal{O}_{X,x}$ has no nilpotent elements, we shall prove that in any local model (V, \mathcal{O}_V) for (X, \mathcal{O}_X) , a germ g at $a \in V$ which vanishes on V belongs to the ideal \mathcal{I} defining \mathcal{O}_V . The Nullstellensatz implies that $g^k \in \mathcal{I}_a$ if k is large enough. But it is then clear that $g \in \mathcal{I}_a$ if $\mathcal{O}_{V,a}/\mathcal{I}_a$ is free from nilpotent elements.

Given an analytic space (X, \mathcal{O}_X) we can associate to it a reduced space in the following way. Let \mathcal{N}_x be the ideal in $\mathcal{O}_{X,x}$ consisting of all nilpotent elements (the nil-radical of 0). Then $\mathcal{N} = U\mathcal{N}_x$ is a coherent sheaf by the Oka-Cartan theorem, for in a local model (V, \mathcal{O}_V) for (X, \mathcal{O}_X) we have $\mathcal{N}_X = (\mathcal{I}'/\mathcal{I})_X$ where \mathcal{I}' is the sheaf of germs vanishing on V and \mathcal{I} the