

§1. Acyclic maps and homotopy equivalences

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **25 (1979)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **22.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

§ 1. ACYCLIC MAPS AND HOMOTOPY EQUIVALENCES

We will use the terminology *CW-space* for a space having the homotopy type of a *CW-complex*. The category of *CW-spaces* is the largest category of spaces for which the Whitehead characterization of homotopy equivalences holds.

(1.1) DEFINITION. *A space X is acyclic provided the integral reduced homology $\tilde{H}_*(X) = 0$.*

In particular, an acyclic space X is path connected, its fundamental group $\pi_1(X)$ is perfect, i.e. $\pi_1(X)$ is equal to its commutator subgroup, and for any constant coefficient module L it follows that $\tilde{H}_*(X, L) = 0$. Recall that a local coefficient system L on X is a module over $\pi_1(X)$ and that

$$H_*(X, L) = H_*(C_*(X) \otimes_{\mathbf{Z}\pi_1(X)} L)$$

where $C_*(\tilde{X})$ is the chain complex over \mathbf{Z} viewed as a $\mathbf{Z}\pi_1(Y)$ -module. In general, $\tilde{H}(X, L) \neq 0$ for an acyclic space and a local coefficient system L .

(1.2) DEFINITION/PROPOSITION. *A map $f: X \rightarrow Y$ between path connected spaces is acyclic provided any of the following equivalent conditions hold:*

- (a) *The homotopy fibre F of $f: X \rightarrow Y$ is an acyclic space.*
- (b) *For any local coefficient system L on Y the induced morphism*

$$f_* : H_*(X, f^*L) \rightarrow H_*(Y, L)$$

*is an isomorphism where f^*L is the induced local system on X .*

- (c) *The induced morphism*

$$f_* : H_*(X, f^*\mathbf{Z}\pi_1(Y)) \rightarrow H_*(Y, \mathbf{Z}\pi_1(Y))$$

is an isomorphism.

- (d) *For the universal covering $\tilde{Y} \rightarrow Y$ of Y the map $X \times_Y \tilde{Y} \rightarrow \tilde{Y}$ defined by f induces an isomorphism*

$$H_*(X \times_Y \tilde{Y}) \rightarrow H_*(\tilde{Y}).$$

Proof. For (a) implies (b), we use the Serre spectral sequence for the fibration $F \xrightarrow{i} X \xrightarrow{f} Y$ where

$$E^2 = H_* (Y, H_* (F, i^* f^* L)) \Rightarrow H_* (X, f^* L).$$

Since $i^* f^* L$ is trivial on F , statement (a) gives $\tilde{H}_* (F, i^* f^* L) = 0$ and the edge morphism $H_* (X, f^* L) \rightarrow H_* (Y, L) = E^2_{*,0}$, which is induced by f , is an isomorphism.

Clearly (b) implies (c), which is a special case of (b), and for (c) implies (d) we use the following morphism of fibrations

$$\begin{array}{ccc} & \pi = \pi_1(Y) & \\ & \swarrow \quad \searrow & \\ X \times_Y \tilde{Y} & \longrightarrow & \tilde{Y} \\ \downarrow & & \downarrow \\ X & \xrightarrow{f} & Y. \end{array}$$

This induces a morphism of the Serre spectral sequences which on the E^2 -level is the given isomorphism from (c)

$$E^2 = H_* (X, f^* \mathbf{Z}\pi_1(Y)) \rightarrow H_* (Y, \mathbf{Z}\pi_1(Y)) = E^2.$$

Hence by the spectral mapping theorem $H_* (X \times_Y \tilde{Y}) \rightarrow H_* (\tilde{Y})$ is an isomorphism.

For (d) implies (a), note that $F \rightarrow X \times_Y \tilde{Y}$ is the fibre of $X \times_Y \tilde{Y} \rightarrow \tilde{Y}$. Since $H_* (X \times_Y \tilde{Y}) \rightarrow H_* (\tilde{Y})$ is an isomorphism on the horizontal edge of the spectral sequence, we see $\tilde{H}_0(F) = 0$. Moreover, assuming inductively that $\tilde{H}_j(F) = 0$ for $i < n$, we deduce that $\tilde{H}_n(F) \doteq 0$ by looking at the spectral sequence terms $E^r_{0,n}$ which is $H_n(F)$ for $r = 2$ and zero for $r > n + 1$. This completes the proof the equivalence of (a), (b), (c), and (d).

(1.3) PROPOSITION. *If $f : X \rightarrow Y$ is an acyclic map, then $f_* : \pi_1(X) \rightarrow \pi_1(Y)$ is an epimorphism with kernel a perfect normal subgroup.*

Proof. Since the fibre F of f is connected, the induced homomorphism f_* is an epimorphism, and since $\pi_1(F)$ is perfect, $\ker(f_*) = \text{im}(\pi_1(F) \rightarrow \pi_1(X))$ is perfect.

(1.4) PROPOSITION. Let $f: X \rightarrow Y$ be a map between path connected spaces. Then $\pi_i(f): \pi_i(X) \rightarrow \pi_i(Y)$ is an isomorphism for all $i \geq 0$ if and only if f is acyclic and $\pi_1(f)$ is an isomorphism.

Proof. Let $F \rightarrow X$ be the homotopy fibre of f . The second conditions say that $\pi_1(F)$ is perfect and abelian respectively. Thus $\pi_1(F) = 0$ and on simply connected spaces F the homotopy $\pi_i(F) = 0$ if and only if the homology $\tilde{H}_i(F) = 0$. The proposition follows now from an application of the homotopy exact sequence.

(1.5) COROLLARY. A map $f: X \rightarrow Y$ between path connected CW-spaces is a homotopy equivalence if and only if f is acyclic and $\pi_1(f)$ is an isomorphism.

This is an immediate application of the Whitehead criterion for homotopy equivalence applied to (1.4).

In section 3 we will see that the subgroups $\ker(\pi_1(f))$ classify acyclic maps $f: X \rightarrow Y$ from X .

(1.6) Remark. Cohomology with local coefficients can be used to characterize acyclic maps. As with homology

$$H^*(X, L) = H^*(\text{Hom}_{\mathbb{Z}\pi_1(X)}(C^*(\tilde{X}), L))$$

defines cohomology with local coefficients. Then a map $f: X \rightarrow Y$ between path connected spaces is acyclic if and only if $f^*: H^*(Y, L) \rightarrow H^*(X, f^*L)$ is an isomorphism for each local coefficient system L on Y . The direct implication is checked exactly as (a) implies (b) using cohomology in (1.2). Conversely we show that $X \times_Y \tilde{Y} \rightarrow \tilde{Y}$ defined by f induces an isomorphism $H^*(\tilde{Y}) \rightarrow H^*(X \times_Y \tilde{Y})$. This is done as (c) implies (d) in (1.2) and as in (d) implies (a) in (1.2) we have $\tilde{H}^*(F) = 0$. Using the universal coefficient theorem, we deduce that $\tilde{H}_*(F) = 0$ and F is acyclic.

The cohomology characterization of acyclic maps is useful in obstruction theory.

(1.7) Remark. Let $f: X \rightarrow Y$ be an acyclic map and \bar{Y} a connected covering of Y . Then the induced map $\bar{f}: X \times_Y \bar{Y} \rightarrow \bar{Y}$ is also acyclic. This follows directly from (1.2, (d)) or from the fact that f and \bar{f} have the same fibre. When \bar{Y} is the universal covering of Y , the space $X \times_Y \bar{Y} = \tilde{X}_N$ is the covering of X with fundamental group $N = \ker(\pi_1(f))$.