

2. Proof of Theorem B

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **25 (1979)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **24.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

2. PROOF OF THEOREM B

2.1. The complex analytic case. Let G be a reductive complex algebraic group. Then G is the universal complexification of a compact real Lie group $G^{\mathbf{R}}$ [9], [8, XVII.5].

Suppose that G acts linearly on $V = \mathbf{C}^n$, and that X is a germ at 0 of an invariant closed analytic subset of V . Let I be the ideal in $\mathbf{C}\{x\} = \mathbf{C}\{x_1, \dots, x_n\}$ of germs of holomorphic functions which vanish on X . Suppose that I is generated by $f_1, \dots, f_k$.

For any irreducible complex representation $T: G \rightarrow GL(W)$ of G , we consider the action of G on the space $\text{End}_{\mathbf{C}}(W, W)$ of complex linear endomorphisms defined by

$$(g \cdot \lambda)(w) = T(g) \lambda(w),$$

where $g \in G$, $w \in W$ and $\lambda \in \text{End}_{\mathbf{C}}(W, W)$. For each $i = 1, \dots, k$, we consider the mapping

$$f_i^T: V \rightarrow \text{End}_{\mathbf{C}}(W, W)$$

$$f_i^T(x) = \int_{G^{\mathbf{R}}} f_i(g^{-1}x) T(g) dg,$$

defined in an open neighborhood of 0 where f_i converges. Then f_i^T is equivariant with respect to the actions of $G^{\mathbf{R}}$ on V and $\text{End}_{\mathbf{C}}(W, W)$, and hence with respect to the actions of G (the "unitarian trick"). Furthermore $f_i(gx) = 0$ for all $g \in G$ if and only if $f_i^T(x) = 0$ for all irreducible complex representations T of $G^{\mathbf{R}}$ (cf. [10, 12.2]; this is essentially the Peter-Weyl theorem).

Hence X is defined by the equations

$$f_i^T(x) = 0,$$

where $1 \leq i \leq k$ and T runs over all irreducible complex representations of $G^{\mathbf{R}}$. It follows that X is defined by a finite subset of these equivariant equations.

2.2. The real analytic case. Let G be a reductive real algebraic group. Then the universal complexification $G^{\mathbf{C}}$ of G is a reductive complex algebraic group [8, XVIII.4].

Suppose that G acts linearly on $V = \mathbf{R}^n$, and that X is a germ at 0 of an invariant real analytic subset of V . The complexification $X^{\mathbf{C}}$ of X is a germ at 0 of a complex analytic subset of $V^{\mathbf{C}} = \mathbf{C}^n$. The complexification $X^{\mathbf{C}}$ is invariant under the induced action of $G^{\mathbf{C}}$ on $V^{\mathbf{C}}$.

By the complex analytic case 2.1, there is a linear action of $G^{\mathbf{C}}$ on a finite dimensional complex vector space W , and a germ H at 0 of a $G^{\mathbf{C}}$ -equivariant holomorphic mapping of some neighborhood of $0 \in V^{\mathbf{C}}$ into W , such that $X^{\mathbf{C}} = H^{-1}(0)$.

Let Y be W with its underlying real structure. Then $F = H|V: V \rightarrow Y$ is G -equivariant, and $X = F^{-1}(0)$.

2.3. The algebraic case. Our ground field $\mathbf{k}$ is now arbitrary. Let G be a reductive algebraic group acting linearly on $V = \mathbf{k}^n$, and let X be an invariant algebraic subset of V . Let I be the ideal in $\mathbf{k}[x]$ of polynomials which vanish on X , and $\mathbf{k}[x]_c$ be the subspace of $\mathbf{k}[x]$ of polynomials of degree at most c . Then I and $\mathbf{k}[x]_c$ are invariant subsets of $\mathbf{k}[x]$.

For each $c \in \mathbf{N}$, we define a polynomial mapping

$$F_c: V \rightarrow \text{End}_{\mathbf{k}}(I \cap \mathbf{k}[x]_c, \mathbf{k})$$

by the formula $F_c(x)(h) = h(x)$, where $x \in V$ and $h \in I \cap \mathbf{k}[x]_c$. Then F_c is equivariant and $X \subset F_c^{-1}(0)$ for all $c \in \mathbf{N}$.

We consider the ideal J in $\mathbf{k}[x]$ generated by the coordinate functions of all equivariant polynomial mappings defined on V , which vanish on X . Since J is finitely generated, it suffices to show that $J = I$. Clearly $J \subset I$. On the other hand, suppose $h \in I \cap \mathbf{k}[x]_c$, $h \neq 0$. Let $\{e_j\}_{1 \leq j \leq q}$ be a basis of the vector space $I \cap \mathbf{k}[x]_c$, such that $e_1 = h$. Then h is the first coordinate function of the equivariant mapping F_c , with respect to the dual basis $\{e_j^*\}_{1 \leq j \leq q}$ in $\text{End}_{\mathbf{k}}(I \cap \mathbf{k}[x]_c, \mathbf{k})$. Since $X \subset F_c^{-1}(0)$, then $h \in J$. Hence $J = I$ as required.

This case of Theorem B may also be obtained from a lemma of Cartier [13, p. 25].

3. PROOF OF THEOREM A

The formal power series $\bar{y}(x) \in \mathbf{k}[[x]]^p$ define a local $\mathbf{k}$ -homomorphism $\phi: \mathbf{k}\{x, y\} \rightarrow \mathbf{k}[[x]]$ (or a $\mathbf{k}$ -homomorphism $\phi: \mathbf{k}[x, y] \rightarrow \mathbf{k}[[x]]$ in the algebraic case) by substitution: $h(x, y) \rightarrow h(x, \bar{y}(x))$.