

6. The local fundamental group

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **25 (1979)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **26.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Proposition 5.2 shows that characterizations A5' and A1 are equivalent. Clearly Characterization A5' implies A5; since A5 implies A2, they are all equivalent.

COROLLARY 5.3. *Let G be a small finite subgroup of $GL(2, \mathbb{C})$. Then $G \subset SL(2, \mathbb{C})$ if and only if $\mathbb{C}^2/G$ embeds in codimension one.*

This corollary follows from the above case-by-case analysis. J. Wahl points out that it is also possible to prove it directly, using the following two facts:

Fact 1. Let G be a small finite subgroup of $GL(2, \mathbb{C})$. Then $G \subset SL(2, \mathbb{C})$ if and only if the singularity of $\mathbb{C}^2/G$ is Gorenstein.

This is a special case of [Watanabe]. A germ of a normal two-dimensional complex space is *Gorenstein* if there is a nowhere-vanishing holomorphic two-form on its regular points.

Fact 2. Let V be the germ at $\mathbf{v}$ of a two-dimensional rational singularity. Then V is Gorenstein if and only if V embeds in codimension 1.

Proof. Any singularity embedded in codimension one is Gorenstein. Conversely, suppose V is Gorenstein. Let $\pi: M \rightarrow V$ be the minimal resolution of V , and let $E_1 \cup \dots \cup E_s = \pi^{-1}(\mathbf{v})$ be its exceptional set as in Section 3. Since V is Gorenstein, there is a divisor K on M (the *canonical class*) satisfying the adjunction formula. Furthermore $K \cdot E_i \geq 0$ for all i since the resolution is minimal, so $K \leq 0$ [Artin, bottom of p. 130]. If $K < 0$, then $-K > 0$, so arithmetic genus p of $-K$ satisfies $p(-K) \leq 0$ [Artin, Proposition 1]. On the other hand, $p(-K) = 1 - \chi(-K) = 1$ by the Riemann-Roch Theorem, a contradiction. Hence $K = 0$. Thus $K \cdot E_i = 0$ for all i , so V is a double point and embeds in codimension one, as in the proof that Characterization A3 implies Characterization A2.

6. THE LOCAL FUNDAMENTAL GROUP

Let V be the germ of a normal two-dimensional complex analytic space with an isolated singularity at $\mathbf{v}$. Without loss of generality, we may assume that V is a *good neighborhood* of $\mathbf{v}$, that is, that there is a neighborhood basis V_i of $\mathbf{v}$ in V such that each $V_i - \mathbf{v}$ is a deformation retract of $V - \mathbf{v}$ [Prill]. The *local fundamental group* of V at $\mathbf{v}$ is then defined as $\pi_1(V - \mathbf{v})$. This group is trivial if and only if V is nonsingular at $\mathbf{v}$ [Mumford].

PROPOSITION 5.1 (bis). *The following statement is equivalent to those listed above.*

(d) *The local fundamental group of V is finite.*

It is shown in [Prill, p. 381; Brieskorn 2, p. 344] that conditions (a) and (d) are equivalent.

Characterization A6. The local fundamental group of $f^{-1}(0)$ is finite. Thus Characterizations A5 and A6 are equivalent.

There is an algorithm for computing the local fundamental group of V from a resolution [Mumford], and singularities V with finite, nilpotent and solvable local fundamental group have been classified [Brieskorn 2; Wagreich 2]. When V is a complete intersection, this classification is particularly simple [Durfee 2, Proposition 3.3].

7. VOLUME

Let $f(x, y, z)$ be the germ at the origin 0 of a complex analytic function, and suppose that $f(0) = 0$ and that the origin is an isolated critical point of f . There is an $\varepsilon > 0$ such that $f^{-1}(0)$ intersects all spheres of radius ε' about 0 transversally for $0 < \varepsilon' \leq \varepsilon$. (See Section 12.) For $t \in \mathbb{C}$, let

$$V_t = f^{-1}(t) \cap D_\varepsilon^6$$

where D_ε^6 is the closed disk of radius ε about 0 . The function $f(x, y, z)$ takes the constant value t on V_t , so $\frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy + \frac{\partial f}{\partial z} dz \equiv 0$ there. Hence a nowhere-vanishing holomorphic two-form ω_t on V_t may be defined by the equivalent expressions

$$\omega_t = \frac{dy \wedge dz}{\partial f / \partial x} = \frac{dz \wedge dx}{\partial f / \partial y} = \frac{dx \wedge dy}{\partial f / \partial z},$$

Characterization A7. The integral $\int_{V_0} \omega_0 \wedge \bar{\omega}_0$ is finite.

Note that the form $\omega_0 \wedge \bar{\omega}_0$ takes positive real values. The equivalence of Characterizations A2 and A7 is due to Laufer, and follows easily from his expression for the geometric genus in terms of forms [Laufer 2, Corollary 3.6].

A different formulation of this characterization is due to E. Looijenga (unpublished): Let $\Delta(r) = \{t \in \mathbb{C} : |t| < r\}$, let

$$X(r) = f^{-1}(\Delta(r)) \cap D_\varepsilon^6$$

and let $\text{vol}(X(r))$ be its volume in $\mathbb{C}^3$.