

V. Construction of the Model

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **28 (1982)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **19.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Proposition 2 may be expressed by a \prod_2^0 formula. First it is clear that we can construct a \sum_1^0 -formula ϕ_i that expresses the properties that

1. $P_i : [\mathbb{N}]^{e_i} \rightarrow r_i$ is a primitive recursive partition
2. $z_1 < z_2 < \dots < z_{n_k}$
3. $\{z_1, \dots, z_{n_k}\}$ is homogeneous for P_i
4. $k \leq n_k$
5. $2^{2^{z_1}} \leq n_k$

Proposition 2 asserts that for every k

$$\mathbb{N} \models \exists z_1 \dots \exists z_{n_k} \bigwedge_{i \leq k} \phi_i .$$

V. CONSTRUCTION OF THE MODEL

We now have all the ingredients at hand to construct a non-standard model of Peano arithmetic, and we have only to assemble them according to the specifications of Section II.

Let P_i be an effective enumeration of all primitive recursive partitions $P_i : [\mathbb{N}]^{e_i} \rightarrow r_i$. By Proposition 2 we have that for every k

$$\mathbb{N} \models \exists z_1 \dots \exists z_{n_k} \bigwedge_{i \leq k} \phi_i$$

where ϕ_i is the \sum_1^0 -formula of Section IV expressing the conditions (1)-(5) satisfied by the partition P_i .

Following the prescription given in Section III we let a_{kn_k} be the smallest number such that a_{k_1}, \dots, a_{kn_k} is an increasing sequence satisfying the formula $\bigwedge_{j \leq k} \phi_j$. Now we define the functions h_j by

$$h_0(k) = n_k \quad \text{for every } k$$

and for $j > 0$

$$h_j(k) = \begin{cases} a_{kj} & \text{for } j \leq n_k \\ h_{j-1}(k)^2 & \text{for } j > n_k . \end{cases}$$

Let $\mathcal{F} = \{f \mid f \leq h_j\}$.

Since $1 \leq h_0$ the function 1 is automatically in \mathcal{F} .

By Theorem 2 the sequence $\{h_j\}$ satisfies $\bigwedge_{j < \infty} \phi_j$ in \mathcal{F}/D . We now prove that this implies that the sequence $\{h_j\}$ satisfies the Stability and

Closure Conditions in \mathcal{F}/D . As we saw in Section III it suffices for this purpose to show that for each k

$$\begin{aligned} \mathbf{N} \vdash \exists z_1 \dots \exists z_{n_k} \bigwedge_{\substack{1 \leq i < j, j' < n_k \\ 1 \leq s < k}} [(\forall y < z_i) (\psi_s(y; z_j) \\ \leftrightarrow \psi_s(y; z_{j'}) \wedge z_{j-1}^2 < z_j)] \end{aligned} \quad (*)$$

Let t_i be the length of the sequence y in $\psi_i(y; z)$. Define the partitions $T: [\mathbf{N}]^2 \rightarrow 2$, $Q_i: \mathbf{N} \rightarrow t_i^2 + 1$, and $S_i: [\mathbf{N}]^{2e+1} \rightarrow 2$ by:

$$T(a, b) = \begin{cases} 1 & \text{if } a^2 < b \\ 0 & \text{if not} \end{cases}$$

$$Q_i(a) = \min(a, [t_i \log_2 a] + 1)$$

and for $a \in \mathbf{N}$, $c, c' \in [\mathbf{N}]^e$

$$S_i(a, c, c') = \begin{cases} 1 & \text{if } (\forall y < a) (\psi_i(y; c) \leftrightarrow \psi_i(y; c')) \\ 0 & \text{if not.} \end{cases}$$

The partitions T , Q_i , and S_i are clearly primitive recursive since $\psi_i(y; z)$ is a limited formula. Hence $T, Q_1, \dots, Q_k, S_1, \dots, S_k$ occur in the sequence $\{P_i\}$. Thus by looking sufficiently far in the sequence we can find a set $X_k = \{a_{k1}, \dots, a_{kn_k}\}$ which is homogeneous for $T, Q_1, \dots, Q_k, S_1, \dots, S_k$ with $n_k \geq k, 2^{2^{a_{k1}}}$.

Since $a_{kn_k} > a_{k1}^2$, $T(a_{k1}, a_{kn_k}) = 1$. Hence, by homogeneity,

$$T(a, b) = 1, \quad \text{i.e. } a^2 < b,$$

for all $a < b$ in X_k .

Since $\#X > 1$, and X_k is homogeneous for Q_i , $a_{k1} \geq t_i \log_2 a_{k1}$.

The number of sequences of numbers $< a_{k1}$ of length t_i is $< a_{k1}^{t_i}$. The number of distinct sequences of truth values of length $a_{k1}^{t_i}$ is $< 2^{a_{k1}^{t_i}}$. Now $n_k > 2^{2^{a_{k1}}} > 2^{a_{k1}^{t_i}}$ since $a_{k1} \geq t_i \log_2 a_{k1}$. Thus there are distinct $c, c' > a$ in X_k such that

$$(\forall y < a_{k1}) (\psi_i(y; c) \leftrightarrow \psi_i(y; c')),$$

i.e. $S_i(a_{k1}, c, c') = 1$.

By homogeneity

$$S_i(a, b, b') = 1 \quad \text{for all } a < b, b' \text{ in } X_k$$

proving (*).

We have thereby shown that \mathcal{F}/D is a model of the Peano axioms. Since a_{kn_k} was chosen minimal, Proposition 2 is false in \mathcal{F}/D , and hence independent of the Peano axioms.

Proposition 1 is also false in \mathcal{F}/D . In fact it is provable in Peano arithmetic that Proposition 1 implies Proposition 2. This is a consequence of the following lemma, provable in Peano arithmetic (c.f. Lemma 2.9 in [3]).

LEMMA 2. *Let $P_i : [\mathbb{N}]^e \rightarrow r_i$, $1 \leq i \leq n$, be n partitions. There is a partition $P : [\mathbb{N}]^e \rightarrow r$ such that for all subsets H of \mathbb{N} of cardinality $> e$, H is homogeneous for P if and only if H is homogeneous for all the P_i .*

We may also obtain a purely finitary combinatorial principle which is false in our model.

PROPOSITION 3. *For all natural numbers e , r , and k there exists an N , such that for all partitions $P : [N]^e \rightarrow r$ there exists a subset X of N , with $\# X \geq k$ and $\# X \geq 2^{2^{\min X}}$, which is homogeneous for P .*

This result follows immediately from the infinite Ramsey Theorem by an application of König's Lemma. If we drop the condition that $\# X \geq 2^{2^{\min X}}$, then we obtain the usual finite Ramsey Theorem. Ramsey [11] gave a proof of the latter theorem which is formalizable in Peano arithmetic. Proposition 3 directly yields Proposition 1, for if $P : [\mathbb{N}]^e \rightarrow r$ is a partition and k is a number then by considering the partition $P \upharpoonright [N]^e$, where N is the number provided by Proposition 3 we obtain the required homogeneous set X for $P \upharpoonright [N]^e$ and hence for P . This proof may be carried out in Peano arithmetic. Thus, Proposition 3 is false in our model and independent of the Peano axioms.

VI. A SIMPLER MODEL

The condition in Proposition 1 that $\# X \geq 2^{2^{\min X}}$ can be simplified and so yield a simpler sequence $\{h_i\}$ of functions which define the model \mathcal{F}/D . In this section we describe such a model by using a combinatorial consequence of Ramsey's Theorem which is closer to the proposition proved independent in [3].

PROPOSITION 4. *Let $P : [\mathbb{N}]^e \rightarrow r$ be a primitive recursive partition. For every k there exists a finite subset X of \mathbb{N} , with $\# X \geq k$ and $\# X \geq \min X$, which is homogeneous for the partition P .*

Proposition 4 implies Proposition 1 via the following result, the proof of which is the same as the proof of Lemma 2.14 of [3].