

Affine structures in 2, 3, and 4 dimensions

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **29 (1983)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **19.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

space M' is obtained by gluing, each time along one of the two segments of a or c , as many copies of open sectors bounded by the lines a and c , (each covering an open annulus [5]) as there are letters in the characteristic word. These projective structures on the 2-torus are characterized by their (cyclic) word and the $t = 1$ flow map. In suitable homogeneous coordinates the last is expressed as $f_1 : f_t : (x, y, z) \rightarrow (xe^{\alpha t}, ye^{\beta t}, ze^{\gamma t})$ $\alpha < \beta < \gamma$, $t = 1$.

Remark. Following the curve from its initial point P to its endpoint P' , one can say that the sectors of P and P' were identified by the identity map: in homogeneous coordinates.

$$(x, y, z) \rightarrow (x, y, z)$$

A more general case (see Goldman [5]) is obtained if we identify by any projectivity commuting with f_1 :

$$g : (x, y, z) \rightarrow (xe^{\lambda}, ye^{\mu}, ze^{\nu})$$

$\lambda, \mu, \nu \in \mathbf{R}$.

AFFINE STRUCTURES IN 2, 3, AND 4 DIMENSIONS

In dimension two only the torus admits an affine structure by Benzecri [1] and for all affine structures the developing map is a covering of its image by Nagano-Yagi [7]. The image is affinely equivalent to either the whole plane, the once punctured plane, the half plane or the quarter plane.

We obtain interesting affine structures in dimensions 3 and 4 using respectively the projective and inversive structures in dimension 2 discussed above.

i) A projective transformation of the real projective plane $\mathbf{RP}^2 = \mathbf{R}^3 - \{0\}/\mathbf{R}^*$ (where $\mathbf{R}^* = \mathbf{R} - \{0\}$) lifts to an affine transformation of $V = \mathbf{R}^3 - \{0\}$, unique but for scalar multiplication. Any such commutes with scalar multiplication by a real number $\alpha > 1$ (e.g. $\alpha = 2$).

Thus one may build an affine 3-manifold using as a pattern a projective two manifold (open sets in the projective plane lift to open sets (cones) in V etc.). If we further divide by the action of a compactness is preserved in the construction.

The projective structures on the two torus constructed above yield compact affine 3-manifolds where the developing map is not a covering. In particular, from the example in Figure 7, we can obtain an affine 3-manifold which develops

over the part outside the coordinate axes of $\{X > 0\} \cup \{Z > 0\} \subset \mathbf{R}^3 = \{(x, y, z)\}$, but not as a covering. In these examples the 3-manifold M is a 3-torus.

ii) Similarly, a projective transformation of the complex projective line $\mathbf{CP}^1 = \mathbf{C}^2 - \{0\}/\mathbf{C}^*$, that is to say an orientable conformal or inversive transformation of $S^2 = \mathbf{CP}^1$, lifts to a complex affine transformation of $V = \mathbf{C}^2 - \{0\}$, unique but for scalar multiplication and commuting with scalar multiplication.

We can build a four dimensional affine manifold from an inversive 2-manifold, which is actually a complex affine manifold of \mathbf{C} -dimension 2, and this construction is the analogue of the above over \mathbf{C} , thinking of S^2 as \mathbf{CP}^1 and the conformal transformations as the \mathbf{C} -projective transformation.

Again compactness is achieved if we divide by $\alpha = 2$. Thus using the inversive Example 2 we obtain affine 4-manifolds whose developing image has a complicated boundary related to the non-differentiable Jordan curve. Using Example 3, we obtain an affine four-manifold whose developing image in R^4 omits a Cantor set of two planes passing through the origin.

Using Example 4, we can build affine manifolds whose developing map is not a covering of its image (which is all of $\mathbf{C}^2 - 0$). And we repeat, all the above are actually complex affine structures on compact 4-manifolds.

NOTE 1 (see page 16). *Ehresmann* defined the *development map* as follows. Let $\mathcal{P} \rightarrow M$ be the principle \mathcal{A} -bundle over M , whose points are germs $[x, \kappa]$ of canonical charts $\{x \in U \subset M, \kappa: U \rightarrow A\}$. Define a new topology $\mathcal{F}(\mathcal{P})$ in the set \mathcal{P} by taking as open set the germs at all points $x \in U$ of any given chart $\kappa: U \rightarrow A$. The natural map $d: \mathcal{F}(\mathcal{P}) \rightarrow A$ is an immersion. Choose one component of $\mathcal{F}(\mathcal{P})$ and call it M' . The restriction $d: M' \rightarrow A$ is a development map. The restriction of the natural fibre bundle projection $p: \mathcal{F}(\mathcal{P}) \rightarrow M$ is a covering $M' \rightarrow M$.

NOTE 2 (see page 16). *The fibre bundle picture*. For the simple *local* discussion of *one canonical chart* $U \subset A$, we can describe a trivial fibre bundle $E_U = U \times A \rightarrow U$ by assigning to any $x \in U$ the "heavily osculating" model space $A_x = A$. The manifold U is embedded as the diagonal cross section. $s(U) = \{(y, y)\} = \text{diag}(U \times U) \subset U \times U \subset U \times A$. Its points are the points of tangency of fibre and base manifolds. Finally a foliation \mathcal{F} is defined as the one with horizontal leaves $U \times \{v\} \subset E_U = U \times A$, for $v \in A$.

For the *global* discussion of an \mathcal{A} -structure on a manifold M , we assume \mathcal{A} -compatible canonical charts that are topological embeddings $\kappa: U \hookrightarrow A$ for

small open sets $U \subset M$. A point of the fibre bundle space E over M is by definition a triple

$$\{x, \kappa, v\},$$

where $x \in U \subset M$, $\kappa: U \rightarrow A$ is a canonical chart and $v \in A$, *modulo equivalence* by the action of \mathcal{A} given by $g: \{x, \kappa, v\} \simeq \{x, \kappa', v'\}$ where $\kappa' = g \circ \kappa$, $v' = gv$, $g \in \mathcal{A}$. In E , M is embedded as the "diagonal cross section" $s(M)$, whose points are represented by triples $\{x, \kappa, \kappa(x)\}$. The foliation \mathcal{F} has the local "horizontal" leaves represented by triples $\{U, \kappa, v\}$. For contractible closed curves starting and ending at $x_0 \in M$ in the base space M , the holonomy of the foliation is of course the identity map of the fibre A_{x_0} . As a consequence for closed curves in general, starting and ending at x_0 the holonomy gives the representation of $\pi_1 M$ into the group \mathcal{A} acting on A_{x_0} . "Parallel displacement" of the points of $s(M)$ along the lifting in \mathcal{F} -leaves of curves in the base space ending at x_0 , determines the development map $M' \rightarrow A_{x_0}$.

NOTE 3 (see page 16). *Flat Cartan connections.* Manifolds with canonical (\mathcal{A}, A) -charts are the flat cases (without torsion and without curvature) of *manifolds M with a general (\mathcal{A}, A) -connection.* They are defined in [4] as follows

- (1) A fibre bundle $A \rightarrow E \rightarrow M$ with fibre A over M
- (2) A fixed cross section $s(M)$
- (3) An n -plane field ξ in E transversal to the fibres and transversal to the fixed cross sections, such that
- (4) The holonomy obtained by lifting a closed curve starting and ending at $x_0 \in M$, into all curves tangent to ξ , belongs to \mathcal{A} acting on A_{x_0} . It is in general different for homotopic curves. It is flat if contractible closed curves have trivial holonomy (= identity).

The development of a curve ending in x_0 in M , is obtained by dragging along ξ the corresponding points of $s(M)$ until they arrive in the fibre A_{x_0} . In the flat case homotopic curves with common initial and end points give the same image of the initial point in the end fibre and the development map is achieved.