

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 41 (1995)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: CUBIC FORMS AND COMPLEX 3-FOLDS
Autor: Okonek, Ch. / Van de Ven, A.
Kapitel: 4. Invariants of complex 3-folds
DOI: <https://doi.org/10.5169/seals-61829>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 02.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The number of integral classes in these orbits is therefore finite. We have, however, an even stronger finiteness theorem for stable ternary cubics:

PROPOSITION 7. *Let H be a free \mathbf{Z} -module of rank 3. There exist only finitely many classes of symmetric trilinear forms $F \in S^3 H^\vee$ with a fixed discriminant $\Delta \neq 0$.*

Proof. In terms of Arnhold's invariants S and T , Δ is given by $\Delta = S^3 - T^2$. By a theorem of C. Siegel [Si], the diophantine equation $S^3 - T^2 = \Delta$ has only finitely many integral solution (S, T) for any integer $\Delta \neq 0$. For each of these solutions the corresponding point in $S^3 H_C^\vee / SL(H_C)$ lies outside of the discriminant curve, so that the π -fiber over it is a closed $SL(H_C)$ -orbit. The finiteness of the class number then follows from the Borel/Harish-Chandra theorem.

A famous special case of Siegel's theorem is Bachet's equation $S^3 - T^2 = 2$; it has only the two obvious solutions $(3, \pm 5)$.

REMARK 10. To get finiteness results for ternary cubic forms it is not sufficient to fix the J -invariant (instead of the discriminant): The forms $f_m = X^3 + XZ^2 + Z^3 + mY^2Z$, $m \in \mathbf{Z} \setminus \{0\}$, all have the same J -invariant, but they are not equivalent, even over \mathbf{Q} , since they have bad reduction at different primes $p \mid m$.

4. INVARIANTS OF COMPLEX 3-FOLDS

In this section we begin to investigate the topology of 1-connected, compact, complex 3-folds. After a brief discussion of the possible systems of Chern numbers of almost complex 6-manifolds, we study the behaviour of the topological invariants of complex 3-folds under certain standard constructions, like e.g. branched coverings, or blow-ups of points and curves. Then we describe some interesting examples of 1-connected, non-Kählerian 3-folds, including a new construction method which generalizes the Calabi-Eckmann manifolds. These examples will be needed in the next section in order to realize complex types of cubic forms as cup-forms of complex 3-folds.

4.1 CHERN NUMBERS OF ALMOST COMPLEX STRUCTURES

Let X be a closed, oriented, 6-dimensional differentiable manifold. The tangent bundle of X is induced by a classifying map $t_X: X \rightarrow BSO(6)$ which is unique up to homotopy. By an almost complex structure on X we mean the homotopy class $[\tilde{t}_X]$ of a lifting $\tilde{t}_X: X \rightarrow BU(3)$ of t_X to $BU(3)$.

PROPOSITION 8. *Every closed, oriented, 6-dimensional C^∞ -manifold X without 2-torsion in $H^3(X, \mathbf{Z})$ admits an almost complex structure. There is a 1-1 correspondence between almost complex structures on X and integral lifts $W \in H^2(X, \mathbf{Z})$ of $w_2(X)$. The Chern classes c_i of the almost complex manifold (X, W) are given by $c_1 = W$, $c_2 = \frac{1}{2}(W^2 - p_1(X))$.*

Proof (cf. [W]). The obstructions against lifting t_X to $BU(3)$ lie in the cohomology groups $H^{i+1}(X, \pi_i(SO(6)/_{U(3)}))$, $i = 0, 1, \dots, 5$. Since $SO(6)/_{U(3)} = \mathbf{P}^3$ has only one nontrivial homotopy group $\pi_2(SO(6)/_{U(3)}) \cong \mathbf{Z}$ in dimensions $i \leq 5$, there is in fact only one obstruction $o(t_X) \in H^3(X, \mathbf{Z})$, and this obstruction can be identified with the image of $w_2(X)$ under the Bockstein homomorphism $\beta: H^2(X, \mathbf{Z}/_2) \rightarrow H^3(X, \mathbf{Z})$. Since $H^3(X, \mathbf{Z})$ has no 2-torsion by assumption, $\beta w_2(X)$ must be equal to zero, so that X has at least one almost complex structure $[\tilde{t}_X] \in [X, BU(3)]$. Standard homotopy arguments show now that the map, which assigns to an almost complex structure $[\tilde{t}_X]$ its first Chern class $\tilde{t}_X^* c_1$, induces a 1-1 correspondence between integral lifts $W \in H^2(X, \mathbf{Z})$ of $w_2(X)$ and homotopy classes of liftings of $[t_X]$ to $BU(3)$.

The second Chern class c_2 of the almost complex manifold (X, W) is determined by $W^2 - 2c_2 = p_1(X)$.

The Chern numbers $c_1^3, c_1 c_2, c_3$ of an almost complex manifold X of real dimension 6 satisfy the following congruences: $c_1^3 \equiv 0 \pmod{2}$, $c_1 c_2 \equiv 0 \pmod{24}$, $c_3 \equiv 0 \pmod{2}$. Conversely, given a triple (a, b, c) of integers $a \equiv 0 \pmod{2}$, $b \equiv 0 \pmod{24}$, and $c \equiv 0 \pmod{2}$, there always exists an almost complex manifold X of dimension 6 with Chern numbers $c_1^3 = a$, $c_1 c_2 = b$, $c_3 = c$.

It is not totally clear, however, that one can find a *connected* manifold X with prescribed Chern numbers [H1].

PROPOSITION 9. *Every triple $(a, b, c) \in \mathbf{Z}^{\oplus 3}$ satisfying $a \equiv 0 \pmod{2}$, $b \equiv 0 \pmod{24}$, $c \equiv 0 \pmod{2}$ is realizable as the Chern numbers of an almost complex 6-manifold.*

Proof. Consider the complete intersection $V(f, g) \subset \mathbf{P}^5$ defined by the polynomials $f(z) = z_0^2 + z_1^2 + 2z_2^2 - z_3^2 - z_4^2 - 2z_5^2$, and $g(z) = z_0^4 + z_1^4 + 2z_2^4 - z_3^4 - z_4^4 - 2z_5^4$ [We]. $V(f, g)$ is a singular 3-fold with 90 ordinary double points, and every small resolution V of these nodes is a (not necessarily projective) Calabi-Yau 3-fold with Euler number 4. Suppose now that a prescribed triple $(a, b, c) \in \mathbf{Z}^{\oplus 3}$ is realized by a possibly disconnected almost complex manifold $X = \coprod_{i \in I} X_i$. If we form the connected sum

X' of the X_i , we obtain a connected almost complex manifold X' with Chern numbers $c_1^3 = a$, $c_1 c_2 = b$, but with $c_3 = c - 2(|I| - 1)$.

If $|I| > 1$ take the connected sum of X' with $|I| - 1$ copies of the complex manifold V . Since V is Calabi-Yau, the Chern numbers c_1^3 and $c_1 c_2$ remain unchanged, whereas the Euler number of $X' \#_{|I|-1} V$ becomes $c_3 = c$.

REMARK 11. The above argument has been suggested by F. Hirzebruch after talk at the MPI, in which one of us had sketched a less geometric proof of the proposition.

There is another question which is related to the result above: Fix a closed, oriented, 6-dimensional differentiable manifold X . Which pairs (a, b) of integers with $a \equiv 0 \pmod{2}$ and $b \equiv 0 \pmod{24}$ occur as Chern numbers c_1^3 and $c_1 c_2$ of almost complex structures on X , and in how many ways?

For manifolds with $b_2(X) = 1$ the Chern numbers determine the almost complex structure. For manifolds with $b_2 > 1$ this is no longer true. It is possible to construct infinitely many distinct almost complex structures with the same Chern numbers on a hypersurface of bidegree $(3, 3)$ in $\mathbf{P}^2 \times \mathbf{P}^2$.

An almost complex structure $[\tilde{t}_X]$ on a differentiable 6-manifold X is said to be integrable if \tilde{t}_X is homotopic to the classifying map of a complex 3-fold. We are not aware of any example of an almost complex 6-manifold which is known not to be integrable. On the other hand, it is also unknown whether or not the Chern numbers $c_1^3, c_1 c_2$ of integrable almost complex manifold are topological invariants. The following remark might therefore be of some interest:

PROPOSITION 10. *If the Chern numbers of complex 3-folds are topological invariants, then there exist almost complex structures which are not integrable.*

Proof. Consider a closed, oriented differentiable 6-manifold X without 2-torsion in $H^3(X, \mathbf{Z})$. Fix any almost complex structure on X with first Chern class $W \in H^2(X, \mathbf{Z})$.

Every element $x \in H^2(X, \mathbf{Z})$ defines a new almost complex structure on X with first Chern class $W + 2x$, and it is easy to see that these two almost complex structures have the same Chern numbers if and only if x satisfies the equations $p_1(X) \cdot x = 0$, and $3W^2 \cdot x + 6W \cdot x^2 + 4x^3 = 0$.

Suppose now (X, W) is integrable, $p_1(X) \neq 0$, and choose $x \in H^2(X, \mathbf{Z})$ such that $p_1(X) \cdot x \neq 0$. Then clearly, either none of the almost complex manifolds $(X, W + 2x)$ is integrable, or the Chern numbers of complex 3-folds are not topologically invariant.

REMARK 12. It is very likely that there exist non-integrable almost complex structures on manifolds X as above, but probably this is hard to prove. It is also not unlikely that the Chern numbers of complex 3-folds are not topological invariants. A possible way to check this would be, to run a computer search for 3-folds given by certain standard constructions.

4.2 STANDARD CONSTRUCTIONS

For later use we investigate the topological invariants of complex 3-folds which can be obtained by certain simple standard constructions like complete intersections, simple cyclic coverings, blow-ups of points and curves, and projective bundles.

PROPOSITION 11 (Libgober/Wood). *Let $X \subset \mathbf{P}^{3+r}$ be a smooth complete intersection of multidegree $\underline{d} = (d_1, \dots, d_r)$. Choose a normalized basis $e \in H^2(X, \mathbf{Z})$, and let $\varepsilon \in H^4(X, \mathbf{Z})$ be defined by $\varepsilon(e) = 1$. Then the invariants of X are:*

$$\begin{aligned} F_X(xe) &= dx^3 \quad \text{where } d = \prod_{i=1}^r d_i, \quad w_2(X) \equiv (4 + r - \sum_{i=1}^r d_i)e, \\ p_1(X) &= d(4 + r - \sum_{i=1}^r d_i^2)\varepsilon, \quad \text{and} \\ b_3(X) &= 4 - \frac{d}{6} \left[(4 + r - \sum_{i=1}^r d_i)^3 - 3(4 + r - \sum_{i=1}^r d_i)(4 + r - \sum_{i=1}^r d_i^2) \right. \\ &\quad \left. + 2(4 + r - \sum_{i=1}^r d_i^3) \right]. \end{aligned}$$

Proof. [L/W].

PROPOSITION 12. *Let X be a smooth, 1-connected, complex projective 3-fold, and let $\pi: X' \rightarrow X$ be a simple cyclic covering of degree d branched along a non-singular ample divisor $B \in |L^{\otimes d}|$. X' is smooth, projective, 1-connected, and $\pi^*: H^2(X, \mathbf{Z}) \rightarrow H^2(X', \mathbf{Z})$ is an isomorphism. The invariants of X and X' are related by the formulae:*

$$\begin{aligned} (\pi^*)^* F_{X'} &= dF_X, \quad w_2(X') - \pi^* w_2(X) \equiv (d-1)\pi^* c_1(L), \\ p_1(X') - \pi^* p_1(X) &= (1-d)(1+d)\pi^* c_1(L)^2, \quad \text{and} \\ b_3(X') &= db_3(X) + (d-1)(b_2(B) - 2b_2(X)). \end{aligned}$$

Proof. X' is clearly smooth and projective. By a theorem of M. Cornalba $\pi: X' \rightarrow X$ is a 3-equivalence, i.e. $\pi_*: \pi_i(X') \rightarrow \pi_i(X)$ is bijective for $i \leq 2$, and surjective for $i = 3$ [Co]. X' is therefore 1-connected, and $\pi^*: H^2(X, \mathbf{Z}) \rightarrow H^2(X', \mathbf{Z})$ is an isomorphism. The relation between $F_{X'}$ and F_X is obvious, whereas the formula for $b_3(X')$ follows from $\pi_1(B) = \{1\}$ and standard properties of Euler numbers.

In order to calculate $w_2(X')$ and $p_1(X')$ we compute the Chern classes of X' : $c_1(X') - \pi^*c_1(X) = (1-d)\pi^*c_1(L)$, $c_2(X') - \pi^*c_2(X) = (1-d)\pi^*[c_1(X)c_1(L) - dc_1(L)^2]$.

The latter formulae follow from the description of X' as a divisor in the total space of the line bundle L .

EXAMPLE 9. Let X be a d -fold, simple cyclic covering of \mathbf{P}^3 branched along a smooth surface $B \subset \mathbf{P}^3$ of degree dl , $l \geq 1$. Let $e \in H^2(X, \mathbf{Z})$ correspond to the preimage of a plane in \mathbf{P}^3 . The invariants of X are then given by:

$$F_X(xe) = dx^3, w_2(X) \equiv (4 + (1-d)l)e, p_1(X) = d[4 + (1-d)(1+d)l^2]\varepsilon \\ (\varepsilon(e) = 1), b_3(X) = (d-1)(d^2l^2 - 4dl + 6)dl.$$

PROPOSITION 13. Let $\sigma: \hat{X} \rightarrow X$ be the blow-up of a complex 3-fold X in a point, and let $e \in H^2(\hat{X}, \mathbf{Z})$ be the class of the exceptional divisor. The invariants of \hat{X} and X are related by the following formulae:

$$F_{\hat{X}}(\sigma^*h + xe) = F_X(h) + x^3 \quad \forall h \in H^2(X, \mathbf{Z}), x \in \mathbf{Z}, w_2(\hat{X}) = \sigma^*w_2(X), \\ p_1(\hat{X}) = \sigma^*p_1(X) + 4(e^2 - \sigma^*c_1(X) \cdot e), b_3(\hat{X}) = b_3(X).$$

Proof. Standard arguments, see [G/H]. The Chern classes are related by $c_1(\hat{X}) = \sigma^*c_1(X) - 2e$, $c_2(\hat{X}) = \sigma^*c_2(X)$.

PROPOSITION 14. Let $\sigma: \hat{X} \rightarrow X$ be the blow-up of a complex 3-fold X along a smooth curve C of genus g , and let $e \in H^2(\hat{X}, \mathbf{Z})$ be the class of the exceptional divisor. The invariants of \hat{X} and X are related by:

$$F_{\hat{X}}(\sigma^*h + xe) = F_X(h) - 3h \cdot Cx^2 - \deg N_{C/X}x^3 \quad \forall h \in H^2(X, \mathbf{Z}), \\ x \in \mathbf{Z}, w_2(\hat{X}) \equiv \sigma^*w_2(X) + e, p_1(\hat{X}) = \sigma^*p_1(X) + (e^2 - 2\sigma^*C), \\ b_3(\hat{X}) = b_3(X) + 2g.$$

Proof. [G/H]. The Chern classes are given by $c_1(\hat{X}) = \sigma^*c_1(X) - c$, $c_2(\hat{X}) = \sigma^*(c_2(X) + C) - \sigma^*c_1(X) \cdot e$.

PROPOSITION 15. Let E be a holomorphic vector bundle of rank 2 with Chern classes $c_i(E)$, $i = 1, 2$ over a 1-connected, compact complex surface Y , and let $\pi: \mathbf{P}(E) \rightarrow Y$ be the projective bundle of lines in the fibers of E . The cup-form of $\mathbf{P}(E)$ is given by

$$F_{\mathbf{P}(E)}(h + x\xi) = x[(3h^2) - (3c_1(E) \cdot h)x + (c_1(E)^2 - c_2(E))x^2],$$

where $\xi = c_1(\mathcal{O}_{\mathbf{P}(E)}(1))$, $h \in H^2(Y, \mathbf{Z})$, and $x \in \mathbf{Z}$. The other topological invariants of $\mathbf{P}(E)$ are:

$$\begin{aligned} w_2(\mathbf{P}(E)) &\equiv \pi^*(w_2(Y) + c_1(E)), p_1(E)) \\ &= \pi^*[c_1(Y)^2 - 2c_2(Y) + c_1(E)^2 - 4c_2(E)], b_3(\mathbf{P}(E)) = 0. \end{aligned}$$

Proof. The Leray-Hirsch theorem identifies the cohomology ring $H^*(\mathbf{P}(E), \mathbf{Z})$ with the ring $H^*(Y, \mathbf{Z})[\xi]/\langle \xi^2 + c_1(E) \cdot \xi + c_2(E) \rangle$; this determines the cup-form. In order to calculate the characteristic classes one uses the exact sequence $0 \rightarrow \mathcal{O}_{\mathbf{P}(E)} \rightarrow \pi^*E \otimes \mathcal{O}_{\mathbf{P}(E)}(1) \rightarrow T_{\mathbf{P}(E)} \rightarrow \pi^*T_Y \rightarrow 0$. $b_3(\mathbf{P}(E)) = 0$ follows from $b_1(Y) = 0$ and the Leray-Hirsch theorem.

4.3 EXAMPLES OF 1-CONNECTED NON-KÄHLERIAN 3-FOLDS

Recall that the Hessian of a symmetric trilinear form $F \in S^3 H^\vee$ on a free \mathbf{Z} -module H of finite rank was defined as the composition $H_F: H \xrightarrow{F^t} S^2 H^\vee \xrightarrow{\text{disc}} \mathbf{Z}$. In terms of coordinates ξ_1, \dots, ξ_b on H it is given by the determinant $\det\left(\frac{\partial^2 f}{\partial \xi_i \partial \xi_j}\right)$, where $f \in \mathbf{C}[H_{\mathbf{C}}]_3$ is the homogeneous cubic polynomial associated with F .

PROPOSITION 16. *Let F be a symmetric trilinear form whose Hessian vanishes identically. Then F is not realizable as cup-form of a Kählerian 3-fold.*

Proof. Let X be a complex 3-fold with a Kähler metric g . The Kähler class $[\omega_g] \in H^2(X, \mathbf{R})$ defines a multiplication map $\cdot [\omega_g]: H^2(X, \mathbf{R}) \rightarrow H^4(X, \mathbf{R})$, which is an isomorphism by the Hard Lefschetz Theorem [G/H]. In section 3.1 we have seen that this is not possible if the Hessian of the cup-form vanishes.

COROLLARY 6. *Cubic forms $f \in \mathbf{C}[H_{\mathbf{C}}]_3$ which depend on strictly less than $b = \text{rk}_{\mathbf{Z}} H$ variables are not realizable as cup-forms of Kählerian 3-folds with $b_2 = b$.*

By considering the Hessian of a cup-form over the reals one obtains further conditions.

DEFINITION 4. *Let $F \in S^3 H^\vee$ be a symmetric trilinear form on a free \mathbf{Z} -module of rank b .*

The Hesse cone of F is the subset $\mathcal{H}_F \subset H_{\mathbf{R}}$ defined by $\mathcal{H}_F := \{h \in H_{\mathbf{R}} \mid (-1)^b \det(F^t(h)) < 0\}$.

The index cone \mathcal{J}_F of F is the subset $\mathcal{J}_F := \{h \in \mathcal{H}_F \mid F^t(h) \in S^2 H_{\mathbf{R}}^\vee\}$ has signature $(1, -1, \dots, -1)$.

Clearly \mathcal{J}_F is an open subcone of \mathcal{H}_F which coincides with \mathcal{H}_F iff $b \leq 2$.

THEOREM 5. Let $F_X \in S^3 H^2(X, \mathbf{Z})^\vee$ be the cup-form of a smooth projective 3-fold with $h^{0,2}(X) = 0$. Then F_X has a non-empty index cone.

Proof. Let $h \in H^2(X, \mathbf{Z})$ be the dual class of a hyperplane section Y in some projective embedding. The inclusion $i: Y \hookrightarrow X$ induces a monomorphism $i^*: H^2(X, \mathbf{Z}) \rightarrow H^2(Y, \mathbf{Z})$ by the weak Lefschetz theorem. The symmetric bilinear form $F_X^t(h) \in S^2 H^2(X, \mathbf{Z})^\vee$ is simply the pull-back of the cup-form of Y under the inclusion i^* ; it is therefore non-degenerate by the Hard Lefschetz theorem [L]. Applying the Hodge index theorem to Y we see that the real bilinear form $F_X^t(h) \in S^2 H^2(X, \mathbf{R})^\vee$ must have one positive and $b - 1$ negative eigenvalues. In other words: $h \in I_{F_X}$.

REMARK 13. This result has two applications: it provides topological ‘upper bounds’ for the ample cone of a projective 3-fold with $h^{0,2} = 0$, and it gives further restrictions on symmetric trilinear forms to be realizable as cup-forms of projective 3-folds with $h^{0,2} = 0$ if $b \geq 4$.

These applications will be discussed in section 5.

We will now describe examples of 1-connected, non-Kählerian, complex 3-folds and determine their topological structure.

EXAMPLE 10 (Calabi-Eckmann). E. Calabi and B. Eckmann have defined complex structures X_τ , depending on a parameter τ , on the product $S^3 \times S^3[C/E]$. Their manifolds are principal fiber bundles over $\mathbf{P}^1 \times \mathbf{P}^1$ whose fiber and structure group is the elliptic curve $E_\tau = \mathbf{C}/\mathbf{Z} \oplus \mathbf{Z}\tau$, $\text{Im}(\tau) > 0$.

The Calabi-Eckmann manifolds are homogeneous, non-Kählerian 3-folds of algebraic dimension 2.

EXAMPLE 11 (Maeda). H. Maeda has generalized the Calabi-Eckmann construction. He constructed fiber bundles X'_τ over Hirzebruch surfaces \mathbf{F}_n , $n \geq 0$, whose fiber and structure group are an elliptic curve E_τ and $\text{Aut}(E_\tau)$ respectively [M]. X'_τ is again diffeomorphic to $S^3 \times S^3$, and therefore non-Kählerian. Maeda’s manifolds X'_τ are homogeneous if and only if $n = 0$ in which case they are Calabi-Eckmann 3-folds.

The Calabi-Eckmann construction can also be generalized in the following way:

Let $S^2 \tilde{\times} S^4$ be the non-trivial S^4 -bundle over S^2 , i.e. $S^2 \tilde{\times} S^4$ is the unique 1-connected, closed, oriented, differentiable 6-manifold with $H_2(S^2 \tilde{\times} S^4, \mathbf{Z}) \cong \mathbf{Z}$ and $b_3 = 0$, whose cup-form and Pontrjagin class vanish, but whose Stiefel-Whitney class w_2 is non-zero.

THEOREM 6. *For any integer $b \geq 0$ there exist compact complex 3-folds X_b , and X_b^- if $b \geq 1$, which are homeomorphic to $\#_b S^2 \times S^4 \#_{b+1} S^3 \times S^3$, and $S^2 \tilde{\times} S^4 \#_{b-1} S^2 \times S^4 \#_{b+1} S^3 \times S^3$.*

Proof. Let Y be a 1-connected, compact complex surface with $p_g(Y) = 0$ and $b_2(Y) \geq 2$, and let $E = \mathbf{C}/\Gamma$ be the elliptic curve associated to the lattice $\Gamma \subset \mathbf{C}$. We want to construct the required 3-folds as total spaces of principal E -bundles over Y . Let $c: H_2(Y, \mathbf{Z}) \rightarrow \Gamma$ be an arbitrary epimorphism. The corresponding cohomology class $c \in H^2(Y, \Gamma)$ defines a topological principal bundle over Y with fiber and structure group $E = \mathbf{C}/\Gamma$ as follows immediately from the identification of the classifying space $BE \simeq K(\Gamma, 2)$.

Let $\mathcal{O}_Y(E)$ be the sheaf of germs of holomorphic maps from Y to E . We have a short exact sequence $0 \rightarrow \Gamma \rightarrow \mathcal{O}_Y \rightarrow \mathcal{O}_Y(E) \rightarrow 0$ and a corresponding exact cohomology sequence

$$\rightarrow H^1(Y, \mathcal{O}_Y) \rightarrow H^1(Y, \mathcal{O}_Y(E)) \xrightarrow{\delta} H^2(Y, \Gamma) \rightarrow H^2(Y, \mathcal{O}_Y) \rightarrow$$

By our assumptions δ is an isomorphism, so that every topological principal E -bundle admits a holomorphic structure. Let X be the total space of such a bundle corresponding to a surjective map $c: H_2(Y, \mathbf{Z}) \rightarrow \Gamma$. The homotopy sequence of the fibration $p: X \rightarrow Y$ yields the sequence

$$0 \rightarrow \pi_2(X) \xrightarrow{p^*} \pi_2(Y) \rightarrow \pi_1(E) \rightarrow \pi_1(X) \xrightarrow{p^*} \pi_1(Y) \rightarrow 0.$$

Since Y is 1-connected, $\pi_2(Y)$ can be identified with $H_2(Y, \mathbf{Z})$, and then the boundary map $\pi_2(Y) \rightarrow \pi_1(E)$ becomes the characteristic map $c: H_2(Y, \mathbf{Z}) \rightarrow \Gamma$ of the bundle. This implies $\pi_1(X) = \{1\}$, whereas $H_2(X, \mathbf{Z})$ is given by: $0 \rightarrow H_2(X, \mathbf{Z}) \xrightarrow{p^*} H_2(Y, \mathbf{Z}) \xrightarrow{c} \Gamma \rightarrow 0$.

In particular, $H_2(X, \mathbf{Z})$ is free as a submodule of $H_2(Y, \mathbf{Z})$, and by dualizing the last sequence we obtain an identification (via p^*)

$$H^2(X, \mathbf{Z}) = H^2(Y, \mathbf{Z})/\Gamma^\vee.$$

The cup-form F_X of X is therefore trivial. In order to calculate $p_1(X)$ and $w_2(X)$, we use the exact sequence of tangent sheaves: $0 \rightarrow T_{X/Y} \rightarrow T_X$

$\rightarrow p^*T_Y \rightarrow 0$. Since $T_{X/Y}$ is a trivial bundle, the characteristic classes of X are simply the pullbacks of the corresponding classes of Y . But the map $p^*: H^4(Y, \mathbf{Z}) \rightarrow H^4(X, \mathbf{Z})$ is zero, since $\langle p^*(\varepsilon) \cup p^*(\alpha), [X] \rangle = \langle \varepsilon \cup \alpha, p_*[X] \rangle = 0$ for all classes $\varepsilon \in H^4(Y, \mathbf{Z})$, and $\alpha \in H^2(Y, \mathbf{Z})$.

Thus $p_1(X) = 0$, and $w_2(X)$ is the residue of $w_2(Y) \in H^2(Y, \mathbf{Z}_{/2})$ modulo $\Gamma^\vee / {}_2\Gamma^\vee$.

The Euler characteristic of X is zero, so that from $b_2(X) = b_2(Y) - 2$ we find $b_3(X) = 2(b_2(Y) - 1)$. The system of invariants associated to the manifold X is therefore given by

$$(b_2(Y) - 1, H^2(Y, \mathbf{Z}) / \Gamma^\vee, w_2(Y) \pmod{\Gamma^\vee / {}_2\Gamma^\vee}, 0, 0, 0),$$

i.e. X is diffeomorphic to

$$\#_{b_2(Y)-2} S^2 \times S^4 \#_{b_2(Y)-1} S^3 \times S^3 \text{ if } w_2(Y) \in \Gamma^\vee / {}_2\Gamma^\vee,$$

and to $S^2 \tilde{\times} S^4 \#_{b_2(Y)-3} S^2 \times S^4 \#_{b_2(Y)-1} S^3 \times S^3$ if $b_2(Y) \geq 3$, and $w_2(Y) \notin \Gamma^\vee / {}_2\Gamma^\vee$.

EXAMPLE 12 (Kato). In the two papers [K1], [K2] M. Kato studies the class of compact, complex 3-folds X containing smooth rational curves with neighborhoods biholomorphic to those of projective lines in \mathbf{P}^3 . On this class of 3-folds, called class L , he defines a semi-group structure $+$ with neutral element \mathbf{P}^3 .

Kato's connecting operation $+$ is defined by removing 'lines' $L_i \subset X_i$ from 3-folds $X_i, i = 1, 2$, and by identifying the complements $X_i \setminus L_i$ along open sets $U_i \setminus L_i$ obtained from suitable neighborhoods $U_i \subset X_i$.

Starting with a certain elliptic fiber space X_1 over the blow-up of $\mathbf{P}^1 \times \mathbf{P}^1$ in a point, he constructs a sequence of 3-folds $X_n := X_1 + X_{n-1}, n \geq 2$. The 3-folds X_n are 1-connected spin-manifolds with $H_2(X_n, \mathbf{Z}) = \mathbf{Z}$. Their cup-forms F_{X_n} , and their Pontrjagin classes $p_1(X_n)$ are in terms of a (normalized) generator $e_n \in H^2(X_n, \mathbf{Z})$ and its dual class $\varepsilon_n \in H^4(X_n, \mathbf{Z})$ given by $F_{X_n}(xe_n) = (n-1)x^3$, and $p_1(X_n) = 4(n-1)\varepsilon_n$ ($\varepsilon_n(e_n) = 1$). The third Betti-number of X_n is $4n$.

In particular, X_1 is diffeomorphic to $S^2 \times S^4 \#_2 S^3 \times S^3$, and X_2 is diffeomorphic to $\mathbf{P}^3 \#_4 S^3 \times S^3$. It is interesting to note that the Chern-numbers $c_1^3, c_1 c_2$ of the X_n are $c_1^3 = 64(1-n)$, $c_1 c_2 = 24(1-n)$, i.e. they satisfy $8c_1 c_2 = 3c_1^3$. For projective manifolds of general type this equality is characteristic for ball quotients [Y].

EXAMPLE 13 (Twistor spaces). Let $p: Z \rightarrow M$ be the twistor fibration of a closed, oriented Riemannian 4-manifold (M, g) . Z carries a natural almost complex structure which is integrable if and only if g is self-dual [A/H/S].

Examples of 1-connected 4-manifolds which admit self-dual structures are S^4 , $\#_n \mathbf{P}^2$, and $K3$ -surfaces.

The total spaces of their twistor fibrations are 1-connected complex 3-folds which may be Moishezon for S^4 and $\#_n \mathbf{P}^2$ [C], but which are usually non-Kähler [Hi]. We leave it to the reader to calculate the topological invariants of these 3-folds. There is an interesting relation between Twistor spaces of connected sums and Kato's connection operation $+$ for class L manifolds [K2], [D/F].

EXAMPLE 14 (Oguiso). In a recent preprint [O1] K. Oguiso constructs examples of 1-connected, Moishezon Calabi-Yau 3-folds with very interesting cup-forms. He proves that for every integer $d \geq 1$ there exists a smooth complete intersection X'_d of type $(2, 4)$ in \mathbf{P}^5 which contains a non-singular rational curve C_d of degree d with normal bundle $N_{C_d/X_d} = \mathcal{O}_{C_d}(-1)^{\oplus 2}$.

The 3-fold X'_d can now be flopped along C_d , i.e. C_d can be blown up to $\mathbf{P}(N_{C_d/X_d}) \cong \mathbf{P}^1 \times \mathbf{P}^1$, and then 'blown down in the other direction'. The resulting 3-fold X_d is a 1-connected Moishezon manifold with trivial canonical bundle and cup-form F_{X_d} given by $F_{X_d}(xe_d) = (d^3 - 8)x^3$. Here $e_d \in H^2(X_d, \mathbf{Z})$ is the normalized generator corresponding to the strict transform of the negative of a hyperplane section of X'_d . The Pontrjagin class of X_d is $p_1(X_d) = (112 + 4d)\varepsilon_d$ where $\varepsilon_d \in H^4(X_d, \mathbf{Z})$ denotes the generator with $\varepsilon_d(e_d) = 1$. Since the Euler-number does not change under a flop we have $b_3(X_d) = 180$ for every d .

5. COMPLEX 3-FOLDS WITH SMALL b_2

In this section we investigate the following natural problem: Which cubic forms can be realized as cup-forms of compact complex 3-folds? For small b_2 something can be said: Any core of a 1-connected, closed, oriented differentiable 6-manifold with $H_2(X, \mathbf{Z}) \cong \mathbf{Z}$ is homotopy equivalent to the core of a 1-connected complex 3-fold. In the case $b_2 = 2$, at least every discriminant Δ is realizable by a complex manifold. If $b_2 = 3$ we can realize all types of complex cubics with one exception, the union of a smooth conic and a tangent line. In addition to these realization results we prove a finiteness