

3. The τ_I -spherical functions

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **45 (1999)**

Heft 3-4: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **20.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

3. THE τ_l -SPHERICAL FUNCTIONS

Let $\mathfrak{U}(\mathfrak{g})$ be the universal enveloping algebra of the complexification $\mathfrak{g}_{\mathbb{C}}$ of $\mathfrak{g} = \mathfrak{sp}(1, n)$. The elements of $\mathfrak{U}(\mathfrak{g})$ are considered as left-invariant differential operators on $G = \mathrm{Sp}(1, n)$ acting on C^∞ functions on the right:

$$f(g; X) := \left. \frac{d}{dt} f(g \exp tX) \right|_{t=0} \quad (f \in C^\infty(G), X \in \mathfrak{g}, g \in G).$$

We adopt Harish-Chandra's notation $f(g; D)$ for the image of $f \in C^\infty(G)$ under the right action of $D \in \mathfrak{U}(\mathfrak{g})$. The set of K -invariant elements of $\mathfrak{U}(\mathfrak{g})$ is denoted by $\mathfrak{U}(\mathfrak{g})^K$.

3.1. THEOREM ([Go], Theorems 8, 10 and 14; [GaV], Theorems 1.3.14, 1.4.5 and Proposition 1.4.4). *Let $l \in \mathbf{N}/2$ be fixed. Let ζ be a complex-valued continuous function on G satisfying (2.9), (2.10) and $\zeta(e) = 1$ (e is the unit element of G). The following statements are mutually equivalent.*

1. *The mapping $f \mapsto \int_G f(g) \zeta(g) dg$ is an algebra homomorphism of $\mathcal{D}(G; \chi_l)$ into \mathbb{C} .*

2. *ζ satisfies the functional equation*

$$(3.15) \quad \int_K \zeta(kg_1k^{-1}g_2) dk = \zeta(g_1)\zeta(g_2)$$

for all $g_1, g_2 \in G$.

3. *ζ is a common eigenfunction of the elements of $\mathfrak{U}(\mathfrak{g})^K$.*

A function ζ satisfying the equivalent conditions of Theorem 3.1 is called a *spherical function of type τ_l (and height 1)* or briefly a *τ_l -spherical function*¹⁾.

Observe that Condition 3 implies in particular that every τ_l -spherical function is analytic on G because $\mathfrak{U}(\mathfrak{g})^K$ contains an elliptic differential operator.

¹⁾ Our definition of τ_l -spherical function ζ is formally less restrictive than Godement's, which also requires ζ to be quasi-bounded with respect to some seminorm on G (cf. [Go], p. 519). It is possible to show (e.g. [GaV], Theorem 1.3.14) that each of the conditions for ζ given by Theorem 3.1 is equivalent to the existence of an irreducible Fréchet representation (T, \mathcal{H}) – with a d_l -dimensional K -isotypic subspace $\mathcal{H}(\tau_l)$ of type τ_l – for which

$$(*) \quad \zeta = \frac{1}{d_l} \mathrm{tr}[E(\tau_l)TE(\tau_l)],$$

$E(\tau_l)$ being the projection of \mathcal{H} onto $\mathcal{H}(\tau_l)$. (T, \mathcal{H}) can be chosen to be a Banach representation of G if and only if ζ is quasi-bounded with respect to some seminorm on G . In Section 5 we will determine, for each τ_l -spherical function ζ , an irreducible *Hilbert* representation (T, \mathcal{H}) for which $(*)$ holds. It follows, in particular, that the condition of quasi-boundedness is, in our case, automatically satisfied.

For complex-valued functions f on G and F on \mathbf{R} , we set

$$\begin{aligned} f^*(g) &= \overline{f(g^{-1})} & (g \in G) \\ F^*(t) &= \overline{F(-t)} & (t \in \mathbf{R}). \end{aligned}$$

The τ_l -Abel transform of $f \in \mathcal{D}(G; \chi_l)$ is the function $\mathcal{A}_l f$ on \mathbf{R} defined by

$$(3.16) \quad \mathcal{A}_l f(t) = \frac{1}{d_l^2} e^{\rho t} \int_N f(a_t n) dn.$$

Its properties are summarized in the following proposition.

3.2. PROPOSITION. *For all $f \in \mathcal{D}(G; \chi_l)$, $\mathcal{A}_l f$ is a C^∞ function on \mathbf{R} with compact support. If $f, f_1, f_2 \in \mathcal{D}(G; \chi_l)$ and $a_1, a_2 \in \mathbf{C}$, then*

$$(3.17) \quad (\mathcal{A}_l f)^* = \mathcal{A}_l(f^*),$$

$$(3.18) \quad \mathcal{A}_l(a_1 f_1 + a_2 f_2) = a_1 \mathcal{A}_l f_1 + a_2 \mathcal{A}_l f_2,$$

$$(3.19) \quad \mathcal{A}_l(f_1 * f_2) = \mathcal{A}_l f_1 * \mathcal{A}_l f_2.$$

Formula (3.17) is equivalent to the fact that $\mathcal{A}_l f$ is an even function.

Proof. Formulas (3.17)–(3.19) are immediately proven by passing to $\mathcal{D}(G; \tau_l)$. For the last statement, recall that $f(g^{-1}) = f(g)$ for $f \in \mathcal{D}(G; \chi_l)$. \square

The following lemma relates our definition of \mathcal{A}_l to the definition often found in the literature (cf. e.g. [W2], p. 34).

3.3. LEMMA. *For $f \in \mathcal{D}(G; \chi_l)$ one has*

$$\int_N f(ka_t n) dn = \frac{1}{d_l} \chi_l(k) \int_N f(a_t n) dn.$$

Proof. Let $F \in \mathcal{D}(G; \tau_l)$. Then $\int_N F(a_t n) dn$ commutes with $\tau(m)$ ($m \in M$) so with $\tau(k_1)$ ($k_1 \in K_1$), hence is a scalar multiple of the identity. The lemma follows by taking traces. \square

We now use the τ_l -Abel transform to construct τ_l -spherical functions. Because of Proposition 3.2, for any complex number s , the map

$$(3.20) \quad \lambda_s : f \longmapsto \int_{-\infty}^{\infty} \mathcal{A}_l f(t) e^{-st} dt$$

is an algebra homomorphism of $\mathcal{D}(G; \chi_l)$ into \mathbf{C} .

Set

$$(3.21) \quad \alpha_{l,s}(ka_t n) = \frac{1}{d_l} \chi_l(k) e^{-(s+\rho)t}.$$

Since $f = f * d_l \chi_l$ and $\chi_l(k^{-1}) = \chi_l(k)$ for $k \in K$, for every $f \in \mathcal{D}(G; \chi_l)$

$$(3.22) \quad \begin{aligned} \lambda_s(f) &= \frac{1}{d_l} \int_K \int_{-\infty}^{\infty} \int_N f(ka_t n) \chi_l(k) e^{(-s+\rho)t} dk dt dn \\ &= \int_G f(g) \alpha_{l,s}(g) dg \\ &= \int_G f(g) \int_K \alpha_{l,s}(kgk^{-1}) dk dg \\ &= \int_G f(g) \zeta_{l,s}(g) dg \end{aligned}$$

with

$$(3.23) \quad \zeta_{l,s} := \int_K \alpha_{l,s}(kgk^{-1}) dk.$$

One easily checks that $\zeta_{l,s}$ satisfies $\zeta_{l,s} = \zeta_{l,s}^0$, $\zeta_{l,s} * d_l \chi_l = \zeta_{l,s}$ and $\zeta_{l,s}(e) = 1$. Thus $\zeta_{l,s}$ is a τ_l -spherical function. It will be shown in the next section that any τ_l -spherical function is of the form (3.24).

By Remark 2.3, we have

$$(3.24) \quad \zeta_{l,s}(g) = \frac{1}{d_l} \chi_l(k_1) \zeta_{l,s}(a_t) \quad \text{for } g = k_1 k_2 a_t k_2',$$

so $\zeta_{l,s}$ is uniquely determined by its restriction to A .

4. THE DIFFERENTIAL EQUATION FOR THE τ_l -SPHERICAL FUNCTIONS

For a subalgebra \mathfrak{u} of \mathfrak{g} , let $\mathfrak{u}_{\mathbf{C}}$ denote the complex subalgebra of $\mathfrak{g}_{\mathbf{C}}$ generated by \mathfrak{u} . The universal enveloping algebra $\mathfrak{U}(\mathfrak{u})$ of $\mathfrak{u}_{\mathbf{C}}$ is considered as a subalgebra of $\mathfrak{U}(\mathfrak{g})$.

The representation τ_l of K_1 induces differentiated representations of the Lie algebra \mathfrak{k}_1 of K_1 and of the universal enveloping algebra $\mathfrak{U}(\mathfrak{k}_1)$ of $(\mathfrak{k}_1)_{\mathbf{C}}$. We indicate these representations with the same letter τ_l . Let \mathfrak{k}_2 be the Lie algebra of K_2 . Every element $Y \in \mathfrak{k}_{\mathbf{C}}$ can be uniquely decomposed as