

## 2. Hyperbolic geometry and Fuchsian groups

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **45 (1999)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.09.2024**

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

### **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

canonical polygons. Section 4 provides the necessary material from hyperbolic trigonometry, it contains also some lemmas needed later. Section 5 contains the proof of the main theorem and Section 6 gives some applications, mainly concerning hyperelliptic Riemann surfaces. More precisely, I give a new proof of a geometric characterization of hyperelliptic Riemann surfaces which first appeared in [14] (I thank very much Feng Luo who, by his comments on [14], has contributed to the idea of this new proof). I also show (and this is a new result) that the Teichmüller space  $T_g$  for  $g = 2$  can be parametrized by 7 geodesic length functions, taken as homogeneous parameters. This is the optimum parametrization of Teichmüller space by geodesic length functions which one can expect.

I spoke about the content of this paper in lectures of the Troisième Cycle Romand de Mathématiques (Lausanne 1997); I thank the participants for their comments.

## 2. HYPERBOLIC GEOMETRY AND FUCHSIAN GROUPS

The material of this section and of parts of the following section is standard, see for example [1], [4], [5], [6], [7], [8], [15].

DEFINITION. (i)  $\mathbf{H} = \{z = (x, y) \in \mathbf{C} : y > 0\}$  denotes the *upper halfplane*. The *hyperbolic metric* on  $\mathbf{H}$  is given by

$$dz = \frac{1}{y}(dz)_E$$

where  $(dz)_E$  is the standard Euclidean metric on  $\mathbf{C}$  and  $y$  is the imaginary part of  $z$ .

(ii) Define

$$\mathrm{SL}(2, \mathbf{R}) = \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} : ad - bc = 1; a, b, c, d \in \mathbf{R} \right\}$$

and

$$\mathrm{PSL}(2, \mathbf{R}) = \mathrm{SL}(2, \mathbf{R})/\sim$$

with  $A \sim B$  if and only if  $A = \pm B$  for  $A, B \in \mathrm{SL}(2, \mathbf{R})$ . Let  $\gamma \in \mathrm{SL}(2, \mathbf{R})$ ,

$$\gamma = \begin{bmatrix} a & b \\ c & d \end{bmatrix}.$$

Then the action of  $\gamma$  on  $\mathbf{H}$  is defined as

$$\gamma(z) = \frac{az + b}{cz + d}$$

for  $z \in \mathbf{H}$ .

**THEOREM 1.**  $\mathbf{H}$  is a complete Riemannian manifold of constant curvature  $-1$ . The geodesics in  $\mathbf{H}$  are either Euclidean semicircles which are orthogonal to the real axis or vertical half-lines.

**THEOREM 2.**

(i)  $\text{PSL}(2, \mathbf{R}) = \text{Isom}^+(\mathbf{H})$ , the group of orientation preserving isometries of  $\mathbf{H}$ .

(ii) Let  $u$  and  $v$  be geodesics in  $\mathbf{H}$ , let  $z$  be on  $u$  and  $z'$  on  $v$ . Then there exists  $\gamma \in \text{PSL}(2, \mathbf{R})$  with  $\gamma(u) = v$  and  $\gamma(z) = z'$ .

**DEFINITION.** For a measurable subset  $G \subset \mathbf{H}$  define the volume  $\text{vol}(G)$  as

$$\text{vol}(G) = \int_G \frac{dx dy}{y^2}.$$

**REMARK.** The volume is invariant under  $\gamma \in \text{SL}(2, \mathbf{R})$ .

**CONVENTIONS.** (i) Speaking of triangles, quadrilaterals and polygons always means that the sides are hyperbolic geodesic segments in  $\mathbf{H}$ .

(ii) Speaking of *angles* in triangles, quadrilaterals and polygons always means *interior angles*.

**THEOREM 3.** The volume of a polygon with angles  $\alpha_i$ ,  $i = 1, 2, \dots, m$ ,  $m \geq 3$ , is

$$(m - 2)\pi - \sum_{i=1}^m \alpha_i.$$

**DEFINITION.** A Fuchsian group  $\Gamma$  is a discrete subgroup of  $\text{PSL}(2, \mathbf{R})$  where discrete means that the identity matrix is not a cluster point in  $\Gamma$  with respect to the topology induced by the standard topology of  $\mathbf{R}^4$ .

**THEOREM 4.** *Let  $\Gamma$  be a Fuchsian group without elliptic elements (an element  $\gamma \in \text{PSL}(2, \mathbf{R})$  is elliptic if  $|\text{tr}(\gamma)| < 2$  where  $\text{tr}$  is the trace). Then  $\mathbf{H}/\Gamma$  is a complete connected orientable Riemannian manifold of dimension 2 with a metric of constant curvature  $-1$ .*

**DEFINITION.** *A hyperbolic surface is a connected orientable manifold  $M = \mathbf{H}/\Gamma$  as in Theorem 4 (where  $\Gamma$  is a Fuchsian group without elliptic elements).  $M$  is called closed if  $M$  is compact and has no boundary.*

3. FUNDAMENTAL DOMAINS AND CANONICAL POLYGONS

**DEFINITION** (Compare Figure 2). Let  $g \geq 2$  be an integer. A *canonical polygon*  $P(g)$  is a polygon with  $4g$  sides, denoted by  $a_1, \dots, a_{4g}$ , ordered clockwise, and angles  $\alpha_i$  between  $a_i$  and  $a_{i+1}$ ,  $i = 1, \dots, 4g$  (indices are taken modulo  $4g$ ), such that

- (I)  $a_i$  and  $a_{i+2g}$  have the same length,  $i = 1, \dots, 2g$ ;
- (II) the sum of the angles of  $P(g)$  is  $2\pi$ ;
- (III)  $0 < \alpha_i < \pi$ ,  $i = 1, \dots, 4g$ ;
- (IV)  $\alpha_1 = \alpha_{2g+1}$ ;
- (V)  $\sum_{i=1}^g \alpha_{2i-1} + \sum_{i=g+1}^{2g} \alpha_{2i} = \sum_{i=1}^g \alpha_{2i} + \sum_{i=g+1}^{2g} \alpha_{2i-1}$ .

I shall speak of condition (I) (or (II) or (III) or (IV) or (V) ) referring to this definition.




FIGURE 2  
A canonical polygon  $P(g)$  for  $g = 2$