

4. On K-homology

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **49 (2003)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **26.04.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

For R a fixed set of representatives for G/H , the map

$$\begin{aligned}\varphi_R: \text{Ind}_H^G(S_{\tilde{D}}) &\rightarrow S_{\tilde{D}} \\ f &\mapsto \{f(r)\}_{r \in R}\end{aligned}$$

is well-defined by H -equivariance of the elements of $S_{\tilde{D}}$ and one checks that it defines a G -equivariant isometric bijection. Similarly for the adjoint operators.

The following example is a particular case of the previous lemma.

EXAMPLE 3.2. Let us look at the case $\tilde{M} = M \times G$. A section $\tilde{s} \in C_c^\infty(\tilde{M}, \pi^*E)$ is an element $\tilde{s} = \{s_g\}_{g \in G}$ where $s_g \in C^\infty(M, E)$ and $s_g = 0$ for all but finitely many g 's. Note that $L^2(\tilde{M}, \pi^*E)$ can be identified with $\ell^2(G) \otimes L^2(M, E)$. Now

$$\tilde{D}\tilde{s} = \{Ds_g\}_{g \in G} \in C_c^\infty(\tilde{M}, \pi^*F)$$

and hence $S_{\tilde{D}}$ may be identified with $\ell^2(G) \otimes S_D \cong \ell^2(G)^d$, where $d = \dim_{\mathbb{C}}(S_D)$. In this identification the projection P onto $S_{\tilde{D}}$ becomes the identity in $M_d(\mathcal{N}(G))$ and thus

$$\dim_G(S_{\tilde{D}}) = \sum_{i=1}^d \langle e, e \rangle = d = \dim_{\mathbb{C}}(S_D).$$

A similar argument for D^* shows that in this case not only does the L^2 -Index of $\tilde{D}$ coincide with the Index of D , but also the individual terms of the difference correspond to each other. This is not the case in general, see Example 2.2.

4. ON K -HOMOLOGY

Many ideas of this section go back to the seminal article by Baum and Connes [3], which has been circulating for many years and has only recently been published.

An elliptic pseudo-differential operator D on the closed manifold M can also be used to define an element $[D] \in K_0(M)$, the K -homology of M , and according to Baum and Douglas [4], all elements of $K_0(M)$ are of the form $[D]$. The index defined in Section 2 extends to a well-defined

homomorphism (cf. [4])

$$\text{Index}: K_0(M) \rightarrow \mathbf{Z},$$

such that $\text{Index}([D]) = \text{Index}(D)$. On the other hand, the projection $\text{pr}: M \rightarrow \{pt\}$ induces, after identifying $K_0(\{pt\})$ with $\mathbf{Z}$, a homomorphism

$$(*) \quad \text{pr}_*: K_0(M) \rightarrow \mathbf{Z},$$

which, as explained in [4], satisfies

$$\text{pr}_*([D]) = \text{Index}([D]).$$

More generally (cf. [4]), for a not necessarily finite CW-complex X , every $x \in K_0(X)$ is of the form $f_*[D]$ for some $f: M \rightarrow X$, and $K_0(X)$ is obtained as a colimit over $K_0(M_\alpha)$, where the M_α form a directed system consisting of closed Riemannian manifolds (these homology groups $K_0(X)$ are naturally isomorphic to the ones defined using the Bott spectrum; sometimes, they are referred to as K -homology groups with *compact supports*). The index map from above extends to a homomorphism

$$\text{Index}: K_0(X) \rightarrow \mathbf{Z},$$

such that $\text{Index}(x) = \text{Index}([D])$ if $x = f_*[D]$, with $f: M \rightarrow X$.

We now consider the case of $X = BG$, the classifying space of the discrete group G , and obtain thus for any $f: M \rightarrow BG$ a commutative diagram

$$\begin{array}{ccc} K_0(M) & \xrightarrow{\text{Index}} & \mathbf{Z} \\ f_* \downarrow & & \parallel \\ K_0(BG) & \xrightarrow{\text{Index}} & \mathbf{Z}. \end{array}$$

Note that (*) from above implies the following naturality property for the index homomorphism.

LEMMA 4.1. *For any homomorphism $\varphi: H \rightarrow G$ one has a commutative diagram*

$$\begin{array}{ccc} K_0(BH) & \xrightarrow{\text{Index}} & \mathbf{Z} \\ (B\varphi)_* \downarrow & & \parallel \\ K_0(BG) & \xrightarrow{\text{Index}} & \mathbf{Z}. \quad \square \end{array}$$

We now turn to the L^2 -index of Section 2. It extends to a homomorphism

$$\text{Index}_G: K_0(BG) \rightarrow \mathbf{R}$$

as follows. Each $x \in K_0(BG)$ is of the form $f_*(y)$ for some $y = [D] \in K_0(M)$, $f: M \rightarrow BG$, M a closed smooth manifold and D an elliptic operator on M . Let $\tilde{D}$ be the lifted operator to $\tilde{M}$, the G -covering space induced by $f: M \rightarrow BG$. Then put

$$\text{Index}_G(x) := \text{Index}_G(\tilde{D}).$$

One checks that $\text{Index}_G(x)$ is indeed well-defined, either by direct computation, or by identifying it with $\tau(x)$, where τ denotes the composite of the assembly map $K_0(BG) \rightarrow K_0(C_r^*G)$ with the natural trace $K_0(C_r^*G) \rightarrow \mathbf{R}$ (for this latter point of view, see Higson-Roe [10]; for a discussion of the assembly map see e.g. Kasparov [12], or Valette [14]). The following naturality property of this index map is a consequence of Lemma 3.1.

LEMMA 4.2. *For $H < G$ the following diagram commutes:*

$$\begin{array}{ccc} K_0(BH) & \xrightarrow{\text{Index}_H} & \mathbf{R} \\ \downarrow & & \parallel \\ K_0(BG) & \xrightarrow{\text{Index}_G} & \mathbf{R}. \quad \square \end{array}$$

Atiyah's L^2 -Index Theorem 2.1 for a given G can now be expressed as the statement (as already observed in [10])

$$\text{Index}_G = \text{Index}: K_0(BG) \rightarrow \mathbf{R}.$$

5. ALGEBRAIC PROOF OF ATIYAH'S L^2 -INDEX THEOREM

Recall that a group A is said to be *acyclic* if $H_*(BA, \mathbf{Z}) = 0$ for $* > 0$. For G a countable group, there exists an embedding $G \rightarrow A_G$ into a countable acyclic group A_G . There are many constructions of such a group A_G available in the literature, see for instance Kan-Thurston [11, Proposition 3.5], Berrick-Varadarajan [5] or Berrick-Chatterji-Mislin [6]; these different constructions are to be compared in Berrick's forthcoming work [7]. It follows that the suspension ΣBA_G is contractible, and therefore the inclusion $\{e\} \rightarrow A_G$