

**Zeitschrift:** Helvetica Physica Acta  
**Band:** 72 (1999)  
**Heft:** 1

**Artikel:** Algebraic localisation of linear response in networks with algebraically decaying interaction, and application to discrete breathers in dipole-dipole systems  
**Autor:** Baesens, Claude / MacKay, Robert S.  
**DOI:** <https://doi.org/10.5169/seals-117167>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 03.07.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

# Algebraic localisation of linear response in networks with algebraically decaying interaction, and application to discrete breathers in dipole–dipole systems

By Claude Baesens<sup>1</sup> and Robert S. MacKay

Nonlinear Centre, DAMTP, Silver Street, Cambridge CB3 9EW, United Kingdom

(19.II.1998)

*Abstract.* Networks of units for which the coupling decreases algebraically with distance between them are studied. It is proved that the linear response to an algebraically localised force is algebraically localised if the response exists and is unique. Applications are given to equilibria of networks of bistable units and to breathers in networks of oscillating dipoles.

## 1 Introduction

The response of a network with exponentially decaying interaction to an exponentially localised force is itself exponentially decaying, provided the response exists and is unique (by response we always mean bounded response) (for a proof in a general setting, see [1]). Many systems of physical interest, however, have algebraically decaying interaction. For example, the potential energy of dipole–dipole interaction decays like the inverse cube of their separation, and that of van der Waals forces (due to correlations in dipole moment fluctuations) like the inverse sixth power. Is there an analogous result for algebraically decaying interaction? In this paper we give an affirmative answer.

Our interest in this problem was stimulated by work on “discrete breathers” in a monolayer of carbon monoxide molecules adsorbed on the surface of ruthenium [2]. Discrete breathers are spatially localised time-periodic vibrations of a network of oscillators. In this

---

<sup>1</sup>On leave from Centre de Dynamique des Systèmes Complexes, Université de Bourgogne, Dijon, France.

case, the oscillators are the  $C - O$  stretches and they are considered to be coupled by dipole-dipole interaction, the ruthenium substrate being regarded as rigid. The strategy of [3] can be easily adapted to the space  $\ell_1$  of spatially summable perturbations from equilibrium and hence to prove existence of  $\ell_1$ -breathers, i.e. whose oscillation amplitudes are summable over space, which implies spatial decay to equilibrium but with no particular rate (as the decay could be lacunary). The question arises whether a better spatial decay result can be proved.

In this paper, we prove algebraic upper bounds for the linear response to an algebraically localised force for a network with algebraically decaying coupling (provided the response exists and is unique), give a general application to continuation problems, and apply it to the dipole-dipole breather problem. We conclude by assessing how close our results are to optimal.

## 2 Basic result

Let  $(S, d)$  be a countable metric space. For each  $s \in S$ , let  $X_s$  and  $Y_s$  be Banach spaces, with norms  $|\cdot|_{X_s}$ ,  $|\cdot|_{Y_s}$ , respectively. Let  $X = \{x \in \times_{s \in S} X_s : \|x\|_X < \infty\}$  for some norm  $\|\cdot\|_X$  based on  $(|\cdot|_{X_s})_{s \in S}$ , and suppose it is complete. Define  $Y$  similarly. We will drop the subscripts on the symbols for the norms when the context makes them clear. Let  $L : X \rightarrow Y$  be a bounded linear map of matrix type<sup>2</sup>, i.e. one which can be written as

$$(Lx)_r = \sum_{s \in S} L_{rs} x_s,$$

with respect to components in  $\times_{r \in S} Y_r$  and  $\times_{s \in S} X_s$ . Let  $\alpha \geq 0$ .

**Definition:** A linear map  $L : X \rightarrow Y$  of matrix type is said to be  $\alpha$ -algebraically local if there exists  $\zeta > 0$  such that the modified operator  $L^\zeta$  defined by

$$L_{rs}^\zeta = (1 + \zeta d(r, s))^\alpha L_{rs}$$

is a bounded operator from  $X$  to  $Y$ .

**Definition:** A configuration  $y \in Y$  is said to be  $\alpha$ -algebraically localised about site  $o \in S$ , if there exists  $\zeta > 0$  such that  $y^\zeta$ , defined by

$$y_s^\zeta = (1 + \zeta d(o, s))^\alpha y_s,$$

is also in  $Y$ . Similarly for  $x \in X$ .

Note that if  $L^\zeta$  and  $y^\zeta$  are bounded for some  $\zeta > 0$  then so are  $L^z$  and  $y^z$  for all  $z \geq 0$ : for  $z < \zeta$  use  $1 + zd \leq 1 + \zeta d$  (recall  $d \geq 0$ ), and for  $z > \zeta$  use  $1 + zd \leq (z/\zeta)(1 + \zeta d)$ .

---

<sup>2</sup>We are grateful to Mark Johnston for pointing out that we should have stated this assumption explicitly in [1] also.

**Theorem 1** *If  $L : X \rightarrow Y$  is  $\alpha$ -algebraically local,  $y \in Y$  is  $\alpha$ -algebraically localised about  $o \in S$ ,  $L$  is invertible, and  $x = L^{-1}y$ , then  $x$  is  $\alpha$ -algebraically localised about  $o \in S$ . In fact,  $\|L^z - L\| < \|L^{-1}\|^{-1}$  for  $z$  small enough, and for such  $z$ ,*

$$\|x^z\| \leq \frac{\|y^z\|}{\|L^{-1}\|^{-1} - \|L^z - L\|}.$$

**Lemma 1** *If  $L$  is  $\alpha$ -algebraically local then  $L^z$  depends continuously on  $z \in \mathcal{R}_+$ .*

**Proof:** We treat separately the two cases  $\alpha \geq 1$  and  $0 \leq \alpha < 1$ .

Suppose  $\alpha \geq 1$ . If  $0 \leq p < q$  then

$$q^\alpha - p^\alpha \leq (q - p)\alpha q^{\alpha-1}.$$

Given  $0 \leq a < b$ , choose  $c \geq b$ . Given  $r, s \in S$ , write  $d$  for  $d(r, s)$ . Then

$$\begin{aligned} |L_{rs}^b - L_{rs}^a| &= \left[ \left( \frac{1+bd}{1+cd} \right)^\alpha - \left( \frac{1+ad}{1+cd} \right)^\alpha \right] |L_{rs}^c| \\ &\leq (b-a)d\alpha \frac{(1+bd)^{\alpha-1}}{(1+cd)^\alpha} |L_{rs}^c| \\ &\leq \frac{b-a}{c} \alpha |L_{rs}^c|. \end{aligned}$$

Thus

$$\|L^b - L^a\| \leq \frac{b-a}{c} \alpha \|L^c\|.$$

If  $0 \leq \alpha < 1$  and  $0 \leq p < q$  then  $(1+q)^\alpha - (1+p)^\alpha \leq q^\alpha - p^\alpha$ . Given  $0 \leq a < b$ , choose  $c \geq b$ . Then

$$\begin{aligned} |L_{rs}^b - L_{rs}^a| &\leq \frac{(b^\alpha - a^\alpha) d^\alpha}{(1+cd)^\alpha} |L_{rs}^c| \\ &\leq \frac{b^\alpha - a^\alpha}{c^\alpha} |L_{rs}^c|. \end{aligned}$$

Thus

$$\|L^b - L^a\| \leq \frac{b^\alpha - a^\alpha}{c^\alpha} \|L^c\|.$$

□

**Lemma 2** *If  $o, r, s \in S$  and  $z \geq 0$ , then  $\frac{1+z d(r,o)}{1+z d(s,o)} \leq 1 + z d(r, s)$ .*

**Proof:** We treat separately the two cases  $d(r, o) \geq d(s, o)$  and  $d(r, o) < d(s, o)$ .

If  $d(r, o) \geq d(s, o)$  then since  $d$  is a metric,  $d(r, s) \geq d(r, o) - d(s, o)$ , and so

$$1 + z d(r, s) \geq 1 + z d(r, o) - z d(s, o).$$

It follows that

$$\begin{aligned} (1 + z d(s, o))(1 + z d(r, s)) &\geq (1 + z d(s, o))(1 + z d(r, o) - z d(s, o)) \\ &= 1 + z d(r, o) + z^2 d(s, o)(d(r, o) - d(s, o)) \\ &\geq 1 + z d(r, o), \end{aligned}$$

and hence the desired result.

$$\text{If } d(r, o) < d(s, o) \text{ then } \frac{1+z d(r, o)}{1+z d(s, o)} < 1 \leq 1 + z d(r, s). \quad \square$$

**Proof of Theorem 1:** If  $Lx = y$  then

$$\sum_{s \in S} L_{rs} \left( \frac{1 + z d(r, o)}{1 + z d(s, o)} \right)^\alpha x_s^z = y_r^z.$$

So

$$(L + \hat{L}^z) x^z = y^z, \quad (2.1)$$

where

$$\hat{L}_{rs}^z = \left[ \left( \frac{1 + z d(r, o)}{1 + z d(s, o)} \right)^\alpha - 1 \right] L_{rs}.$$

If  $z \geq 0$  and  $d(r, o) \geq d(s, o)$ , then applying Lemma 2, we obtain

$$|\hat{L}_{rs}^z| \leq |L_{rs}^z - L_{rs}|. \quad (2.2)$$

If  $d(r, o) < d(s, o)$ , interchanging the roles of  $r$  and  $s$  in Lemma 2 and taking the reciprocal yields

$$\frac{1 + z d(r, o)}{1 + z d(s, o)} \geq \frac{1}{1 + z d(r, s)}.$$

Using the standard inequality  $x + 1/x \geq 2$ , this leads to

$$1 - \left( \frac{1 + z d(r, o)}{1 + z d(s, o)} \right)^\alpha \leq (1 + z d(r, s))^\alpha - 1.$$

Hence (2.2) holds for  $d(r, o) < d(s, o)$  also. Thus

$$\|\hat{L}^z\| \leq \|L^z - L\|.$$

But  $L^z$  depends continuously on  $z$  by Lemma 1, and  $L^0 = L$ , so

$$\|L^z - L\| < \|L^{-1}\|^{-1}$$

for  $z$  near 0. Thus  $(L + \hat{L}^z)$  is invertible for such  $z$  and the theorem follows from (2.1).  $\square$

### 3 Application to continuation problems

As was done in [1] for exponential decay, we now show that under suitable conditions the continuation of a non-degenerate algebraically localised solution of a 1-parameter family of problems remains algebraically localised.

Suppose  $G : \mathcal{R} \times X \rightarrow Y$ ,  $(\varepsilon, x) \mapsto G_\varepsilon(x)$ , is  $C^1$ . We adopt the notation  $D$  for the derivative with respect to  $x \in X$  and  $\partial/\partial\varepsilon$  for the derivative with respect to  $\varepsilon \in \mathcal{R}$ , unless  $\varepsilon$  is the only variable in which case we write  $d/d\varepsilon$ .

Given a solution  $x_0$  of  $G_0(x) = 0$  at which  $DG_0$  is invertible then by the implicit function theorem there is a neighbourhood of  $0 \in \mathcal{R}$  on which  $G_\varepsilon(x) = 0$  has a unique solution  $x(\varepsilon)$  near 0, and this solution satisfies

$$\frac{dx}{d\varepsilon} = -L_\varepsilon^{-1} \frac{\partial G_\varepsilon}{\partial \varepsilon}(x(\varepsilon)) \quad (3.1)$$

where  $L_\varepsilon = DG_\varepsilon(x(\varepsilon))$ . The solution can be continued with respect to  $\varepsilon$  as long as  $L_\varepsilon$  remains invertible.

We suppose a “reference state” in  $X$ , which we shall shift to the origin and hence denote by 0. We say a family of linear operators  $P(\mu)$ ,  $\mu$  in a set  $M$ , is uniformly  $\alpha$ -algebraically local if there exists  $\zeta > 0$  and  $p \in \mathcal{R}$  such that  $\|P^z(\mu)\| \leq p$  for all  $\mu \in M$ .

**Theorem 2** *If  $x(0)$  is  $\alpha$ -algebraically localised about site  $o \in S$  (with respect to reference state 0), with continuation  $x(\varepsilon)$  for  $\varepsilon \in [0, \varepsilon_1]$  with  $L_\varepsilon$  invertible and  $\alpha$ -algebraically local there,  $Q = \frac{\partial G_\varepsilon}{\partial \varepsilon}(0)$  is  $\alpha$ -algebraically localised about  $o$  for  $\varepsilon \in [0, \varepsilon_1]$ ,  $G \in C^2$  and  $P = \frac{\partial}{\partial \varepsilon} DG$  is uniformly  $\alpha$ -algebraically local for  $\varepsilon \in [0, \varepsilon_1]$  and  $\|x\| \leq \sup \{\|x(\varepsilon)\| : 0 \leq \varepsilon \leq \varepsilon_1\}$ , then  $x(\varepsilon)$  is  $\alpha$ -algebraically localised about  $o$  for all  $\varepsilon \in [0, \varepsilon_1]$ .*

**Proof:** Choose  $z > 0$  so that  $\|L^z - L\| < \|L^{-1}\|^{-1}$ . By compactness of  $[0, \varepsilon_1]$  there are common bounds,  $\|Q^z\| \leq q$ ,  $\|L^{-1}\|^{-1} - \|L^z - L\| \geq c^{-1} > 0$ , and by the assumption on  $P$ ,  $\|P^z\| \leq p$ . For  $x \in X$ , let

$$\|x\|_z = \|x^z\|.$$

Then  $\|x(0)\|_z < \infty$  and our aim is to prove that  $\|x(\varepsilon)\|_z < \infty$  for all  $\varepsilon \in [0, \varepsilon_1]$ .

First we show that  $\frac{\partial G_\varepsilon}{\partial \varepsilon}(x)$  is algebraically localised whenever  $x$  is. Now

$$\frac{\partial G_\varepsilon}{\partial \varepsilon}(x) = Q(\varepsilon) + \int_0^1 P(\varepsilon, tx) x \, dt.$$

Using Lemma 2,  $\|Px\|_z \leq \|P^z\| \|x^z\|$ , so  $\left\| \frac{\partial G_\varepsilon}{\partial \varepsilon}(x) \right\|_z \leq q + p \|x\|_z$ .

Next we use (3.1) and Theorem 1 to obtain

$$\left\| \frac{dx}{d\varepsilon} \right\|_z \leq c (q + p \|x\|_z).$$

Integrating this differential inequality from  $x(0)$  we obtain

$$\|x(\varepsilon)\|_z \leq \|x(0)\|_z e^{cp\varepsilon} + (e^{cp\varepsilon} - 1) \frac{q}{p} < \infty,$$

so  $x(\varepsilon)$  remains  $\alpha$ -algebraically localised.  $\square$

Extension to multi-dimensional parameter spaces is easy to achieve by taking one-dimensional paths.

## 4 Breathers in dipole–dipole networks

We recall from [3,4] that in many networks of oscillators, time-periodic solutions of compact support at the uncoupled limit can be continued to a range of weak coupling. The continuation process does not necessarily impose any spatial decay on the resulting solutions. Instead of performing the continuation in a space with some a priori spatial decay properties, we prefer to deduce spatial decay properties afterwards. This is because the continuation process often works equally well starting from solutions of the uncoupled network with no spatial decay properties, and yields interesting solutions like multi-breathers. But if the solution at the uncoupled limit is spatially localised then we can often deduce spatial decay of the continuation. For example, exponential decay was obtained in [3] for the continuation of solutions of compact support for 1D nearest neighbour chains and results were quoted there extending this conclusion to a variety of situations. Subsequently, using the idea of [1], this result was generalised in [4] to all networks with exponentially decaying interaction.

Here we apply Theorem 2 to prove algebraic decay in space for time-periodic solutions of some dipole–dipole networks obtained by continuation of solutions of compact support at the uncoupled limit.

Suppose  $S$  is a  $\mathbb{Z}^D$ -lattice with Euclidean metric, and suppose interaction potential

$$\varepsilon W(x) = \sum_{r \neq s} \frac{\varepsilon (x_r - x_s)^2}{2d(r, s)^K} \quad (4.1)$$

and let

$$U_\varepsilon(x) = \sum_r V(x_r) + \varepsilon W(x) \quad (4.2)$$

for some local potential  $V$ .

For simplicity, instead of breathers we begin by considering the problem of continuing equilibria of  $U_\varepsilon$  from  $\varepsilon = 0$ , for  $V$  with (at least) two non-degenerate critical points, without loss of generality at  $x = 0$  and  $1$ . So we consider the implicit function problem

$$G_\varepsilon(x) := DU_\varepsilon(x) = 0.$$

Then  $L = D^2W$  has off-diagonal elements

$$L_{rs} = \varepsilon d(r, s)^{-K}.$$

Use the  $\ell_p$  metric, any  $p \in [1, \infty]$ , for the spaces  $X$  and  $Y$ . Then  $L$  is bounded iff  $K > D$  (independently of  $p$ ). We will assume this inequality, else the energy due to displacing a single site is infinite. The modified operator  $L^z$  is bounded iff  $\alpha < K - D$ . Thus we see that  $L$  is algebraically local if  $K > D$ , and we can choose any  $\alpha \in [0, K - D)$  and any  $z > 0$ . For interacting dipoles ( $K = 3$ ) on a surface ( $D = 2$ ) we can take any  $\alpha \in [0, 1]$ , for example  $\alpha = 0.99$ .

Applying Theorem 2 to an initial equilibrium with  $x_o = 1, x_r = 0$  for  $r \neq o$ , we obtain  $\|x(\varepsilon)\|_z < \infty$ . For  $p = \infty$  this says that

$$|x_r(\varepsilon)| \leq \frac{C}{(1 + z d(r, o))^\alpha},$$

which is an algebraic decay result, though not as fast as for the coupling. Stronger results, however, are obtained with  $p < \infty$ . In particular,  $p = 1$  gives

$$\sum_r |x_r(\varepsilon)| (1 + z d(r, o))^\alpha < \infty. \quad (4.3)$$

If  $x(\varepsilon)$  is “full”, in the sense that  $|x_r|$  lies between two positive constants times a decreasing function of distance, then this implies that  $|x_r| \leq C d(r, o)^{-K}$ , but our method does not guarantee that  $x(\varepsilon)$  is full so in general (4.3) is the strongest bound that we obtain. In the next section we give examples where the true solution has  $|x_r| \sim C d(r, o)^{-K}$ . Thus our use of Lemma 2 must be throwing away a lot in the cases  $p > 1$ . This merits further investigation.

To prove algebraic localisation of breathers for a network with potential (4.2), instead of having two critical points we just require  $V$  to have a non-degenerate local minimum, without loss of generality at 0. We study the implicit function problem  $G_{\varepsilon, T}(x) = 0$ , where  $G_{\varepsilon, T} : C_p^2(S_T^1, \mathcal{R}^S) \rightarrow C_p^0(S_T^1, \mathcal{R}^S)$  is defined by

$$[G_{\varepsilon, T}(x)]_r(t) = \ddot{x}_r(t) + \frac{\partial U_\varepsilon}{\partial x_r}(x(t)),$$

and  $C_p^k(S_T^1, \mathcal{R}^S)$  is the space of  $C^k$  functions from a circle  $S_T^1$  of length  $T$  to  $\mathcal{R}^S$ , with the norm

$$\|x\|^p = \sum_{s \in S} |x_s|_k^p,$$

for  $1 \leq p < \infty$ , or  $\|x\| = \sup_{s \in S} |x_s|_k$  for  $p = \infty$ , where  $|\cdot|_k$  is the  $C^k$ -norm, and then we will apply Theorem 2 (cf. [4]).

It can be checked that  $L := DG_{\varepsilon, T, x}$  is bounded if  $K > D$  and is  $\alpha$ -algebraically local for all  $\alpha < K - D$ . If  $x(0)$  is a breather for  $\varepsilon = 0$  with only one site  $s = o$  excited and  $L$  is invertible there then by the implicit function theorem we obtain a breather  $x(\varepsilon)$  for  $\varepsilon$  small. Using Theorem 2, for all  $\varepsilon$  to which the breather can be continued with  $L$  invertible and  $z$  small enough, we deduce that  $\|x(\varepsilon)\|_z < \infty$ . Hence  $x(\varepsilon)$  is  $\alpha$ -algebraically localised about site  $o$ .

Again the strongest result is obtained for  $p = 1$ , though approximate analysis below suggests that it is not quite optimal.

The same analysis applies to problems with algebraically decaying interaction of more than just two body form.

## 5 True decay

For networks with algebraically decaying interaction, our theorems give upper bounds on the spatial decay of linear responses and of solutions of nonlinear problems obtained by continuation from localised solutions. For comparison, we compute here the asymptotics of the decay for some simple 1D problems.

Let

$$L_\varepsilon = I - \varepsilon \Delta : \mathcal{R}^{\mathbb{Z}} \rightarrow \mathcal{R}^{\mathbb{Z}},$$

where

$$\Delta_{rs} = |r - s|^{-K} \quad \text{for } r \neq s, \quad \Delta_{ss} = 0,$$

for some  $K \in \mathcal{N}$ , and use  $\ell_1$  norm. The operator  $\Delta$  is bounded if  $K > 1$ , so  $L_\varepsilon$  is  $\alpha$ -algebraically local for all  $\alpha < K - 1$  and  $L_\varepsilon$  is invertible for  $\varepsilon < \|\Delta\|_1^{-1}$ . Given  $y \in \ell_1$  and  $k \in S_{2\pi}^1$ , let

$$\hat{y}(k) = \frac{1}{2\pi} \sum_r y_r e^{-ikr},$$

so

$$y_r = \int_{-\pi}^{\pi} \hat{y}(k) e^{ikr} dk.$$

Then the solution  $x \in \ell_1$  of  $L_\varepsilon x = y$  is given by

$$\hat{x}(k) = \frac{\hat{y}(k)}{\lambda(k, \varepsilon)},$$

where

$$\begin{aligned} \lambda(k, \varepsilon) &= 1 - \varepsilon \sum_{n \neq 0} |n|^{-K} e^{ikn} \\ &= 1 - 2\varepsilon g_K(k), \end{aligned}$$

where

$$g_K(k) = \sum_{n \geq 1} n^{-K} \cos nk.$$

For example, for  $K = 2$ , we have

$$g_2(k) = \frac{\pi^2}{6} + \frac{k^2}{4} - \frac{\pi k}{2} \quad \text{for } 0 \leq k \leq 2\pi$$

and then repeated periodically. In particular,  $g_K$  is  $C^{K-2}$  and piecewise  $C^{K-1}$  with jumps in the  $(K - 1)$ st derivative.

Now suppose  $y_0 = 0$ ,  $y_r = |r|^{-P}$  for  $r \neq 0$ , some  $P \in \mathcal{N}$ , then  $y \in \ell_1$  for  $P > 1$  and is  $\alpha$ -algebraically localised about  $r = 0$  for  $\alpha < P - 1$ . The Fourier transform

$$\hat{y}(k) = \frac{1}{\pi} g_P(k).$$

Hence

$$\hat{x}(k) = \frac{1}{\pi} \frac{g_P(k)}{(1 - 2\varepsilon g_K(k))}.$$

For  $\varepsilon \neq 0$ , this function is  $C^{N-2}$  and piecewise  $C^{N-1}$  with jumps in the  $(N-1)$ st derivative, where  $N = \min(K, P)$ . Thus by Fourier analysis (e.g. [5]), there exists  $C \neq 0$  such that  $|x_r| \sim C|r|^{-N}$ .

Compare our bound

$$\sum_{r \in \mathcal{Z}} (1 + z|r|)^\alpha |x_r| \leq C'(\alpha, z) \quad \text{for all } \alpha < N - 1,$$

which is slightly weaker.

Next we consider continuation of equilibria for the 1D case of the example (4.2). The equilibria are given by

$$V'(x_r) + \varepsilon \sum_{s \neq r} |r - s|^{-K} (x_r - x_s) = 0.$$

We know that the continuation of the initial solution  $x_0 = 1, x_r = 0 (r \neq 0)$  is in  $\ell_1$  for  $K > 1$ , thus  $|x_r| \rightarrow 0$  as  $|r| \rightarrow \infty$ , and so it makes sense to separate the linearised part of the equation about 0 from the remainder:

$$\omega_0^2 x_r + \varepsilon \sum_{s \neq r} |r - s|^{-K} (x_r - x_s) = y_r := \omega_0^2 x_r - V'(x_r) = O(x_r^2),$$

where  $\omega_0^2 = V''(0)$ . It follows that

$$\hat{x}(k) = \frac{\hat{y}(k)}{\omega_0^2 + 2\varepsilon (g_K(k) - g_K(0))}.$$

Now  $y_r = O(x_r^2)$  implies that  $y_r$  decays more rapidly than  $x_r$  with  $r$  and hence that  $\hat{y}$  is strictly smoother than  $\hat{x}$  (a little more work is required to give a rigorous statement here). Hence we deduce that the smoothness of  $\hat{x}$  is that of the denominator, i.e.  $\hat{x}$  is  $C^{K-2}$  and piecewise  $C^{K-1}$  with jumps in the  $(K-1)$ st derivative. It follows that there exists  $C(\varepsilon) \neq 0$  such that

$$x_r \sim C(\varepsilon) |r|^{-K}.$$

The same sort of argument can be attempted to find the asymptotics of the spatial decay for breathers in this model. We recall that in the case of finite range interactions in a translation invariant lattice, the decay in space of each time-Fourier component of a breather is asymptotically exponential with exponent given in the simplest cases by the smallest imaginary part of the roots  $k$  of the continuation of the dispersion relation for

linearised solutions about equilibrium to complex wave number [6] (strictly speaking, the analysis given there needs extending to estimate the effect of the nonlinear remainders, including interaction of different time-Fourier harmonics, and this effect can lead to different results for the spatial decay of some of the harmonics, as was realised in [7]; a rigorous analysis can be obtained by bounding the spatial Fourier transform of the  $m^{\text{th}}$  harmonic in a strip  $|\text{Im } k| \leq \delta_m$ ).

In the case of algebraically decaying interactions, the dispersion relation does not have an analytic continuation to complex wave numbers, and in any case the problem always reduces to a linear inhomogeneous one rather than linear homogeneous, because of the infinite range of the interaction. We can write the equations of motion as

$$\ddot{x}_r + \omega_0^2 x_r + \varepsilon \sum_{s \neq r} |r - s|^{-K} (x_r - x_s) = y_r := \omega_0^2 x_r - V'(x_r) = O(x_r^2).$$

If the breather has frequency  $\omega_b$ , its time-Fourier harmonics separate on the left but are mixed on the right. We obtain for the  $m^{\text{th}}$  time-Fourier component  $x^m$  of the breather

$$\hat{x}^m(k) = \frac{\hat{y}^m(k)}{\omega_0^2 - m^2 \omega_b^2 + 2\varepsilon (g_K(k) - g_K(0))}.$$

It is not easy to make a rigorous argument in this case, but it is plausible that  $y^m$  decays in space faster than  $x^m$ , at least for the fundamental ( $m = 1$ ), and hence the smoothness of  $\hat{x}^m$  would be given by the denominator again, so the oscillation amplitude would decay in space like  $C|r|^{-K}$ .

## References

- [1] Baesens, C., MacKay, R.S., *Exponential localization of linear response in networks with exponentially decaying coupling*, Nonlinearity **10** (1997) 931–940.
- [2] Jakob, P., *Dynamics of the C – O stretch overtone vibration of CO/Ru(001)*, Phys. Rev. Lett. **77** (1996) 4229–32.
- [3] MacKay, R.S., Aubry, S., *Proof of existence of breathers for time-reversible or Hamiltonian networks of weakly coupled oscillators*, Nonlinearity **7** (1994) 1623–43.
- [4] Sepulchre, J.-A., MacKay, R.S., *Localised oscillations in conservative and dissipative networks of weakly coupled autonomous oscillators*, Nonlinearity **10** (1997) 679–713.
- [5] Zygmund, A., *Trigonometric series* (Cambridge University Press, 1959).
- [6] Flach, S., *Conditions on the existence of localized excitations in nonlinear discrete systems*, Phys. Rev. E **50** (1994) 3134–42.
- [7] Flach, S., *Obtaining breathers in nonlinear Hamiltonian lattices*, Phys. Rev. E **51** (1995) 3579–87.