Zeitschrift:	Schweizer Archiv für Tierheilkunde SAT : die Fachzeitschrift für Tierärztinnen und Tierärzte = Archives Suisses de Médecine Vétérinaire ASMV : la revue professionnelle des vétérinaires
Herausgeber:	Gesellschaft Schweizer Tierärztinnen und Tierärzte
Band:	132 (1990)
Heft:	8
Artikel:	Cytokeratin filaments of the liver of BALB/C-Mice as a sensitive model for liver damages : computer-aided characterization with the help of the image analysing system "IBAS"
Autor:	Steinmann, J. / Kohler, M. / Aufderheide, M.
DOI:	https://doi.org/10.5169/seals-593705

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 01.07.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch



Fig. 1: Granulomatous reaction with the presence of multinucleated giant cells (arrows). Hematoxylin and eosin, 160x.

ted in the cytoplasm of degenerating tubular cells. In few cases leptospiral antigen was detected in the cytoplasm of giant cells. Moreover, these cells showed a variable positive reaction when tested with the monoclonal antibody against vimentin (fig. 2). In 4 out of 12 cases isolation attempts for leptospires were carried out. All the 4 cases resulted positive and the isolates were belonging to the *pomona* serogroup.

The presence of multinucleated giant cells in renal lesions strikingly suggestive of granulomata has been described in cattle naturally or experimentally infected with serovar *pomona* (1, 2). These giant cells were considered to develop from tubular epithelium. However, in our



Fig. 2: Intense immunoreactivity for vimentin is detectable in the cytoplasm of two giant cells (arrows). Immunoperoxidase for vimentin, hematoxylin counterstain, 200x.

cases some findings suggest the monocytic-macrophagic origin of these cells. In fact their cytoplasm was PAS positive, reacted positively to vimentin immunolabelling and they were frequently observed adjoining to degenerating tubules. Therefore, it may be suggested that the granulomatous-like reaction might be directed against tubular structures containing leptospiral antigen.

References

1. Hadlow W. J., Stoenner H. G. (1955): Am. J. Vet. Res. 16, 45–56. — 2. Spradbrow P. B., Seawright A. A. (1963): Austral. Vet. J. 39, 423–433.

Hannover Medical School, Inst. Experimental Pathology, FRG

CYTOKERATIN FILAMENTS OF THE LIVER OF BALB/C-MICE AS A SENSITIVE MODEL FOR LIVER DAMAGES. COMPUTER-AIDED CHARACTERIZATION WITH THE HELP OF THE IMAGE ANALYSING SYSTEM «IBAS»

J. Steinmann, M. Kohler, M. Aufderheide, U. Mohr The cytoskeleton represents a complex structural network which is of major importance both for the structural integrity and the physiologic processes of the cell (1; 2; 3). Damages of the cytokeratin filament organisation and dysregulation of the keratin metabolism are regarded as responsible mechanisms leading to alteration of cell function, cell death or transformation (4; 5).

The aim of the present investigation was to test whether liver-damaging influences can be detected through changes in the liver cytokeratin filaments as one important component of the liver cytoskeleton. The iron storage behaviour of the liver was investigated as a comparative marker for induced liver alterations (data not shown).

Material and methods

In the animal experiment 10 male and 10 female mice per group were treated according to UCHIDA and HIRONO (14) with dimethyl-

nitrosamine (DMNA/Phenobarbital, group II) or colchizine (group III), respectively. The control group (group I) remained untreated. The histologic changes in different locations of the liver tissue were characterized by light and fluorescence microscopy (primary antibody AB: Antikeratin Bio Yeda, secondary AB: AMCA-Antiguinea pig) and keratin alterations were quantified with the image analysing system «IBAS« (Zeiss). After image-preparing steps, areas without tissue components were eliminated to get the reference area. Subdividing the grey value spectrum of the remaining picutre into phases with grey values ranging from 0–90, 91–150 and 151–255 it was possible to discriminate weakly, moderately and intensively fluorescing keratin components and to measure their corresponding area. The area percentage of the reference area and the areas with low (LKF_{Ind}), moderate MKF_{Ind}) and high fluorescence intensity (HKF_{Ind}) were defined as keratin indices of the corresponding fluorescence

intensity. For statistical evaluation an ANOVA-test with the SASsystem (software release No 148270010) was carried out.

Results

Group II developed hepatocellular adenomas with an overall tumor incidence of 55% (11/20). 36% of the tumor bearing animals showed multiple adenomas. The colchizine-treated group showed some intranuclear inclusion bodies. Investigation of the liver cytokeratin filaments in the DMNA group by indirect immunofluorescence showed a marked tendency towards keratinization of the liver cells with different morphologic features. This strong keratinization did not indicate any sex-dependent behavior and could be measured quantitatively in the form of a highly significant increase of the parameter HKF_{Ind} compared to the control (p < 0.001). The colchizine group did not display any structural changes compared with the control group, but image analysis of the keratin filaments revealed that the males of the colchizine group had significantly greater values for the parameter LKF ind whereas the values for the parameter HKF Ind, were significantly smaller than for the control males (p <0.001). No group showed lobe-specific differences concerning the induced keratin alterations (see figure 1).

Discussion

In the liver cells of the DMNA-treated group a general condensation of the cytoplasmic keratin network could be observed. It was partly accompanied by both non-filamentous keratin conglomerates which were in close contact to the cytoplasmic network and with a general irregular thickening of the pericytoplasmic keratin bands with a clod-like morphology.

Thus the classical concept of Mallory body (MB) formation, the best known pathologic keratin alteration in liver cells, (6; 7; 8; 9; 11; 12; 13) has to be extended in as far that MB formation can also be a combination of keratin conglomeration which leads to polymorphic non-filamentous structures and a condensation of the cytoplamic

Institute of Pathology, Hannover Veterinary School, FRG

MORPHOLOGICAL CHARACTERISATION OF BOVINE MESOTHELIOMA

N. Stockhofe-Zurwieden, M. Stöber, J. Pohlenz

Mesotheliomas are rare tumors in the bovine species with an incidence of about 1.6% (Klopfer et al., 1983). Generally they are described as diffuse tumorous proliferations on the peritoneal surface but they also occur on the pleural and pericardial serosa.

Material and methods

In this retrospective study 13 cases of bovine mesotheliomas of the last 16 years were re-evaluated. 10 of the 13 cases were necropsied, from 3 cases only biopsy specimens were available. The material had been fixed in 10% formalin. Paraffin-embedded tissues stained with hematoxylin-eosin were examined by light microscopy. For additional histochemical studies tissue sections were stained with Periodicacid-Schiff (PAS) with and without diastase digestion, alcian-blue and Meyers mucicarmine. To detect ferruginous bodies sections



Fig. 1: Area percentage (%) of the different keratin indices of both sexes of the control, DMNA and colchizine groups

keratin network. Besides this keratinization, a general loss of keratin components can be a further expression of toxic damages of liver cells, as observed after cholchizine treatment. Liver cytokeratin filaments could be morphologically and quantitatively characterized as sensitive indicators for induced liver damages.

References

Fey E. G. et al. (1884): J. Cell Biol. 98, 1973–1984. — 2. Fey E.
G., Penman S. (1985): Hepatology 5, (1), 152–158. — 3. Gerace L.
(1986): Trends biochem. 11, 443–446. — 4. Borenfreund E. et al.
(1981): Hepatology 1, (5), 408–415. — 5. Mukunda B. R. (1987): Human Pathol. 18, (1). — 6. Christoffersen P. (1972): Acta path. microbiol. scand. Section A 80. — 7. Denk H. et al. (1979): Proc. Natl. Acad. Scie. USA 76, (8), 4112–4116. — 8. Denk H. et al.
(1981): Hepatology 9–20. — 9. Denk H., Franke W. W. (1981): Europ. J. Cell Biol. 23,241–249. — 10. Kimoff R. J., Huang S. N.
(1981): Lab. Invest. 45. — 11. French S. W. (1983): Pathol. Lab. Med. 107. — 12. Okanoue T.M. et al. (1985): Lab. Invest. 53, (5), 534–540. — 13. Mukunda B. R. (1987): Human Pathol. 18, (1). — 14. Uchida E. Hirono E. (1979): Gann. 70, 639–644.

were stained with Turnbull-blue. Electronmicroscopical investigations were carried out in two cases on glutaraldehyde (5%) fixed tissue.

Results

Out of the 13 cases 5 animals were younger than 10 weeks, 3 animals between 4 months and 1 year and 5 animals were older than 5 years. The latter group consisted of cows only, whereas the first two goups included 5 males and 1 heifer.

The peritoneal cavity was predominantly affected. The tumor spread over the visceral and parietal surface of the serosa. The scrotal cavity was involved in 4 bull calves, whereas in 6 cases the pleural, pericardial and peritoneal serosa contained tumors. Macroscopically the tumors revealed a diffuse distribution pattern with some of them having a pedunculated growth and others being sessile on the surface.