

Atrichops crassipes (Meigen, 1820) (Diptera, Athericidae), a species not so new for Switzerland

Autor(en): **Sartori, Michel / Lods-Crozet, Brigitte / Derleth Sartori, Pascale**

Objektyp: **Article**

Zeitschrift: **Mitteilungen der Schweizerischen Entomologischen Gesellschaft = Bulletin de la Société Entomologique Suisse = Journal of the Swiss Entomological Society**

Band (Jahr): **84 (2011)**

Heft 3-4

PDF erstellt am: **19.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-403029>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Atrichops crassipes (Meigen, 1820) (Diptera, Athericidae), a species not so new for Switzerland

MICHEL SARTORI¹, BRIGITTE LODS-CROZET^{1,2}, PASCALE DERLETH
SARTORI^{1,2}, JEAN PERFETTA³, PASCAL STUCKI⁴ & HEINRICH VICENTINI⁵

¹ Musée cantonal de zoologie, Palais de Rumine, Place Riponne 6, CH-1005 Lausanne;
michel.sartori@vd.ch

² Laboratoire du Service des Eaux, Sols et Assainissement (SESA), Ch. des Boveresses 155, CH-1066
Epalinges

³ Service de l'écologie de l'eau (SECOE-DIM), case postale 78, CH-1211 Genève 8

⁴ Aquabug, case postale 1643, CH-2001 Neuchâtel

⁵ Gewässerökologie, Wasserschöpfli 48, CH-8055 Zürich

The genus *Atrichops* Verrall, 1909, with the species *A. crassipes* (Meigen, 1820) is documented for the first time in Switzerland, although its first mention dates back to 1995, but remained unnoticed.

Keywords: new record, Switzerland, aquatic insect, Diptera.

INTRODUCTION

Athericidae (watersnipe-flies) is a small family of primitive Brachycera related to Tabanidae and Rhagionidae (Stuckenberg 1973; Rozkosny & Nagatomi 1997). The larvae are always aquatic and live in fast flowing streams or in more lentic habitats (Thomas 1981, 1997). They are top predators and feed on other invertebrates such as Diptera (Chironomidae), Ephemeroptera or Plecoptera (Thomas 1975).

Up to now, only two species, out of the 10 known to occur in Europe, have been reported from Switzerland (Merz 1998): *Atherix ibis* (Fabricius, 1798) and *Ibis-sia marginata* (Fabricius, 1781) which are both relatively common in stony streams. A third one, *Atrichops crassipes* (Meigen, 1820) has been mentioned by Merz (1998) as a potential candidate to live in Switzerland. In fact, we discovered during the writing of this note, that it has been already cited by Oertli (1995) in a species list of macroinvertebrates living in a pond near Geneva. This citation remained unnoticed up to now, although published in an entomological journal. Here we document further on the presence of this species in several Swiss streams and rivers, its most common habitat.

Atrichops crassipes (Meigen, 1820)

Material:

Geneva: Seymaz, embouchure, 502970/115070, 392 m, 1 larva; Claparede, 503220/116000, 403 m, 3 larvae; De Haller, 504095/117390, 417 m, 2 larvae, II. 2007. **Nant du Paradis**, embouchure, 507215/12065, 428 m, 3 larvae, II. 2007. **Hermance**, Pont de Bouringe, 508225/128395, 380 m, 1 larva, XI. 2007; Pont de

Crevy, 507695/126490, 415 m, 2 larvae, V. 2007. Marnot, embouchure, 508600/127400, 395 m, 1 larva, V. 2007. **Rhone**, Touvière, 488100/114740, 347 m, 1 larva, XII. 2008. **Arve**, Ecole-de-Médecine, 499440/116815, 375 m, 1 larva, V. 2008. All coll. SECOE.

Vaud: **Venoge**, Bussigny/Lausanne, 531710/ 155175, 375 m, 3 larvae; Vufflens-la-Ville, 530415/ 158980, 405 m, 2 larvae; Denges, 531505/152915, 380 m, 3 larvae, all coll. 16.II.2010 SESA. Denges, 1 larva, 16.IV.2011, coll. P. Derleth.

Jura: **Vendline**, Beurnevésin aval, frontière, 576345/261000, 419 m, 2 larvae, 31.III.2011, coll. P. Stucki.

Lucerne: **Suhre**, Oberkirch, 663459/225401, 506 m, 1 larva, 25. IV. 2007, 3 larvae, 28. VIII. 2007, 2 larvae, 16. IV. 2003, 3 larvae, 14. VII. 2003; Büron, 648740/228840, 487 m, 1 larva, 15. IV. 2003, coll. H. Vicentini. **Aabach**, Hitzkirch, 661390/230220, 473 m, 2 larvae, 15. IV. 2003, coll. V. Lubini.

A. crassipes is known from Western Europe, extending its range up to Poland and Romania (Roskosny 2011). It has been recently mentioned from Hungary for the first time (Muranyi *et al.* 2009). The larvae are psammophilous, living in sand and fine substrate deposited by the current in streams and rivers, where they feed largely on Chironomidae (Thomas 1975). They also colonize the deposits of some dam-lakes (Gagneur 1981) or ponds (Oertli 1995). The morphology of the larvae is very characteristic and cannot be confused with any other species, even among watersnipe-flies. Abdominal segments 6 and 7 bear each four tracheal projections, (two dorsal and two lateral), and the anal segment two more. They are oriented rearwards and upwards to the surface of sediments (Fig. 1).

This merolimnic species goes through five aquatic larval instars before pupation in the banks, and the single available study reports a semivoltine cycle, i.e. development lasts two years (Gerke & Böttger 2001). At the moment, no adults have been caught but according to Thomas (1981) adults fly between June and July. As for other Athericidae, females lay their eggs in batches on the underside of leaves above the watercourse (Gerke & Böttger 2001), and first instar larvae drop into the water shortly after hatching.

In the Canton de Vaud, no specimens have been caught before 2010. Up to now, the Venoge River is the only watercourse to be colonized by *A. crassipes*. At the mentioned localities, the river is between 8 and 10 m wide, the current speed is around 0.2 and 0.4 m/s. Substrate composition and bank vegetation strongly vary between localities, suggesting that the availability of adequate fine substrate is the main parameter for its setting up.

It has been reported that *A. crassipes* belongs to species on which global climate change has a positive impact (Daufresne *et al.* 2003). This could explain the actual scarcity of data for this species in Switzerland, its low altitudinal range (all localities below 510 m), and the fact that it could be a new immigrant, as documented for instance for the mayfly *Habrophlebia eldae* (Wagner *et al.* 2007).

ACKNOWLEDGEMENTS

Our sincere thanks to Alain Thomas (Toulouse) for his comments and to Verena Lubini (Zurich) for sharing her data.




Fig. 1 : *Atrichops crassipes* (Meigen, 1820), larva. A: general habitus, B: Larva digging in the sediment, C: close view of the mouthparts, especially the mandibles.

REFERENCES

- Daufresne, M., Roger, M.C., Capra, H. & Lamouroux, N. 2003. Long-term changes within the invertebrate and fish communities of the Upper Rhône River: effects of climatic factors. — *Global Change Biology* 10: 124-140.
- Gagneur, J. 1981. Sur la biologie d'*Atrichops crassipes* Meigen (Diptères, Athericidae) dans le réservoir de Cajarc (Lot). — *Bulletin de la Société d'Histoire naturelle de Toulouse* 117: 231-237.
- Gerke, N. & Böttger, K. 2001. The life cycle of *Atrichops crassipes* Meigen, 1820 (Diptera: Athericidae) at the lower Schierenseebrook, a lake outflow in the North German Lowland. — *Aquatic Insects* 23: 85-92.
- Merz, B. 1998. Athericidae. In: Merz, B., Bächli, G., Haenni, J.-P. & Gonseth, Y. (eds), *Diptera – Checklist*, p. 149. — *Fauna Helvetica* 1, CSCF-SEG, Neuchâtel.
- Muranyi, D., Tarjanyi, N. & Schöll, K. 2009. First record of the genus *Atrichops* Verrall, 1909 in Hungary (Diptera: Athericidae). — *Opuscula Zoologica, Budapest* 40: 103-105.
- Oertli, B. 1995. Impact de la création d'un biotope sur la conservation de la biodiversité : les macroinvertébrés de l'étang du Bois-Vieux (Genève). — *Bulletin romand d'Entomologie* 13: 11-24.
- Rozkosny, R. 2011. Fauna Europaea: Athericidae. In: Papp, T. & Beuk, P. (eds), *Fauna Europaea: Diptera, Brachycera. Fauna Europaea version 2.4*, <http://www.faunaeur.org>
- Rozkosny, R. & Nagatomi, A. 1997. Family Athericidae. In: Papp, L. & Darvas, B. (eds), *Contributions to a manual of Palaearctic Diptera*, volume 2, pp. 439-446. — *Science Herald, Budapest*.
- Stuckenberg, B.R. 1973. The Athericidae, a new family in the lower Brachycera (Diptera). — *Annals of the Natal Museum* 21: 649-673.
- Thomas, A.G.B. 1975. Diptères torrenticoles peu connus: III.- Les Athericidae du sud de la France (Régime alimentaire des larves: aspect qualitatif) (Brachycera, Orthorrhapha). — *Annales de Limnologie* 11: 169-188.
- Thomas, A.G.B. 1981. Travaux sur la taxonomie, la biologie et l'écologie d'insectes torrenticoles du Sud-ouest de la France (Ephéméroptères et Diptères: Dixidae, Cecidomyiidae, Rhagionidae et Athericidae), avec quelques exemples de perturbations par l'homme. — Thèse de doctorat, n° 988, Université Paul Sabatier, Toulouse, 330 pp.
- Thomas, A.G.B. 1997. Rhagionidae and Athericidae, Snipe-flies. In: Nilsson, A.N. (ed.), *Aquatic Insects of North Europe*, volume 2, pp. 311-320. — *Apollo Books, Stenstrup*.
- Wagner, A., Lubini, V. & Vicentini, H. 2007. *Habrophlebia eldae* Jacob & Sartori, 1984 (Ephemeroptera: Leptophlebiidae) neu für die Schweizer Fauna. — *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 80: 91-98.

(received September 15, 2011, accepted November 7, 2011)